

TABLE OF CONTENTS

INTRODUCTION

The Northern Global Change Research Program
Richard A. Birdsey, John L. Hom, and Marla Emery..... 1

MONITORING AND PREDICTING REGIONAL ENVIRONMENTAL CHANGE

Delineation of Climate Regions in the Northeastern United States
Arthur T. DeGaetano..... 7

Enhancement of Regional Wet Deposition Estimates Based on Modeled Precipitation Inputs
James A. Lynch, Jeffrey W. Grimm, and Edward S. Corbett..... 8

RESPONSES OF NORTHERN TREE SPECIES TO REGIONAL STRESS

Effects of Calcium Fertilization and Acid Mist on Calcium Concentration and Cold Tolerance of
Red Spruce Needles
G. R. Strimbeck, David R. Vann, and Arthur H. Johnson..... 10

Ozone-Induced Accelerated Foliar Senescence: Implications for Toxicity and Compensation
Eva J. Pell, Bryan W. Brendley, and Judith P. Sinn..... 13

Root Growth and Respiration of Aspen in Response to Ozone and Elevated Carbon Dioxide
M. D. Coleman, R. E. Dickson, and J. G. Isebrands..... 20

Effects of Ozone and CO₂ on the Growth and Physiology of Aspen
D. F. Karnosky, Z. Gagnon, R. E. Dickson, P. Pechter, M. Coleman, Olevi Kull,
Anu Sober, and J. G. Isebrands..... 21

Interacting Effects of Ozone and CO₂ on Growth and Physiological Processes in Northern Forest
Trees
J. G. Isebrands and D. F. Karnosky..... 22

The Chronic Response of Yellow-Poplar and Eastern White Pine to Ozone and Elevated Carbon
Dioxide: Three-Year Summary
Joanne Rebbeck..... 23

Effects of Carbon Dioxide Enrichment on Response of Pitch Pine Grown at Different Nutrient
Levels to Aluminum
George A. Schier and Carolyn J. McQuattie..... 31

Elevated CO₂ Compensates for Water Stress in Northern Red Oak
Patricia T. Tomlinson and Paul D. Anderson..... 32

Modeling Impacts of CO₂, Ozone, and Climate Change on Tree Growth
George E. Host, Gary W. Theseira, and J. G. Isebrands..... 33

Ozone Exposure, Uptake, and Response of Different-Sized Black Cherry Trees
Todd S. Fredericksen, John M. Skelly, Kim C. Steiner, and Thomas E. Kolb..... 38

Water Use in Forest Canopy Black Cherry Trees and Its Relationship to Leaf Gas Exchange and Environment B. J. Joyce, K. C. Steiner, and J. M. Skelly.....	41
Effects of Elevated CO ₂ and Ozone on Phenolic Glycosides of Trembling Aspen James K. Nitao, Muraleedharan G. Nair, William J. Mattson, Daniel A. Herms, Bruce A. Birr, Mark D. Coleman, Terry M. Trier, and J.G. Isebrands.....	42
Variable Performance of Outbreak Defoliators on Aspen Clones Exposed to Elevated CO ₂ and O ₃ Daniel A. Herms, William J. Mattson, David N. Karowe, Mark D. Coleman, Terry M. Trier, Bruce A. Birr, and J. G. Isebrands.....	43
Genetic Aftereffects of Increased Temperature in <i>Larix</i> Michael S. Greenwood and Keith W. Hutchison.....	56
Physiological and Genotypic Responses of Central Hardwood Species to Allelochemicals, Other Stresses, and Their Interactions J. W. Van Sambeek, John E. Preece, Nadia E. Navarrete, and George Rink.....	63
RESPONSES OF ECOSYSTEM PROCESSES TO REGIONAL STRESS	
The Howland Integrated Forest Study (HIFS) - Ecosystem Research on Atmospheric Influences Governing Forest Function Ivan J. Fernandez, Lindsey E. Rustad, Jeffrey A. Simmons, and James W. McLaughlin.....	67
Biogeochemical Cycling of Carbon, Nitrogen, and Sulfur at the Howland Integrated Forest Study Site, Howland, Maine James W. McLaughlin, Ivan J. Fernandez, Stewart M. Goltz, Lindsey E. Rustad, and Larry Zibilske.....	68
Elemental Cycling Response of an Adirondack Subalpine Spruce-Fir Forest to Atmospheric and Environmental Change Andrew J. Friedland and Eric K. Miller.....	77
Internal Uptake and Assimilation of Gaseous Nitric Acid by Western Forest Tree Species John L. Hom, Marek Krywult, Andrzej Bytnerowicz, and Kevin E. Percy.....	82
Climatic and Pollution Influences on Ecosystem Processes in Northern Hardwood Forests Kurt S. Pregitzer, David D. Reed, Glenn D. Mroz, Andrew J. Burton, John A. Witter, and Donald A. Zak.....	83
Effects of Elevated CO ₂ and Shade on the Decomposition of Senesced Tree Foliage: Impacts on Microbial Activity Michael G. Kaufman, R. Malcolm Strand, Mark E. Kubiske, William J. Mattson, Daniel A. Herms, Edward D. Walker, Kurt S. Pregitzer, and Richard W. Merritt.....	89
Effects of Elevated CO ₂ and Shade on the Decomposition of Senesced Tree Foliage: Impacts on the Growth and Survival of Treehole Mosquitoes R. Malcolm Strand, Daniel A. Herms, Michael G. Kaufman, Mark E. Kubiske, William J. Mattson, Edward D. Walker, Kurt S. Pregitzer, and Richard W. Merritt.....	101
Cation Depletion in New England Forests Richard Hallett and James Hornbeck.....	111

Aluminum Mobilization and Calcium Depletion in the Forest Floor of Red Spruce Forests in the Northeastern United States Gregory B. Lawrence, Mark B. David and Walter C. Shortle.....	112
Calcium Status of the Forest Floor in Red Spruce Forests of the Northeastern U.S. - Past, Present and Future Mark B. David, Gregory B. Lawrence, Walter C. Shortle, and Scott W. Bailey.....	118
Putrescine: A Marker of Stress in Red Spruce Trees Rakesh Minocha, Walter C. Shortle, Gregory B. Lawrence, Mark B. David, and Subhash C. Minocha.....	119
Long-term Changes in the Acidity of a DeKalb Forest Soil in the Mid-Region of the Susquehanna River Watershed Joy R. Robert and William E. Sharpe.....	131
Experimental Soil Warming Effect on C, N, and Major Element Cycling in a Low Elevation Spruce-Fir Forest Soil Lindsey E. Rustad, Ivan J. Fernandez, and Stephanie Arnold.....	132
Soil Respiration and Net N Mineralization Along a Climate Gradient in Maine Jeffrey A. Simmons, Ivan J. Fernandez, and Russell D. Briggs.....	140
Increasing Soil Temperature in a Northern Hardwood Forest: Effects on Elemental Dynamics and Primary Productivity Patrick J. McHale, Myron J. Mitchell, Dudley J. Raynal, and Francis P. Bowles.....	146
Microhabitat Effects of Litter Temperature and Moisture on Forest-Floor Invertebrate Communities Tim A. Christiansen, Sue A. Perry, and William B. Perry.....	153
Effects of Acidic Precipitation on Leaf Decomposition Rates, Microbial Biomass, and Leaf Pack Macroinvertebrates in Six Streams on the Allegheny Plateau of West Virginia Erik S. Engstrom, Sean K. Meegan, Sue A. Perry, and William B. Perry.....	154
Leaf Litter Processing in West Virginia Mountain Streams: Effects of Temperature and Stream Chemistry Jacquelyn M. Rowe, William B. Perry, and Sue A. Perry.....	157
Leaf Litter Decomposition and Elemental Change in Three Appalachian Mountain Streams of Different pH Steven W. Solada, Sue A. Perry, and William B. Perry.....	161
Methane Evolution from Minnesota Peatlands Elon S. Verry.....	165
Nutrient Cycling in Aggrading Forests: 50 Years of Research at Pack Forest, New York Edwin H. White, Charles A. Z. Buxbaum, and Christopher A. Nowak.....	166
Relative Nitrogen Mineralization and Nitrification Potentials in Relation to Soil Chemistry in Oak Forest Soils Along a Historical Deposition Gradient. Ralph E. J. Boerner and Elaine Kennedy Sutherland.....	167

Mortality Patterns in Oak Forests Along a Climatic and Acid Deposition Gradient Elaine Kennedy Sutherland, Daniel A. Yaussy, and Ralph E. J. Boerner.....	168
--	-----

FOREST AND LANDSCAPE RESPONSES TO REGIONAL STRESS AND MANAGEMENT ACTIVITIES

Thirty-three Year Changes in Above- and Below-Ground Biomass in Northern Hardwood Stands in Vermont A. H. Johnson and G. R. Strimbeck.....	169
---	-----

Evaluation of Methodology for Detecting/Predicting Migration of Forest Species Dale S. Solomon and William B. Leak.....	175
--	-----

Growth Responses of White Oak and Black Oak to Drought Stress Across Gradients of Acid Deposition and Microclimate David LeBlanc and Robert Haack.....	183
---	-----

Assessing the Ability of Plants to Respond to Climatic Change Through Distribution Shifts Mark W. Schwartz.....	184
--	-----

Overview of Current Research on Atmospheric Interactions With Wildland Fires Warren E. Heilman.....	192
--	-----

Synoptic Circulation and Temperature Pattern During Severe Wildland Fires Warren E. Heilman.....	196
---	-----

Atmospheric Mesoscale Simulations of Regional Wildland Fire Episodes: Looking for Weather-Related Factors and Scales of Interaction Brian E. Potter.....	197
---	-----

HUMAN-FOREST INTERACTIONS AND REGIONAL CHANGE

Spatial Patterns in Carbon Storage in a Lake States' Landscape J. C. Bell, D. F. Grigal, P. C. Bates, and C. A. Butler.....	198
--	-----

Carbon Storage in Managed Forests of the Northern Great Lake States Jeanette L. Rollinger and Terry F. Strong.....	203
---	-----

Temperature and Earthworm Effects on C and N Dynamics in Oak Stands Along an Urban-Rural Land Use Gradient R. V. Pouyat, P. M. Groffman, M. M. Carreiro, P. Bohlen, and R. W. Parmelee.....	208
--	-----

AN INTEGRATED NATIONAL MODEL OF THE EFFECTS OF GLOBAL CHANGE ON U.S. FORESTS

Preparation of Forest Inventory and Analysis (FIA) and State Soil Geographic Data Base (STATSGO) Data for Global Change Research in the Eastern United States Louis R. Iverson, Anantha M. G. Prasad, and Charles T. Scott.....	209
--	-----

Regeneration Attributes of Tree Species in the Central Hardwoods Type Elaine Kennedy Sutherland, Betsy J. Hale, and David M. Hix.....	215
--	-----

Herbivorous Insects and Climate Change: Potential Changes in the Spatial Distribution of Forest Defoliator Outbreaks David W. Williams and Andrew M. Liebhold.....	216
---	-----

Predicting the Effects of Tropospheric Ozone on Forest Productivity in the Northeastern U.S. Scott V. Ollinger, John D. Aber, and Peter B. Reich.....	217
Catalog of Long-Term Research Conducted Within the USDA Forest Service, Northeastern Forest Experiment Station Hope R. Barrett.....	226
BOREAL FORESTS AND GLOBAL CHANGE	
Carbon Balance of the Alaskan Boreal Forest John Yarie and Tim Hammond.....	227
An Energy Balance Model for Forest Canopies: A Case Study S. M. Goltz and James A. Smith.....	228
Temporal and Spatial Trends of Fluxes and Concentrations of CO ₂ Above and Within the Canopy at Howland, Maine: Preliminary Results S. M. Goltz.....	229
Dynamics of Calcium Concentration in Stemwood of Red Spruce and Siberian Fir Kevin T. Smith, Walter C. Shortle, Rakesh Minocha, and Vladislav A. Alexeyev.....	230

FOREWORD

The Northern Global Change Program meeting was held March 14-16, 1995, in Pittsburgh, Pennsylvania. Its purpose was to chronicle the research activities of the Northern Global Change Program over the past five years, and outline the implications of these research results for natural resource management. We thank the authors for their participation and for promptly submitting their papers in both paper and electronic form.

CARBON BALANCE OF THE ALASKAN BOREAL FOREST

John Yarie and Tim Hammond¹

Determination of the carbon balance in a broad forest region like the Alaskan boreal forest requires the development of a number of important environmental (state factors) classes to allow for the development of carbon balance estimates. We have used the following factors to develop a regional classification of the state: (1) average monthly temperature May through September (Approximate growing season); (2) total precipitation May through September; and (3) the domain, division and province levels of the ecoregions classification of Alaska (see McNab and Avers 1994).

In addition the following classifications to further subdivide and describe the boreal forest regions of the state are being developed: (1) seasonal land cover classification is being developed by the USGS EROS field office in Anchorage; and (2) age structure of the vegetation in the forest areas.

A forest dynamics model (GAFED) for the Alaskan boreal forest will be run to determine the effects of anticipated global change on the carbon dynamics within the regions of the biogeoclimatic classification. This approach should result in an accurate representation of the carbon dynamics of the forest within an area of land the size of Alaska.

¹Dept. of Forest Sciences, University of Alaska, Fairbanks, AK 99775-7200.

AN ENERGY BALANCE MODEL FOR FOREST CANOPIES: A CASE STUDY

S. M. Goltz¹ and James A. Smith²

The use of thermal scanning devices to map underlying terrain surface temperatures has been recognized as a potential tool for estimating evapotranspiration and latent heat flux densities in forest canopies. Ecologically, knowledge of the temperature distribution of phyto-elements and the underlying soil surface is directly applicable to understanding photosynthetic, heat and mass transfer and soil decomposition rates. Such knowledge of the current state and surface characterization of terrestrial ecosystems is a critical starting point for understanding Global Change processes and modeling the underlying energy exchanges.

A steady-state thermal radiance model to compute thermal exitance and energy balance components within forest canopies is presented. The model treats fully leafed canopies as discrete ensembles of leaves partitioned into slope-angle and height classes. Short-wave energy flux absorbed within the canopy is estimated by solving simplified radiosity equations. Sensible heat exchange is estimated using a logarithmic wind profile above the canopy and a modified exponential profile within the canopy. The latent heat boundary layer resistance is estimated from site-specific measurements summarizing the effects of solar irradiance, air temperature and vapor pressure deficit on stomatal conductance. Comparisons are presented of the basic model with field measurements conducted at a dense spruce-fir forest study site at Howland, Maine. Field data included eddy correlation values of latent and sensible heats as well as IR measurements of foliage temperatures. For clear days the resulting root mean square error in modeled versus measured canopy temperatures was 1.2 °C. Corresponding errors in latent and sensible heat flux energy budget terms were 30 and 32 W m⁻², respectively. For partly cloudy days the root mean square error in predicted temperature was 1.0 °C and corresponding errors in latent and sensible heat were 40 and 110 W m⁻².

¹University of Maine, Orono, ME 04473

²Laboratory for Terrestrial Physics-NASA/GSFC, Greenbelt, MD 20771

TEMPORAL AND SPATIAL TRENDS OF FLUXES AND CONCENTRATIONS OF CO₂ ABOVE AND
WITHIN THE CANOPY AT HOWLAND, MAINE: PRELIMINARY RESULTS

S. M. Goltz¹

In order to develop and evaluate models of net carbon exchange, we have collected profiles of CO₂ through and above the canopy for extended periods over three years as well as collected short-term trial data of diurnal CO₂, water vapor, and sensible heat fluxes above the canopy as measured by eddy correlation. For fluxes, carbon dioxide concentrations were measured with a LI-COR model LI-6251 fast response IRGA housed at the base of the tower. The air sampled from 26 m (7 m above the canopy) was brought through tubing to the IRGA. Both vertical air movement and water vapor density were measured adjacent to the inlet of this tubing at the same elevation using a Campbell Scientific model CA-27 1-D sonic anemometer and a Campbell Scientific model KH-20 krypton hygrometer. For eddy flux analysis, sampling of the three sensors occurred at a frequency of 10 Hz during sampling periods of 30 min. The CO₂ concentration profile through the forest was determined from sequentially collected samples at three levels within the canopy and at 7 meters above canopy using a system consisting of a dedicated LI-6252 CO₂ analyzer, switching valves, pumps, Teflon inlet tubes, calibration gases, and dedicated datalogger (model 21X, Campbell Scientific). These data are used to determine a storage flux of CO₂ associated with the change in CO₂ concentration within the air column. The effects of atmospheric stability, stomatal conductance, and soil moisture on concentrations and fluxes are described.

¹University of Maine, Orono, ME 04469.

DYNAMICS OF CALCIUM CONCENTRATION IN STEMWOOD OF RED SPRUCE AND SIBERIAN FIR

Kevin T. Smith¹, Walter C. Shortle¹, Rakesh Minocha¹, and Vladislav A. Alexeyev²

Abstract: The atmospheric deposition of strong acid anions such as sulfate and nitrate shifts the ion exchange equilibrium in the rooting zone of sensitive forests. Red spruce and other northern coniferous forests are especially sensitive to deposition due to the shallow rooting of trees in a mor-type forest floor. Initially, the deposition of strong acid ions mobilizes essential cations such as calcium (Ca) from ion-exchange sites on soil organic matter. Hypothetically, this mobilization would result in a brief period of increased availability for root uptake. Evidence for this temporary period of increased uptake of essential Ca is the subject of this report. Radial trends in stemwood calcium concentration [Ca] occurred in a common pattern in two sample collections of red spruce from the northeastern United States and in one sample collection of Siberian fir from south-central Siberia, Russia. The [Ca] was measured in wood segments comprising rings that formed during 1871-90, 1891-1910, 1911-30, 1931-50, 1951-70 and 1971-90. For each core, the relative increase or decrease in [Ca] for consecutive periods of wood formation was determined. Previous research indicated that under equilibrium conditions, [Ca] in stemwood decreased in more recently formed wood due to declining numbers of Ca binding sites. Consistent with expectation, the relative frequency of positive change was low among most consecutive periods of growth. Contrary to expectation, however, the frequency of positive increases (48 percent) in [Ca] doubled in 1951-70 compared to 1931-50; this increased frequency was significantly greater ($P \leq 0.01$) than between all other periods.

INTRODUCTION

Declining health, productivity, and diversity of some northern forests have been linked to the rapid increase in emissions of oxides of nitrogen and sulfur (Fig. 1). Evidence that supports this linkage of pollutant emission and deposition to deterioration in forest condition and stability is indirect and subject to alternate explanations (Friedland and others 1993, Johnson and others 1994, Miller and others 1992, and Schlegel and others 1992).




Figure 1. Bars represent percent change in bidecadal mean emissions of S and N oxides (derived from NAPAP 1992). For example, the bar for the 1931-50 bidecade represents an increase in emissions of 25 percent compared to the 1911-30 bidecade. Scatter plots represent the mean percent frequency of positive change in [Ca] from the previous bidecade. Means were calculated from the two data sets of red spruce in the northeastern United States and one data set of Siberian fir from the Sayan Mountains in Siberia, Russia. Each error bar represents the standard error associated with the mean.

¹USDA Forest Service, Northeastern Forest Experiment Station, Durham, New Hampshire 03824 USA.

²V. N. Sukachev Institute for Forestry, Laboratory for Forest Ecosystem Research, Krasnoyarsk, 660036, Russia.

Deposition of strong acid anions such as nitrate and sulfate onto the forest floor tends to mobilize essential base cations e.g., Ca^{+2} and Mg^{+2} , and cations that are antagonistic to the uptake of essential bases, e.g., Al^{3+} (Asp and Berggren 1990; Joslin and others 1988, Kuhn and others 1995). Northern coniferous forests are especially sensitive to the deposition of strong acid anions (Majdi and Persson 1993, Tomlinson 1990). Sensitivity to deposition is greatest where essential cation storage and uptake are from ion-exchange sites in thick, well-defined mor-type forest floor (Cronan 1991). Such mobilization may have three consequences. First, root uptake of essential bases may increase temporarily due to a period of increased availability (Bondiotti and others 1990, Shortle and others, in press). A hypothetical marker of increased availability of essential Ca is presented in this report. Second, root uptake of essential bases may decrease due to the antagonistic binding of Al^{3+} to the ion-exchange sites (Johnson and Fernandez 1992; Smith and others, in press). Third, root uptake of essential bases may continue to decrease as essential bases leach out of the forest floor, eventually impoverishing the rooting zone (Federer and others 1989). These consequences may or may not occur in a continuous series, and the sequence may be interrupted or reversed due to the replenishment of essential bases to the forest floor through stand disturbance.

We have proposed an etiological sequence, the aluminum-induced calcium deficiency syndrome, which may place at risk the health and productivity of sensitive spruce-fir forests in the northeastern United States (Shortle and Smith 1988). Our published reports and those of cooperators tend to support the linkage of pollutant deposition, mobilization and depletion of essential bases, and tree response. Preliminary research on several tree species indicated that stemwood sap increased in pH and/or divalent cation concentration in the mid-1900's (Bondiotti and others 1990; Momoshima and Bondiotti 1990). These changes would be expected during the hypothetical period of enhanced availability due to cation mobilization initiated or enhanced by atmospheric deposition. A comprehensive historical review indicated a marked, widespread decline in essential divalent cations in mor soil of northern forests in the mid- to late-1900's (Shortle and Bondiotti 1992). The ratio of Al:Ca in fine root tips, a hypothetical marker of antagonistic conditions in the rooting zone, was greater at higher elevations and in lower layers of the forest floor (Smith and others, in press). New methods for the extraction (Minocha and others 1994) and determination (Minocha and others 1990) of polyamines allowed the evaluation of the usefulness of these growth regulators as markers of stress caused by Al. Cultures of woody plant tissue dosed with Al indicated that both the timing of the dose and the duration of exposure affected cellular processes in a dose-dependent manner (Minocha and others 1992, Zhou and others 1995). Consequently, the effect of mobilization and antagonistic processes may depend on the phenological condition of the tree.

The time course for these consequences of mobilization is not well established. Rates of cation mobilization and depletion of essential bases differ among forest locations. We do know that the concentration and total amounts of essential bases in the forest floor is limited. Unfortunately, the natural availability of Al is essentially inexhaustible. As the proportion of exchangeable Al increases with respect to exchangeable Ca in the rooting zone, forest stands may begin to decline due to suppressed growth and increased vulnerability to pests and pathogens (Shortle and Smith 1988).

This report presents data on the dynamics of changes in radial trends of calcium concentration [Ca] in stemwood of red spruce from the northeastern United States and Siberian fir (*Abies sibirica* Ledebour) from south-central Siberia, Russia. The [Ca] in these trees have been published (Shortle and others, in press). Ca was emphasized because it is the base element most required for tree growth, and Ca availability in these forest soils may be potentially deficient. The two species were chosen as important components of potentially sensitive forest types. The two regions selected were as widely separated as possible while still being similar habitats.

METHODS

We analyzed the [Ca] in stemwood of three groups of northern conifers. The first data set, NE1, was derived from a previously published report (Bondiotti and others 1990) on divalent cation concentrations in red spruce ($n = 33$ trees) from the northeastern United States (Fig. 2). Data in NE1 were obtained through various methods of sample acquisition, preparation, and analysis. Trends in [Ca] of most individual trees in NE1 were not available, though published data summaries allowed comparison with the other data sets in this investigation. The second data set, RU1, was from Siberian fir ($n = 20$ trees) collected in August 1991 from the Sayan Mountains of the southern Krasnoyarsk region of Siberia, Russia (ca. 53°N , 93°E) (Fig. 2). The third data set, NE2, was from red spruce ($n =$

20 trees) collected in June 1992 from Big Moose Lake, New York, Cone Pond and Crawford Notch, New Hampshire, and Kossuth, Maine, in the Adirondack-New England highlands (ca. 44°N, 69-74°W) (Fig. 2).




Figure 2. Polar and Mercator projection maps showing the locations of data sets NE1 and NE2 in the northeastern US and data set RU1 in south-central Siberia, Russia

For RU1 and NE2 we used identical methods of sample collection and analysis. We removed stemwood cores with an increment borer (12 mm inside diam.) from mature, canopy trees (30 to 60 cm d.b.h.) at 130 cm above groundline. Cores were placed on racks to air-dry on the day they were collected. For chemical analysis we selected single cores from five trees from each of four locations within the sampling areas of RU1 and NE2. Selected cores were free of knots, pitch, and visible indicators of previous injury or infection. Air-dried cores were glued into grooved wooden blocks and sanded with a series of successively finer grits. Following sanding, ring widths were measured and crossdated. Decadal periods of ring formation were marked on the block surface adjacent to the mounted core. Minocha and Shortle (1993) described the preparation, extraction, and analysis of increment cores for cation concentrations. Shavings were removed from each mounted core with a 3.2 mm Ti-coated twist bit drilled perpendicular to the transverse surface at the middle of each decadal growth period. Because of the variable width of growth rings and the fixed diameter of the bit, samples contained varying numbers of growth rings within the

fixed decadal limits. Periods of suppressed, juvenile growth at tree centers were not sampled. Thirty mg of wood shavings from each drill hole were extracted in 6 ml of 0.01 M HCl for three freeze-thaw cycles (Minocha and Shortle 1993). The [Ca] was determined by direct-coupled plasma emission spectroscopy. Each concentration was considered as the concentration for that decadal period of growth. Mean concentrations within each core were calculated for 1891-1910, 1911-30, 193-50, 1951-70, and 1971-90. Comparisons between adjacent bidecadal periods were chosen to lessen the effect of decadal variability and to compensate in part for the width of conducting sapwood during the growth of the tree. For each core, the [Ca] in successive bidecadal periods were scored as having increased, decreased, or remained the same relative to the preceding period. Previous research showed a trend of decreasing [Ca] with increasing radius in the stemwood of red spruce (Momoshima and Bondietti 1990). Consequently, we expected a comparatively high frequency of decreasing [Ca] in wood formed in successively more recent time periods. To test this expectation, the percent frequency of increased [Ca] in successively formed wood was calculated for NE1, RU1, and NE2. Prior to statistical analysis, the percent frequencies were converted using the arcsin-square root transformation to normalize the distribution and stabilize the variance (Bartlett 1947). The transformed percent frequencies of increased concentrations were analyzed by one-way ANOVA. Each adjacent pair of bidecadal periods was considered as a treatment (total of four treatments) and each data set as a replicate ($n = 3$). When justified by a significant F-test ($P < 0.01$), mean frequencies were compared by Fisher's protected LSD test.

RESULTS

The [Ca] tended to decrease along the stemwood radius for most increment cores of both red spruce and Siberian fir (Figs. 3-4). Superimposed on some of the trends of decreased concentration were periods of increased [Ca]. The frequency pattern of positive change in bidecadal mean [Ca] was similar for all three data sets (Fig. 1).

The frequency of increased [Ca] for 1931-50 period to 1951-70 period (48 percent) was significantly greater than for all other comparisons of successive bidecadal periods of wood formation ($P \leq 0.01$, Table 1). The frequencies of positive increases among all other adjacent bidecadal periods did not differ significantly (overall untransformed mean = 21 percent, $SD = 7$).

Table 1. Transformed mean frequencies of positive change in Ca concentration in conifer stemwood

Bidecade ^a	Mean ^b	Standard deviation
1891-1910	0.399	0.072
1911-1930	0.553	0.036
1931-1950	0.516	0.096
1951-1970	0.762	0.031
1971-1990	0.378	0.084

^aThe value for the 1891-1910 bidecade represents the frequency of increased [Ca] from 1871-90 to 1891-1910.

^bRaw frequencies of increased [Ca] for the NE1, NE2, and RU1 data sets were converted with the arcsin-square root transformation prior to statistical analysis. The frequency of increased concentration for the 1951-70 bidecade was significantly greater than for all other periods ($P < 0.01$, Fisher's protected lsd = 0.178). There were no significant differences for other comparisons.




Figure 3 Decadal change in [Ca] in stemwood of red spruce (dataset NE2). For example, the bar for 1950 represents the percent change in [Ca] from 1940 to 1950.




Figure 4. Decadal change in [Ca] concentration in stemwood of Siberian fir (data set RU).

DISCUSSION

Stemwood samples of red spruce and Siberian fir contained similar anomalous increases in [Ca]. The overall trend in most increment cores was a decrease in concentration in more recently formed wood (Figs. 3-4). This general trend was consistent with ion-exchange characteristics previously described for red spruce (Bondiotti and others 1990, Momoshima and Bondiotti 1990). In wood, Ca occurs as a divalent cation bound to ion-exchange sites, primarily anionic carboxylic acid groups of pectins and primary-wall materials (Demarty and others 1984). Ion binding experiments with red spruce wood indicated that the number of available ion-exchange sites per unit of wood tended to decrease with increasing stem diameter (Bondiotti and others 1990). Consequently, when pH and [Ca] in sap were constant, [Ca] in stemwood decreased in more recently formed wood. The underlying mechanism for the progressive decrease in ion-exchange sites has not been established.

The exception to the trend of decreasing [Ca] was the relatively high frequency of increased concentrations within the 1951-70 rings relative to those formed from 1931 to 1950 (Fig. 1). Previous research with red spruce interpreted this increase as a signal of the temporary mobilization of bases in the tree rooting zone (Bondiotti and others 1990; Momoshima and Bondiotti 1990). Such a period of cation mobilization would have been expected to result from anionic loading of sensitive forest soil with the atmospheric deposition of nitrates and sulfates (Shortle and Bondiotti 1992). The mobilization of chemical bases in soil that was recorded in stemwood chemistry was coincident in time with the greatest increase in SO₂ and NO_x emissions (Fig. 1). The presence of a similar signal of increased [Ca] within Siberian fir emphasizes the need to explain this anomaly more fully in future research.

Hypothetically, [Ca] in healthy stemwood could increase due to increasing numbers of ion-exchange sites, increasing the pH of sap or of [Ca] in the xylem stream, or some combination of these. We reject the explanation of increased numbers of ion-exchange sites as all evidence shows decreasing numbers of exchange sites along the stemwood radius (Bondiotti and others 1990; Smith and Shortle, in press). The concentration of exchangeable Ca in stemwood relative to the [Ca] and pH of the sap has the characteristics of a Donnan equilibrium in which 60 to 80 percent of binding sites are occupied with Ca (Momoshima and Bondiotti 1990). An increase in saturation of stemwood binding sites with Ca from 60 to 80 percent can be achieved by an increase in pH from 5.0 to 5.5 or an increase in [Ca] in the sap from 0.5 to 2.5 mequiv per L, or combinations of the two. This relationship is complicated somewhat in that other divalent, e.g., Mg, and monovalent (K) cations also participate in the chemistry of sap and stemwood. Nevertheless, the higher [Ca] within wood formed during 1951-70 relative to the previous 20 years suggest a concomitant change in sap chemistry.

What internal or external factors could account for changes in sap chemistry in two widely separated parts of the northern coniferous forest (Smith and Shortle, in press)? Potential internal factors include the constitutive transformation of sapwood to heartwood. The sapwood:heartwood boundary was present in 1951-70 period for essentially all trees in the three data sets. If this maturation process was responsible, we expect that a substantially greater frequency of increased [Ca] than the observed 48 percent would have occurred between the periods of 1931-50 and 1951-70 period. Also, there are no published reports of the accumulation of Ca or increased pH at or near the sapwood:heartwood boundary in any species of spruce or fir. The internal, inducible factors that affect wood and sap chemistry include stem infection and the response of trees to injury and infection (Smith and Shortle, in press). The similarity in timing, pattern, and geographic separation argues against a common pathological explanation. Base-cation mobilization might increase the concentrations of Ca available to roots, resulting in increased concentrations within wood. Such mobilization could occur from disturbances such as fire, though there are no records of fire on these sites during this time period. Climatic events that combine the effects of rain and prolonged hot summers might produce pulses of anions into the soil water, resulting in mobilization of base cations. However, no such climatic event has been documented for these widely separated forests. Base-cation mobilization due to the deposition of strong acid anions is coincident in time and consistent with known mechanisms that would increase the frequency of increased [Ca] in wood formed in 1951-70. The greatest increase in rates of emissions in the northeastern United States occurred during this period (Fig. 1). Pollution monitoring data for the Sayan Mountain region apparently are not available. Although this part of Siberia is remote, it is not necessarily free of the influence of regional, industrial pollution.

Because of the legislative response to public concern, total emissions of SO₂ and NO_x have begun to decrease (Fig. 1). However, the extent and severity of damage to forest stands due to previous emissions are not known. The biogeochemical links among emission, deposition, essential element uptake, and forest health are poorly understood. Seemingly, some shifts in the biogeochemical equilibrium have occurred, though it is unclear whether a new, stable equilibrium has been reached.

ACKNOWLEDGMENTS

We were assisted in this work by Dr. Vecheslav Kharuk (Sukachev Institute of Forestry), Kenneth R. Dudzik (USDA Forest Service), and Paul J. Krusic and Stephanie Long (University of New Hampshire).

LITERATURE CITED

- Asp, H.; Berggren, D. 1990. Phosphate and calcium uptake in beech (*Fagus sylvatica*) in the presence of aluminum and natural fulvic acids. *Physiologia Plantarum*. 80: 307-314.
- Bartlett, M. S. 1947. The use of transformations. *Biometrics*. 3: 39-52.
- Bondietti, E. A., Momoshima, N., Shortle, W. C. Smith, K. T. 1990. A historical perspective on divalent cation trends in red spruce stemwood and the hypothetical relationship to acidic deposition. *Canadian Journal of Forest Research*. 20: 1850-1858.
- Cronan, C. S. 1991. Differential adsorption of Al, Ca, and Mg by roots of red spruce (*Picea rubens* Sarg.). *Tree Physiology*. 8: 227-237.
- Demarty, M., Morvan, C.; Thellier, M. 1984. Calcium and the cell wall. *Plant Cell and Environment* 7: 441-448.
- Federer, C. A.; Hornbeck, J. W.; Tritton, L. M.; Martin, C. W.; Pierce, R. S.; Smith, C. T. 1989. Long-term depletion of calcium and other nutrients in eastern US forests. *Environmental Management*. 13: 593-601.
- Friedland, A. J.; Miller, E. K.; Boyce, R. L. 1993. Element distribution in red spruce (*Picea rubens*) fine roots; evidence for aluminum toxicity at Whiteface Mountain. Discussion. *Canadian Journal of Forest Research*. 23: 1732-1733.
- Johnson, A. H.; Andersen, S. B.; Siccama, T. G. 1994. Acid rain and soils of the Adirondacks. I. Changes in pH and available calcium, 1930-1984. *Canadian Journal of Forest Research*. 24: 39-45.
- Johnson, D. W.; Fernandez, I. 1992. Soil-mediated effects of atmospheric deposition on eastern U.S. spruce-fir forest. In: Eagar, C.; Adams, M. B., eds. *Ecology and decline of red spruce in the eastern United States*. New York: Springer-Verlag. p. 205-221.
- Joslin, J. D.; Kelley, J. M.; Wolfe, M. H.; Rustad, L. E. 1988. Elemental patterns in roots and foliage of mature spruce across a gradient of soil aluminum. *Water, Air and Soil Pollution*. 40: 375-390.
- Kuhn, A. J.; Bauch, J.; Schröder, W. H. 1995. Monitoring uptake and contents of Mg, Ca and K in Norway spruce as influenced by pH and Al, using microprobe analysis and stable isotope labelling. *Plant and Soil*. 168-169: 135-150.
- Majdi, H.; Persson, H. 1993. Spatial distribution of fine roots, rhizosphere and bulk-soil chemistry in an acidified *Picea abies* stand. *Scandinavian Journal of Forest Research*. 8: 147-155.

- Miller, E. K.; Huntington, T. G.; Johnson, A. H.; Friedland, A. J. 1992. Aluminum in soil solutions from a subalpine spruce-fir forest at Whiteface Mountain, New York. *Journal of Environmental Quality*. 21: 345-352.
- Minocha, R.; Minocha, S. C.; Long, S. L.; Shortle, W. C. 1992. Effects of aluminum on DNA synthesis, cellular polyamines, polyamine biosynthetic enzymes and inorganic ions in cell suspension cultures of a woody plant, *Catharanthus roseus*. *Physiologia Plantarum*. 85: 417-424.
- Minocha, R.; Shortle, W. C. 1993. Fast, safe, and reliable methods for extraction of major inorganic cations from small quantities of woody plant tissues. *Canadian Journal of Forest Research*. 23: 1645-1654.
- Minocha, R.; Shortle, W. C.; Long, S. L.; Minocha, S. C. 1994. A rapid and reliable procedure for extraction of cellular polyamines and inorganic ions from plant tissues. *Journal of Plant Growth Regulation*. 13: 187-193.
- Minocha, S. C., Minocha, R., and Robie, C. A. 1990. High performance liquid chromatographic method for the determination of dansyl-polyamines. *Journal of Chromatography*. 511: 177-183.
- Momoshima, N.; Bondietti, E. A. 1990. Cation binding in wood: applications to understanding historical changes in divalent cation availability to red spruce. *Canadian Journal of Forest Research*. 20: 1840-1849.
- National Acid Precipitation Assessment Program. 1992 Report to Congress. Washington, D.C.: National Acid Precipitation Assessment Program. 130 p.
- Schlegel, H.; Amundson, R. G.; Hüttermann, A. 1992. Element distribution in red spruce (*Picea rubens*) fine roots; evidence for aluminum toxicity at Whiteface Mountain. *Canadian Journal of Forest Research*. 22: 1132-1138.
- Shortle, W. C.; Bondietti, E. A. 1992. Timing, magnitude, and impact of acidic deposition on sensitive forest sites. *Water, Air and Soil Pollution*. 61: 253-267.
- Shortle, W. C.; Smith, K. T. 1988. Aluminum-induced calcium deficiency syndrome in decline red spruce. *Science*. 240: 1017-1018.
- Shortle, W. C.; Smith, K. T.; Minocha, R.; Alexeyev, V. Similar patterns of change in stemwood calcium concentration in red spruce and Siberian fir. *Journal of Biogeography*. [in press].
- Smith, K. T.; Shortle, W. C. Tree biology and dendrochemistry. In: Dean, J. S.; Meko, D. M.; Swetnam, T. W., eds. *Tree rings, environment, and humanity*. Tucson. Radiocarbon. [in press].
- Smith, K. T.; Shortle, W. C.; Ostrofsky, W. D. Aluminum and calcium in fine root tips of red spruce collected from the forest floor. *Canadian Journal of Forest Research*. [in press].
- Tomlinson, G. H. 1990. Effects of acid deposition on the forests of Europe and North America. Boca Raton, FL: CRC Press. 281 p.
- Zhou, X.; Minocha, R.; Minocha, S. 1995. Physiological responses of suspension cultures of *Catharanthus roseus* to aluminum: changes in polyamines and inorganic ions. *Journal of Plant Physiology*. 145: 277-284.