



**The VEMAP Integrated Database for Modelling United States
Ecosystem/Vegetation Sensitivity to Climate Change**

T. G. F. Kittel, N. A. Rosenbloom, T. H. Painter, D. S. Schimel

Journal of Biogeography, Volume 22, Issue 4/5, Terrestrial Ecosystem Interactions with
Global Change, Volume 2 (Jul. - Sep., 1995), 857-862.

Stable URL:

<http://links.jstor.org/sici?sici=0305-0270%28199507%2F199509%2922%3A45%3C857%3ATVIDFM%3E2.0.CO%3B2-N>

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

Journal of Biogeography is published by Blackwell Science, Inc.. Please contact the publisher for further permissions regarding the use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/blacksci-inc.html>.

Journal of Biogeography

©1995 Blackwell Science, Inc.

JSTOR and the JSTOR logo are trademarks of JSTOR, and are Registered in the U.S. Patent and Trademark Office. For more information on JSTOR contact jstor-info@umich.edu.

©2002 JSTOR

The VEMAP integrated database for modelling United States ecosystem/vegetation sensitivity to climate change

T. G. F. KITTEL*†, N. A. ROSENBLOOM*‡, T. H. PAINTER*, D. S. SCHIMEL†§ and VEMAP MODELLING PARTICIPANTS¶ *Climate System Modeling Program, University Corporation for Atmospheric Research, Box 3000, Boulder, CO 80307-3000, U.S.A., †Natural Resource Ecology Laboratory, Colorado State University, Fort Collins, CO 80523, U.S.A., ‡Department of Geology and Institute for Arctic and Alpine Research, University of Colorado, Boulder, CO 80309, U.S.A. and §Climate and Global Dynamics Division, National Center for Atmospheric Research, Box 3000, Boulder, CO 80307-3000, U.S.A.

Abstract. For the Vegetation/Ecosystem Modelling and Analysis Project (VEMAP), we developed a model database of climate, soils and vegetation that was compatible with the requirements of three ecosystem physiology models and three vegetation life-form distribution models. A key constraint was temporal, spatial and physical consistency among data layers to provide these daily or monthly time step models with suitable common inputs for the purpose of model inter-comparison. The database is on a 0.5° latitude/longitude grid for the conterminous United States. The set has both daily and monthly representations of the same long-term climate. Daily temperature and precipitation were stochastically simulated with WGEN and daily solar radiation and humidity empir-

ically estimated with CLIMSIM. We used orographically adjusted precipitation, surface temperature and surface wind-speed monthly means to maintain consistency among these fields and with vegetation distribution. Vegetation classes were based on physiognomic and physiological properties that influence biogeochemical dynamics. Soils data include characteristics of the 1–4 dominant soils per cell to account for subgrid variability.

Key words. Climate, soils, integrated database, spatial interpolation, ecosystem physiological modelling, vegetation life-form distribution modelling, United States.

INTRODUCTION

Continental and global simulations of ecosystem physi-

¶VEMAP modelling participants are: Jerry M. Melillo (Chair), Yude Pan, David W. Kicklighter and A. David McGuire (Marine Biological Laboratory, Woods Hole, MA 02543, U.S.A.); Ronald P. Neilson (USDA Forest Service, Corvallis, OR 97333, U.S.A.) and Jesse Chaney (Department of Botany and Plant Pathology, Oregon State University, Corvallis, OR 97333, U.S.A.); Dennis S. Ojima, Rebecca McKeown, William J. Parton and William M. Pulliam (NREL, Colorado State Univ., Ft. Collins, CO 80523, U.S.A.); I. Colin Prentice and Alex Haxeltine (University of Lund, Lund, Sweden); Steven W. Running, Lars L. Pierce, Ramakrishna R. Nemani and E. Raymond Hunt, Jr (School of Forestry, University of Montana, Missoula, MT 59812, U.S.A.); Thomas M. Smith, Brian Rizzo (Department of Environmental Science, University of Virginia, Charlottesville, VA 22903, U.S.A.) and F. Ian Woodward (Department of Animal and Plant Sciences, University of Sheffield, Sheffield, S10 2TN, U.K.).

National Center for Atmospheric Research is sponsored by the National Science Foundation.

Corresponding author: Timothy Kittel, CSMP, UCAR, P.O. Box 3000, Boulder, CO 80307-3000, U.S.A.

ology and vegetation distribution and their sensitivity to climate and CO₂ change have recently been evaluated in a number of modelling studies (e.g. Melillo *et al.*, 1993; Monserud, Tchebakova & Leemans, 1993; Neilson, 1993; Ojima *et al.*, 1993; Running & Nemani, 1991). Comparison of these results is restricted by the use of different driving-variable and boundary-condition datasets and different scenarios of altered forcing. In addition, with few exceptions (cf. Pastor & Post, 1988), studies have not evaluated the joint response of ecosystem nutrient cycling and biome redistribution to altered forcing across large domains.

The Vegetation/Ecosystem Modelling and Analysis Project (VEMAP) is a multi-institutional, international effort whose goal is to evaluate the sensitivity of terrestrial ecosystem and vegetation processes to altered climate forcing and elevated atmospheric CO₂. The project's objectives are (1) inter-comparison of the climate and CO₂ sensitivity of three ecosystem physiology models (plant production and soil biogeochemical process models) and three vegetation life-form distribution models (biome distribution models) and (2) the one-way linkage of vegetation distri-

bution and ecosystem physiology models to evaluate the joint response of these processes to altered conditions. To accomplish these objectives, the models require common boundary conditions and driving variables so that differences in model results arise only from the models and their implementation rather than from differences in inputs. In addition, a consistent dataset was required for paired simulations with vegetation distribution and ecosystem physiology models.

The three biome distribution models in the VEMAP set are BIOME2 (Prentice *et al.*, 1992; Haxeltine, Prentice, Cresswell, 1995, DOLY (Woodward *et al.*, 1995), and MAPSS (Neilson, 1995). The three ecosystem physiology models are BIOME-BGC/GESSys (Running & Hunt, 1993), CENTURY (Parton *et al.*, 1987, 1994), and TEM (Melillo *et al.*, 1993; McGuire *et al.*, 1993). DOLY and BIOME-BGC are daily time step models, while the others run with monthly inputs.

In this paper, we describe briefly the VEMAP integrated model input database and its development. Access to the database is open to the modelling community via the internet on the World Wide Web site:

<http://www.cgd.ucar.edu:80/vemap/>

or contact the corresponding author.

DATABASE REQUIREMENTS AND CONCEPTUAL FRAMEWORK

Current-climate driving variables required by the suite of models were both daily and monthly fields of minimum and maximum surface air temperature, precipitation, total incident solar radiation, surface air humidity and surface wind-speed. Boundary conditions included soil properties (including texture and water holding capacity) and vegetation type. Altered climate scenarios were also needed for the same set of driving variables. The domain for the VEMAP simulations was the conterminous United States, using a 0.5° latitude/longitude interval grid.

Beyond supplying needed variables on a specified grid, additional requirements for the model inter-comparison dataset were (1) physical consistency among driving variables and boundary conditions as required for any stand-alone model run and (2) the need to match differing individual model requirements for the same variables in a manner that did not violate the conceptual basis of any of the models. Physical consistency among variables was achieved in the climate database by (1) using monthly mean data developed with spatial interpolation techniques that account for effects of topography on climate and (2) using covariance information and empirical relationships among related variables in the generation of daily data. Altered climate scenarios were also constrained to maintain some degree of physical consistency.

In the soils dataset, physical consistency was incorporated by representing a grid cell's soil by a set of dominant soil profiles, rather than by a simple cell average of properties. Because soil processes, such as soil water balance, are non-linearly related to soil texture and other soil parame-

ters, simulations based on dominant soil profiles and their frequency distribution can account for soil dynamics that would be lost if averaged soil properties were used.

Matching model requirements was also a key constraint in the development of consistent daily and monthly climate datasets. Daily time step models require realistic daily variance structure and, as a result, daily 'normals' (e.g. 30-year averages by day of year) would not suffice. On the other hand, the monthly time step models generally require long-term monthly climatological data. Therefore, the daily climate dataset had to have daily variances and covariances characteristic of an actual weather record but maintain, on a monthly basis, the same climate as the long-term climatology. This was accomplished by (1) stochastically generating daily climates for each grid cell based on temporal statistical properties of nearby weather stations and (2) constraining the monthly means of the created daily record to match those of the cell's long-term climate. These processes are described in more detail in the next section. Matching model requirements also played a role in the classification of vegetation types. We based the vegetation classes on physiognomic and physiological characteristics that best corresponded to the conceptual perspectives of both classes of models.

DATASET DEVELOPMENT

Grid description

The VEMAP grid is a 0.5° latitude/longitude grid with cell centres at 0.5° and 1.0° points and covers the conterminous United States (Fig. 1). The dataset includes fields of cell latitude, longitude, elevation and area. Elevation was aggregated from 10-minute Navy Fleet Numeric Oceanographic Center (NFNO, 1985) data (D. Kicklighter and A. D. McGuire, pers. comm.).

Climate variables

For the model inter-comparison, the common climate input dataset needed to be:

1. Spatially consistent, with climate data reflecting orographic effects.
2. Temporally consistent, with daily and monthly representations of the same long-term base climate.
3. Physically consistent, maintaining relationships among climate variables.

These requirements were met in a three-step process.

Step 1: topographically adjusted interpolation of monthly mean temperature, precipitation and wind-speed. Monthly mean minimum and maximum temperatures from 4613 station normals (NCDC, 1992) were adiabatically adjusted to sea level using algorithms of Marks & Dozier (1992), interpolated to the grid, and then re-adjusted to grid elevations. Mean monthly precipitation was spatially aggregated from a 10-km gridded U.S. dataset




FIG. 1. VEMAP gridded mean January minimum temperature, spatially interpolated with adiabatic adjustment from surface station means.

TABLE 1. VEMAP vegetation classes, based on physiognomy (life-form, leaf seasonal duration and leaf shape) and, for grasslands, physiology (photosynthetic pathway).

Tundra
1 Tundra
Forest
2 Boreal coniferous forest (includes boreal transition & subalpine forests)
3 Maritime temperate coniferous forest
4 Continental temperate coniferous forest
5 Cool temperate mixed forest
6 Warm temperate/subtropical mixed forest
7 Temperate deciduous forest
8 Tropical deciduous forest
9 Tropical evergreen forest
Xeromorphic woodlands and forests
10 Temperate mixed xeromorphic woodland
11 Temperate conifer xeromorphic woodland
12 Tropical thorn woodland
Savannas
13 Temperate/subtropical deciduous savanna
14 Warm temperate/subtropical mixed savanna
15 Temperate conifer savanna
16 Tropical deciduous savanna
Grasslands
17 C3 grasslands (includes short to tall C3 grasslands)
18 C4 grasslands (includes short to tall C4 grasslands)
Shrublands
19 Mediterranean shrubland
20 Temperate arid shrubland
21 Subtropical arid shrubland

developed using PRISM (Daly, Neilson & Phillips, 1994). PRISM models precipitation distribution by (1) dividing the terrain into topographic facets of similar aspect, (2) developing precipitation–elevation regressions for each facet for a given region based on station data, and (3) using these regressions to spatially extrapolate station precipitation to cells that are on similar facets. Mean monthly windspeeds at 10 m height were derived from Marks (1990) based on DOE seasonal wind averages (Elliott *et al.*, 1986). The DOE winds were topographically corrected to account for greater windspeeds over regions with high terrain by Elliott *et al.* (1986).

Step 2: stochastic generation of daily temperature and precipitation data. We used a daily weather generator, a modified version of WGEN (Richardson, 1981; Richardson & Wright, 1984), to statistically simulate daily temperature and precipitation. Monthly means of daily values were matched to the long-term means. Parameterization of WGEN was based on daily records from 870 stations (Shea, 1984; Eddy, 1987). WGEN created records that realistically represent daily variances and temporal autocorrelation (e.g. persistence of dry and wet days) and maintain the physical relationship between daily precipitation and temperature. For example, days with precipitation tended to have higher minimum and lower maximum temperatures than days with no precipitation. These charac-

teristics of the daily set are required for the daily-based models to adequately simulate water balance and ecological dynamics.

Step 3: empirical estimation of daily and monthly solar radiation and humidity. We used CLIMSIM (a simplified version of MT-CLIM for flat surfaces; Running, Nemani & Hungerford, 1987; Glassy & Running, 1994) to estimate daily total incident solar radiation, daily irradiance and surface humidity based on the daily minimum and maximum temperature and precipitation. The objective of this approach was to maintain physical relationships among temperature, precipitation, solar radiation and humidity on a daily basis. Monthly means of radiation and humidity were derived from the daily values. CLIMSIM determines daily solar radiative inputs based on latitude, elevation, diurnal range of temperature and occurrence of precipitation using algorithms of Gates (1981) and Bristow & Campbell (1984). Mean daily irradiance was also calculated based on day length. CLIMSIM estimates empirically daily vapour pressure and relative humidity by assuming that on a daily basis minimum temperatures reach the dew point. Because this is often not the case in arid regions, we adjusted daily humidity data downward so that the monthly means match those of Marks (1990).

Boundary conditions

Vegetation types were defined physiognomically in terms of dominant life-form and leaf characteristics (including leaf seasonal duration, shape and size) and, in the case of grasslands, physiologically with respect to dominance of species with C3 versus C4 photosynthetic pathway (Table 1). The physiognomic classification criteria were based on our understanding of vegetation characteristics that influence biogeochemical dynamics (cf. Running *et al.*, 1994). The U.S. distribution of these types was based on a 0.5° latitude/longitude gridded map of Kùchler's (1964, 1975) potential natural vegetation (D. Kicklighter and A. D. McGuire, pers. comm.). These vegetation types represent a match between vegetation classes simulated by the biome distribution models and ecosystem types required as boundary conditions by the ecosystem physiology models.

Soil texture, depth and other properties were determined for dominant soil types based on Kern's (1994, 1995) 10-km gridded Soil Conservation Service national-level (NATSGO) database. To aggregate the Kern dataset spatially to the 0.5° grid, we used cluster analysis to group the 10-km subgrid elements into modal soil types. In this statistical approach, cell soil properties are represented by a set of 1–4 modal soil profiles, rather than by an 'average soil' that may not correspond to an actual soil in the region.

Climate scenarios

Climate scenarios for eight climate change experiments were included in the database. Seven of these experiments are from atmospheric general circulation models (GCMs) implemented with a simple 'mixed-layer' ocean representa-

tion that includes heat storage and vertical exchange of heat and moisture with the atmosphere. The experiments are $1 \times \text{CO}_2$ and $2 \times \text{CO}_2$ equilibrium runs and are archived at the National Center for Atmospheric Research (NCAR; Jenne, 1992). These runs are from the Canadian Climate Centre (CCC; Boer, McFarlane & Lazare, 1992), Goddard Institute for Space Studies (GISS; Hansen *et al.*, 1984), Geophysical Fluid Dynamics Laboratory (GFDL, including R15 $4.5^\circ \times 7.5^\circ$ grid runs with and without Q-flux corrections, and an R30 $2.22^\circ \times 3.75^\circ$ grid run; Manabe & Wetherald, 1987; Manabe & Wetherald in Mitchell *et al.*, 1990), Oregon State University (OSU; Schlesinger & Zhao, 1989), United Kingdom Meteorological Office (UKMO; Wilson & Mitchell, 1987). The CCC and GFDL R30 runs are among the high resolution GCM experiments reported in IPCC (1990). We also included in the VEMAP scenarios results for the conterminous United States from a nested regional climate model (RegCM, based on the Pennsylvania State University/NCAR mesoscale model MM4) by Giorgi, Brodeur & Bates (1994) driven by $1 \times$ and $2 \times \text{CO}_2$ GCM runs (Thompson & Pollard, 1995a,b).

Changes in temperature were represented as differences and those for precipitation, solar radiation, vapour pressure (not available for the CCC run) and horizontal windspeed as change ratios. We determined differences for relative humidity based on the VEMAP base climate and climate changes for temperature, vapour pressure and surface pressure. We interpolated change fields spatially to the 0.5° grid using simple interpolation techniques. This provided smoothed change fields that could be applied to the VEMAP base climate to generate altered-climate inputs.

A key issue in the generation of altered climates based on climate model output is the strong possibility of physical inconsistencies in the new climates. Change ratios from the NCAR archive have an imposed upper limit of 5.0, providing some constraint on these changes. In the creation of the new climates, additional checks were that (1) total incident solar radiation did not exceed potential solar input and (2) relative humidity stayed in the range from 0 to 100%.

SUMMARY

The VEMAP database is an integrated dataset for ecological modelling studies covering the conterminous United States. Physical consistency among climate drivers was accomplished by (1) spatial interpolation that incorporates topographic effects and (2) stochastic generation and empirical estimation of daily data that accounts for temporal variance structure and covariance among variables. Vegetation types were defined in a manner compatible with the needs of both ecosystem physiological and vegetation life-form distribution models. We developed modal soil data to provide boundary conditions that more closely represent actual soils than would a coarse-grid average of soil properties. Together these methods assure the consistency among model inputs, and between inputs and model assumptions, required to adequately capture spatial patterns of ecological change.

ACKNOWLEDGMENTS

Development of the VEMAP database was supported by NASA Mission to Planet Earth, Electric Power Research Institute (EPRI), USDA Forest Service Southern Region Global Change Research Program, and by NSF-ATM Climate Dynamics Program through UCAR's Climate System Modeling Program (CSMP). We thank Lou Pitelka, Susan Fox, Tony Janetos and Hermann Gucinski for their support of VEMAP. Thanks to Hank Fisher and Donna Beller for programming and data management support, Gaylynn Potemkin for help with manuscript preparation, Roy Barnes, Chris Daly, Filippo Giorgi, Roy Jenne, Dennis Joseph, Jeff Kern, John Kineman, Danny Marks, Christine Shields and Dennis Shea for access to datasets and model output, and NCAR Climate and Global Dynamics Division for computer systems support. We also wish to thank Genasys II and StatSci for their support.

REFERENCES

- Boer, G.J., McFarlane, N.A. & Lazare, M. (1992) Greenhouse gas-induced climate change simulated with the CCC second-generation general circulation model. *Bull. Am. Met. Soc.* **5**, 1045–1077.
- Bristow, K.L. & Campbell, G.S. (1984) On the relationship between incoming solar-radiation and daily maximum and minimum temperature. *Agric. For. Meteorol.* **31**, 159–166.
- Daly, C., Neilson, R.P. & Phillips, D.L. (1994) A statistical-topographic model for mapping climatological precipitation over mountainous terrain. *J. appl. Meteorol.* **33**, 140–158.
- Eddy, A. (1987) *Interpolated daily surface data: 1881–1985*. Oklahoma Climatological Survey. NCAR dataset DS508.0. Unpublished dataset.
- Elliott, D.L., Holladay, C.G., Barchet, W.R., Foote, H.P. & Sandusky, W.F. (1986) *Wind energy resource atlas*. Solar Technical Information Program, U.S. Department of Energy.
- Gates, D.M. (1981) *Biophysical ecology*. Springer-Verlag, New York.
- Giorgi, F., Brodeur, C.S. & Bates, G.T. (1994) Regional climate change scenarios over the United States produced with a nested regional climate model. *J. Clim.* **7**, 375–399.
- Glassy, J.M. & Running, S.W. (1994) Validating diurnal climatology logic of the MTCLIM model across a climatic gradient in Oregon. *Ecol. Appl.* **4**, 248–257.
- Hansen, J., Lacis, A., Rind, D., Russell, G., Stone, P., Fung, I., Ruedy, R. & Lerner, J. (1984) Climate sensitivity: analysis of feedback mechanisms. *Climate processes and climate sensitivity* (ed. by J.E. Hansen and T. Takahashi). American Geophysical Union, Washington, D.C.
- Haxeltine, A., Prentice, I.C. & Cresswell, I.D. (1995). A coupled carbon and water flux model to predict vegetation structure. *J. Veg. Sci.* (in press).
- IPCC (1990) *Climate change. The IPCC scientific assessment* (ed. by J.T. Houghton, G.J. Jenkins and J.J. Ephraums). Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge.
- Jenne, R.L. (1992) Climate model description and impact on terrestrial climate, pp. 145–164. *Global climate change: implications, challenges and mitigation measures* (ed. by S.K. Majumdar, L.S. Kalkstein, B. Yarnal, E.W. Miller and L.M. Rosenfeld). Pennsylvania Academy of Science.
- Kern, J.S. (1994) Spatial patterns of soil organic carbon in the contiguous United States. *J. Soil Sci. Soc. Am.* **58**, 439–455.

- Kern, J.S. (1995) Geographic patterns of soil water-holding capacity in the contiguous United States. *J. Soil Sci. Soc. Am.* **59**, 1126–1133.
- Küchler, A.W. (1964) *Manual to accompany the map, potential natural vegetation of the conterminous United States*. Spec. Pub. No. 36. American Geographical Society, New York.
- Küchler, A.W. (1975) *Potential natural vegetation in the United States*, 2nd edn (map 1:3,168,000). American Geographical Society, New York.
- Manabe, S. & Wetherald, R.T. (1987) Large-scale changes in soil wetness induced by an increase in carbon dioxide. *J. Atmos. Sci.* **44**, 1211–1235.
- Marks, D. (1990) The sensitivity of potential evapotranspiration to climate change over the continental United States. *Biospheric feedbacks to climate change: the sensitivity of regional trace gas emissions, evapotranspiration, and energy balance to vegetation redistribution* (ed. by H. Gucinski, D. Marks and D.P. Turner), pp. IV-1–IV-31. EPA/600/3-90/078. U.S. Environmental Protection Agency, Corvallis, OR.
- Marks, D. & Dozier, J. (1992) Climate and energy exchange at the snow surface in the alpine region of the Sierra Nevada: 2. Snow cover energy balance. *Water Resources Res.* **28**, 3043–3054.
- McGuire, A.D., Joyce, L.A., Kicklighter, D.W., Melillo, J.M., Esser, G. & Vorosmarty, C.J. (1993) Productivity response of climax temperate forests to elevated temperature and carbon dioxide: a North American comparison between two global models. *Clim. Change*, **24**, 287–310.
- Melillo, J.M., McGuire, A.D., Kicklighter, D.W., Moore III, B., Vorosmarty, C.J. & Schloss, A.L. (1993) Global climate change and terrestrial net primary production. *Nature*, **363**, 234–240.
- Mitchell, J.F.B., Manabe, S., Meleshko, V. & Tokioka, T. (1990) Equilibrium climate change—and its implications for the future. *Climate change. The IPCC scientific assessment* (ed. by J.T. Houghton, G.J. Jenkins and J.J. Ephraums), pp. 131–172. Cambridge University Press, Cambridge.
- Monserud, R.A., Tchebakova, N.M. & Leemans, R. (1993) Global vegetation change predicted by the modified Budyko method. *Clim. Change*, **25**, 59–83.
- NCDC (National Climatic Data Center) (1992) *1961–1990 monthly station normals tape*. U.S. Department of Commerce, data tape TD 9641.
- NFNOG (Navy Fleet Numeric Oceanographic Center) (1985) *10-minute global elevation, terrain, and surface characteristics* (re-processed by NCAR and NGDC), digital dataset. NOAA National Geophysical Data Center Boulder, CO.
- Neilson, R.P. (1993) Vegetation redistribution: a possible biosphere source of CO₂ during climatic change. *Water, Air Soil Pollut.* **70**, 659–673.
- Neilson, R.P. (1995) A model for predicting continental-scale vegetation distribution and water balance. *Ecol. Appl.* **5**, 362–385.
- Ojima, D.S., Parton, W.J., Schimel, D.S., Scurlock, J.M.O. & Kittel, T.G.F. (1993) Modeling the effects of climatic and CO₂ changes on grassland storage of soil C. *Water, Air Soil Pollut.* **70**, 643–657.
- Parton, W.J., Schimel, D.S., Cole, C.V. & Ojima, D.S. (1987) Analysis of factors controlling soil organic levels of grasslands in the Great Plains. *J. Soil Sci. Soc. Am.* **51**, 1173–1179.
- Parton W.J., Ojima, D.S., Cole, C.V., & Schimel, D.S. (1994) A general model for soil organic matter dynamics: sensitivity to litter chemistry, texture and management. *Quantitative modeling of soil forming processes* (ed. by R.B. Bryant and R.W. Arnold), pp. 147–167. Soil Science Society of America Spec. Publ. **39**, Madison, WI.
- Pastor, J. & Post, W.M. (1988) Response of northern forests to CO₂-induced climate change. *Nature*, **334**, 55–58.
- Prencice, I.C., Cramer, W., Harrison, S.P., Leemans, R., Monserud, R.A. & Solomon, A.M. (1992) A global biome model based on plant physiology and dominance, soil properties and climate. *J. Biogeogr.* **19**, 117–134.
- Richardson, C.W. (1981) Stochastic simulation of daily precipitation, temperature and solar radiation. *Water Resources Res.* **17**, 182–190.
- Richardson, C.W. & Wright, D.A. (1984) *WGEN: a model for generating daily weather variables*, ARS-8. U.S. Department of Agriculture, Agricultural Research Service.
- Running, S.W. & Hunt, E.R. (1993) Generalization of a forest ecosystem process model for other biomes, BIOME-BGC, and an application for global-scale models. *Scaling physiological processes: leaf to globe* (ed. by R. Ehleringer and C.B. Field), pp. 141–158. Academic Press, San Diego, CA.
- Running, S.W. & Nemani, R.R. (1991) Regional hydrologic and carbon balance responses of forests resulting from potential climate change. *Clim. Change*, **19**, 349–368.
- Running, S.W., Loveland, T.R. & Pierce, L.L. (1994) A vegetation classification logic based on remote sensing for use in global biogeochemical models. *Ambio*, **23**, 77–81.
- Running, S.W., Nemani, R.R. & Hungerford, R.D. (1987) Extrapolation of synoptic meteorological data in mountainous terrain and its use for simulating forest evapotranspiration and photosynthesis. *Can. J. For. Res.* **17**, 472–483.
- Schlesinger, M.E. & Zhao, Z.C. (1989) Seasonal climate changes induced by doubled CO₂ as simulated by the OSU atmospheric GCM-mixed layer ocean model. *J. Clim.* **2**, 459–495.
- Shea, D.J. (1984) *The annual cycle. Part I. The annual variation of surface temperature over the United States and Canada*. NCAR Technical Note, NCAR/TN-242 + STR.
- Thompson, S.L. & Pollard, D. (1995a) A global climate model (GENESIS) with a land–surface-transfer scheme (LSX). Part 1: Present-day climate. *J. Clim.* **8**, 732–761.
- Thompson, S.L. & Pollard, D. (1995b) A global climate model (GENESIS) with a land–surface-transfer scheme (LSX). Part 2: CO₂ sensitivity. *J. Clim.* **8**, 1104–1121.
- Wilson, C.A. & Mitchell, J.F.B. (1987) A doubled CO₂ climate sensitivity experiment with a global climate model including a simple ocean. *J. Geophys. Res.* **92(D11)**, 13,315–13,343.
- Woodward, F.I., Smith, T.M. & Emanuel, W.R. (1995) A global land primary productivity and phytogeography model. *Global Biogeochem. Cycles* (in press).