

A NEW CONIFER HERBARIUM

William B. Critchfield (Berkeley, Calif.)*

At the Institute of Forest Genetics, Placerville, California¹), we have recently organized a specialized herbarium of pines (*Pinus*) and firs (*Abies*). Incorporated in it are the many specimens that have accumulated since the Institute was established in 1925. Like the Eddy Arboretum, the herbarium is an integral part of our research program, which is concerned with the genetic improvement of conifers native to the western United States. Interspecific hybridization and geographic variability are important parts of the program, and the herbarium is particularly useful in documenting this research.

Because we deal exclusively with coniferous materials, our techniques have been adapted to their bulky nature. For mounting specimens, we use herbarium plastic and sheets of high-grade cardboard cut to the standard size of herbarium paper. Most cones, bark, and other unmounted materials are stored in heavy-duty (0.004-inch thick) polyethylene bags. Fir cones are prevented from disintegrating by several interconnected straps of thickened herbarium plastic.

Our general pine collection includes cones, foliage, seeds, and seedlings of most of the 95 to 100 species in this genus — about 3,000 specimens in all. The regions best represented are the western United States, Mexico and Central America, and east and southeast Asia. The extensive collection of Mexican pines includes specimens collected by E. L. Little, Jr., M. Martinez, N. T. Mirov, and the 1962 North Carolina State College expedition to Mexico (publ. 1963). The newer general collection of *Abies* includes about 500 specimens; only the western North American firs are adequately represented in it. Collections by J. L. Hamrick, O. V. Matthews, G. B. Parker, and K. J. Roller are included. Most of the specimens in these general collections were obtained from single trees, but many recent additions are small (5- to 10-tree) mass collections from single stands. The other genera of conifers have only token representation in the herbarium.

One important function of this herbarium is the documentation of our research program. Examples are collections of the many pines whose turpentine has been analyzed by N. T. Mirov (1961), and specimens of the numerous interspecific pine hybrids produced at Placerville during the past 40 years (Little and Righter 1965).

Another major function of the herbarium at Placerville is the preservation of mass collections of conifers. These collections document a number of comprehensive investigations of geographic variability within species or species complexes (table 1). Most of these studies have been the subjects of Ph. D. theses at the University of California. This kind of material, which is much too bulky to be acceptable to most herbaria, occupies more than half of our herbarium. In most instances the samples encompass the range of the species. Growing at Placerville are offspring of *Pinus attenuata*, *P. contorta*, *P. flexilis*, *P. strobiformis*, *P. radiata*, and *P. sabiniana* parent trees included in these mass collections.

* Pacific Southwest Forest and Range Experiment Station, Forest Service, U.S. Department of Agriculture, Berkeley, California. Abbreviation for *Index Herbariorum*: IFGP.

¹) Mail address: 1912 Carson Road, Placerville, California 95667, U.S.A.

Table 1. — Mass collections of pines in the herbarium of the Institute of Forest Genetics, Placerville, California

Species	Estimated number of trees	Kinds of material	Collected or assembled by	Reference
<i>Pinus attenuata</i>	150	C, F ¹⁾	G. B. Newcomb	Newcomb (1962)
<i>P. contorta</i>	400	C, F, L	W. B. Critchfield	Critchfield (1957)
<i>P. elliotii</i>	250	C, S, SD	A. E. Squillace	Squillace (1966)
<i>P. flexilis</i> ²⁾	300	C, F	M. M. Douglass and J. R. Douglass	Douglass (1955)
<i>P. flexilis</i> and <i>P. strobiformis</i> ³⁾	600	C, F	J. W. Andresen	Steinhoff (1964)
<i>P. monticola</i> ⁴⁾	100	C	B. V. Barnes	
<i>P. muricata</i>	150	C, F	J. W. Duffield	Duffield (1951)
<i>P. ponderosa</i>	2,000	C, F, B	R. Z. Callaham	Callaham (1962) Wells (1964)
<i>P. radiata</i>	300	C, F, S, L	M. B. Forde	Forde (1964)
<i>P. sabiniana</i>	150	C, F, L	J. R. Griffin	Griffin (1964, 1965)

¹⁾ C = cones, F = foliage, L = leaf transections, B = bark, S = seeds, SD = seedlings.

²⁾ Mostly Colorado.

³⁾ *P. strobiformis* from U.S. only.

⁴⁾ Elevational transects from northern Idaho.

REFERENCES

- CALLAHAM, ROBERT Z. 1962 — Geographic variability in growth of forest trees. pp. 311-325 in: Tree growth (ed. T. T. Kozlowski). New York: Ronald Press.
- CRITCHFIELD, WILLIAM B. 1957 — Geographic variation in *Pinus contorta*. Harvard Univ., Cabot Found. Publ. 3, 118 pp.
- DOUGLASS, MARGARET M., and DOUGLASS, J. R. 1955 — The distribution and growth of the limber pine in Colorado. J. Colo.-Wyo. Acad. Sci. 4: 46-47.
- DUFFIELD, JOHN W. 1951 — Interrelationships of the California closed-cone pines with special reference to *Pinus muricata* D. Don. Ph. D. thesis, Univ. Calif., Berkeley. 77 pp.
- FORDE, MARGOT B. 1964 — Variation in natural populations of *Pinus radiata* in California. Parts 1-4. New Zeal. J. Bot. 2: 213-257, 459-501.
- GRIFFIN, JAMES R. 1964 — Cone morphology in *Pinus sabiniana*. J. Arnold Arboretum 45: 260-272.
- 1965 — Digger pine seedling response to serpentinite and non-serpentinite soil. Ecology 46: 801-807.
- LITTLE, ELBERT L., JR., and RIGHTER, FRANCIS I. 1965 — Botanical descriptions of forty artificial pine hybrids. U.S. Dep. Agr. Tech. Bull. 1345, 47 pp.
- MIROV, N. T. 1961 — Composition of gum turpentine of pines. U.S. Dep. Agr. Tech. Bull. 1239, 158 pp.
- NEWCOMB, GENE B. 1962 — Geographic variation in *Pinus attenuata* Lemm. Ph. D. thesis, Univ. Calif., Berkeley. 191 pp.
- NORTH CAROLINA STATE COLLEGE. 1963 — Collecting pine material in Mexico for provenance trials and wood studies. N.C. State Coll. Sch. Forest., Forest Tree Impr. Program, Tech. Rep. 18. 23 pp.
- SQUILLACE, A. E. 1966 — Geographic variation in slash pine. Forest Sci. Monogr. 10. 56 pp.
- STEINHOFF, RAPHAEL J. 1964. — Taxonomy, nomenclature, and variation within the *Pinus flexilis* complex. Ph. D. thesis, Mich. State Univ., 81 pp.
- WELLS, OSBORN O. 1964 — Geographic variation in ponderosa pine. I and II. Silvae Genet. 13: 89-103, 125-132.