

FIRE AND BIRDS IN MARITIME PACIFIC NORTHWEST

MARK H. HUFF, NATHANIEL E. SEAVY, JOHN D. ALEXANDER, AND C. JOHN RALPH

Abstract. Resource managers face the challenge of understanding how numerous factors, including fire and fire suppression, influence habitat composition and animal communities. We summarize information on fire effects on major vegetation types and bird/fire relations within the maritime Pacific Northwest, and pose management-related questions and research considerations. Information on how fire affects birds is limited for the maritime Pacific Northwest, even though fire is an essential process within natural vegetation communities throughout the region. We describe fire regimes, vegetation succession patterns, bird communities, and fire effects on birds for 12 major vegetation types in the region. Fire regimes and fire effects vary considerably within this region due to its diverse topography and climate. Seven of the types have a low- to moderate-severity fire regime and five have a high-severity fire regime with fire-return intervals that span several centuries. Bird communities and effects of fire are best known from the western hemlock type, which has a high-severity fire regime. The postfire stand-initiation stage in this type supports a reasonably distinct avifauna compared to other successional stages, a phenomenon that has been documented for high-severity fire regimes in other regions. In general, there is a high turnover of species after high-severity fires, with a shift primarily from canopy-dwelling to ground-, shrub-, and snag-dwelling species that mostly are not associated with other successional stages. No studies exist that directly address how bird communities are affected by habitat changes from fire suppression in this region. The most likely bird communities vulnerable to these changes are in low-severity, high-frequency fire regimes that include the Douglas-fir type, drier portions of the white fir type, Oregon-oak woodlands and savannas, native grasslands and sclerophyllous shrublands. In general, prescribed fire is not being used for bird conservation in this region. Where prescribed fire is being used to restore fire as an ecological process or more often for reducing potentially hazardous fuels, bird conservation objectives can be achieved as a secondary benefit. New land management policies that will greatly accelerate fuel reduction activities throughout the Pacific Northwest, including use of prescribed fire, are currently being undertaken with limited scientific information on the ecological consequences for bird communities.

Key Words: birds, fire, fire-suppression, forest management, Pacific Northwest, succession.

EL FUEGO Y LAS AVES EN LA PORCIÓN MARÍTIMA DEL NOROESTE

Resumen. Los manejadores de recursos naturales, enfrentan el reto de entender como numerosos efectos, incluyendo incendios y supresión de estos, influyen en la composición del habitat y sus comunidades de animales. Resumimos información de los efectos del fuego en la mayoría de los tipos de vegetación y las relaciones ave-incendios alrededor de las zonas marítimas de la costa del Noroeste pacífico, y postulamos preguntas y consideraciones de investigación relacionadas al manejo. La información de cómo el fuego afecta a las aves es limitada en la región marítima del Noroeste Pacífico, a pesar de que el fuego es un proceso esencial para las comunidades vegetales naturales de dicha región. Describimos regimenes de incendios, patrones de sucesión vegetal, comunidades de aves y los efectos del fuego en las aves, en 12 tipos principales de vegetación en dicha región. Los regimenes y los efectos del fuego varían considerablemente en esta región, debido a la diversidad topográfica y climática. Siete de los tipos tienen un régimen de severidad de incendios de bajo a moderado, y cinco tienen un régimen de severidad alto, con intervalos de repetición de incendios separados por varios siglos. Las comunidades de aves y sus efectos al fuego son mejor conocidos para el tipo de bosque occidental de abeto, el cual tiene un alto régimen de severidad de incendios. El estado de iniciación post incendio de este tipo, soporta a una avifauna razonablemente distinta comparado con otros estados sucesionales, fenómeno el cual ha sido documentado por regimenes severos de incendios altos en otras regiones. En general, existe una alta recuperación de las especies después de incendios con regimenes de alta severidad, con un cambio primordialmente de especies que viven en las copas de los árboles contra las del suelo, arbustos, y tocones, las cuales principalmente no están asociadas con otro estado sucesional. No existen estudios en esta región los cuales muestren directamente cómo las comunidades de aves son afectadas por cambios en el habitat, ocasionados por supresión del fuego. Las comunidades de aves más vulnerables a estos cambios se encuentran en el tipo de baja severidad, con regimenes de incendio de alta frecuencia, los cuales incluye el tipo *Pseudotsuga menziessi*, porciones más secas del tipo de Abies blanco, bosques de encino y sabana de Oregon, en pastizales nativos y en arbustos sclerophyllus. En general, las quemadas prescritas no son utilizadas para la conservación de aves en esta región. Las quemadas prescritas son utilizadas para restaurar, como un proceso ecológico

más a menudo, para reducir potencialmente combustibles riesgosos. Al restaurar con fuego, objetivos para la conservación de aves pueden ser alcanzados, como beneficio secundario. Nuevas políticas de manejo de la tierra en la región del Noroeste Pacífico acelerarán enormemente actividades de reducción de combustible, incluyendo el uso de quemadas prescritas, las cuales son actualmente aplicadas con información científica limitada en cuanto a las consecuencias ecológicas para las comunidades de aves.

Disturbances can modify physical and biological environments and have profound effects on ecological processes, patterns, and interactions (e.g., White 1979). Primary disturbance agents include fires, wind, insect and disease outbreaks, floods, landslides, and human-related activities (Pickett and White 1985). In the Maritime Pacific Northwest, fires have been the most wide-ranging and continuous disturbance agent (Agee 1993), except for forest harvest, over the last few decades. Fire regimes and fire effects vary considerably within this region, primarily due to a diverse climate and topography that ranges from arid lands to rainforest and sea level to mountain peaks >4,300 m. Fire severities range from slight to cataclysmic, and natural fire-return intervals from almost annually to >1,000 yr (Agee 1993). In general, the north and coastal environments of western Washington and northwest Oregon have infrequent, stand-replacing fires that are part of a high-severity fire regime. In southwest Oregon and northwest California, and on the east side of the Cascade Mountains, fire occurs more frequently, a low- to moderate-severity fire regime, often with effects that are less severe, but more variable.

With effective fire suppression that began 80–100 yr ago, the natural patterns of fire have changed, especially in areas where fires burn most frequently (Agee 1993). In areas of low- to moderate-severity fire regimes, effective fire prevention may change habitat composition, shift the composition of biological communities, and lead to unnatural fuel accumulations associated with severe fires that cannot be withstood by historical ecosystems. When fire suppression increases fuel loads and the risk of severe fires, it becomes difficult to address the social and ecological concerns to protect property and lives, sensitive species and their habitat, and air and water quality, thus reinforcing widespread suppression of fires. However, consequences of continuing to suppress fires (i.e., passive management) without corrective measures also are high (Agee 2002). Shifts in fire prevention strategies are underway that propose to ameliorate potentially hazardous fuel conditions in areas of low- to moderate-severity fire regimes by steadily increasing prescribed fires (a fire ignited under known fuel conditions, weather, and topography to achieve specified objectives); thinning tree canopies to create shaded fuel breaks (Agee *et al.*

2000); and introducing other fuel reduction activities as part of a revised National Fire Plan (USDI *et al.* 2001). Such actions, however, often are planned and implemented with minimal understanding or considerations for effects on biota. Ecological objectives, if stated in planning documents, are almost always secondary to those for reducing hazardous fuels.

The purpose of this paper is to summarize information on bird-fire relations within the Maritime Pacific Northwest (hereafter Pacific Northwest), an area that encompasses the east slope of the Cascade Ranges, and the western portions of Washington, Oregon, and northern California (Fig. 1). This area is roughly equivalent to the range of the Northern Spotted Owl (*Strix occidentalis caurina*), the Northwest Forest Plan (USDA and USDI 1994b), and similar to the Southern Pacific Rainforest and Cascade Mountains physiographic stratifications used for the Breeding Bird Survey (Droege and Sauer 1989; map at <http://www.mbr-pwrc.usgs.gov/bbs/physio.html> [23 July 2004]). Our paper is organized into four parts: (1) a description of environmental conditions, vegetation communities, the role of fire in the major habitats and bird communities of the Pacific Northwest, and a summary of information on the effects of fire on birds; (2) a discussion of the major alterations to low- and moderate-severity fire regimes and their known or hypothesized effects on birds in this region; (3) a discussion of the role of prescribed fire in low- and moderate-severity fire regime and the implications for bird conservation; and (4) the critical management issues and research questions for this region.

ENVIRONMENTAL SETTING AND ROLE OF FIRE

FIRE REGIMES

Fire regimes have been described for the Pacific Northwest forests by Agee (1981, 1990, 1993, and 1998). Fires in high-severity fire regimes usually occurred >100 yr apart with >70% of the vegetative basal area removed; in moderate-severity fire regimes, fires were 25–100 yr apart with 20–70% basal area removed; and in low-severity fire regimes, they averaged 1–25 yr apart with <20% of the basal area removed. Grasslands and shrublands typically are part of a high severity regime, in which fires are




FIGURE 1. Generalized distributions of major vegetation types in the maritime Pacific Northwest (adapted from Kuchler 1964). Two types displayed on the map, ponderosa pine and chaparral (California and montane types combined), are not covered in this chapter because they have large geographic distributions primarily outside the maritime Pacific Northwest.

considerably more frequent than in most forested areas in this regime.

Vegetation patterns after fire depend on fire severity (Agee 1998). Agee found that for the low-, moderate-, and high-severity fire regimes, respectively, burn patch size created by fire tends to be small (~1 ha), medium (up to 300+ ha), and large (up to 10,000+ ha). He also found that the amount of patch edge, or contrasting conditions created by fire, tends to be low, high, and moderate, according to fire regime.

In high-severity fire regime areas of the Pacific Northwest, post fire stand-initiation can be prolonged for many decades as trees reestablish slowly after a fire (e.g., Hemstrom and Franklin 1982, Huff 1995). After burning or logging, this stage can be hastened through replanting trees and suppressing competing vegetation. Once trees establish in high-severity fire regime areas, the stem-exclusion and re-initiation of understory stages may continue for a century or

two, leading to an old-growth stage that can persist for centuries depending on the natural fire rotation. Moderate-severity fire regimes sustain a highly variable forest structure by creating stands of uneven size and age trees and patchiness at landscape scales. Of the three fire regimes, low severity is most likely to create a balanced tree age-class distribution, where each fire only affects pattern and process on a small portion of a landscape (Agee 1998).

ECOLOGICAL UNITS

Geographic distribution of vegetation and fire regimes in the Pacific Northwest are closely linked to complex topographic, moisture, and temperature gradients that can change rapidly with elevation, latitude, geological formations, substrate, and proximity to the Pacific Ocean (Agee 1993). In classifying the environments of Washington, Oregon, and California, Franklin and Dyrness (1973) and Barbour and Major

(1977) divided the states into broad ecological units using two approaches: (1) physiographic provinces based on geography, geology, and soils, and (2) vegetation zones (hereafter types) based on associations of natural plant communities.

In this paper, we use the physiographic area delineated in the Northwest Forest Plan (Forest Ecosystem Management Team 1993, USDA and USDI 1994a, 1994b, 2000) (Fig. 1) to describe distributions of broad vegetation types, fire regimes, and avifauna (Table 1). We aggregated vegetation types into five major ecosystems: lowlands and foothills, coastal forests, lower montane, upper montane, and subalpine. We describe the distribution of 12 vegetation types (Fig. 1), fire regimes, and establishment patterns after fire, using Franklin and Dyrness (1973, 1988) (vegetation of Washington and Oregon), Barbour and Major (1977) (vegetation of California), and Agee (1993) (fire regimes and effects) as our initial source. About 59% of the Pacific Northwest is covered by a high-severity fire regime and about 41% by a low- or moderate-severity fire regime based on vegetation types and fire regimes in Table 1 and Fig. 1. Below we describe the major vegetation types within the five ecosystems.

Lowlands and Foothills Ecosystem

This ecosystem occurs in a relatively dry environment within and near interior valley bottomlands.

The diverse vegetation within the lowlands and foothills is aggregated into one vegetation type, interior valley. It is a mosaic of grasslands (e.g., Johannessen *et al.* 1971, Franklin and Dyrness 1973) and oak savannas, oak woodlands, mixed oak-conifer forests (e.g., Habeck 1961, Thilenius 1968, Smith 1985, Riegel *et al.* 1992); and sclerophyllous shrublands (i.e., chaparral) dispersed sporadically across southwestern Oregon and northern California (Whittaker 1954, Barbour and Major 1977). Generally, fire history has not been well documented in the lowlands and foothills because fires carried by grass and herbs are short duration, low intensity, and high-severity in which most vegetation is consumed and because much of the vegetation has been converted to other uses, such as urban and suburban, cropland, pasture, and forestry.

Oak woodlands and savannas are dominated by Oregon white oak (*Quercus garryana*) in Oregon; California black oak (*Quercus kelloggii*) is an important species in the southern portion of the lowlands and foothills. At present, the ground flora has been so altered in these communities, especially through livestock grazing, that benefits of using fire for restoration are uncertain. Sprouting appears to be the primary process for recruitment. The role fire plays in perpetuating these communities by stimulating sprouts or influencing acorn germination and seedling survival is not well understood (Harrington and Kallas 2002). Sclerophyllous shrublands,

TABLE 1. FOREST VEGETATION TYPES OF THE MARITIME PACIFIC NORTHWEST BY FIVE ECOSYSTEM TYPES AND BY MOISTURE AND FIRE REGIMES.

Ecosystem type	Forest vegetation types	
	Wet to mesic environments/ high-severity fire regime	Dry to mesic environments/ low- to moderate-severity fire regime
Lowlands and foothills		Interior valley—a mosaic of oak woodlands and savannas; oak-conifer; grasslands ^a , and sclerophyllous shrublands ^a .
Coastal forests	Sitka spruce	Coast redwoods.
Lower montane	Western hemlock	Douglas-fir. Mix evergreen hardwood. White fir. Grand fir. (Ponderosa pine ^b). (California and montane chaparral ^{a,b}).
Upper montane	Pacific silver fir	Shasta red fir.
Subalpine	Subalpine fir Mountain hemlock	

^a Occurs in dry to mesic environments and has a high-severity fire regime.

^b Ponderosa pine and California and montane chaparral types were not covered in this chapter because they have large geographic distributions primarily outside the maritime Pacific Northwest.

established near grasslands and oak communities in valley bottoms and serpentine soils, are maintained by fire, and dominated mostly by shrubs, such as *Ceanothus* spp., *Arcostaphyllum* spp., and *Baccharis* spp. High-severity fires are common, and burned areas are readily re-colonized by these shrub species when they resprout and their long-lived seeds quickly respond to scarification from burning. These shrublands appear to be declining in the Pacific Northwest because human development and fire suppression favor tree establishment (Chappell and Kagan 2000).

Coastal Forests Ecosystem

Two major vegetation types occur within the coastal forests, Sitka spruce (*Picea sitchensis*) and coast redwood (*Sequoia sempervirens*), hug the coastline and extend up major river valleys inland where summer fog lingers. Important tree species found in the Sitka spruce type are Sitka spruce western hemlock (*Tsuga heterophylla*) and western red cedar (*Thuja plicata*); grand fir (*Abies grandis*) and Douglas-fir (*Pseudotsuga menziesii*) are minor components (Fonda 1974, Henderson et al. 1989). Fire history has not been documented for this type (Agee 1993). The climatic conditions and the extent of older age classes indicate that it burned infrequently (likely even less often than the wettest areas of the western hemlock type), and belongs to a high-severity fire regime. Fire is more likely to spread into the coastal Sitka spruce type from nearby areas within the lower montane western hemlock type by the occasional dry east winds during rare climatic conditions, than from fires that originate from within the type (Agee 1993). Because fire occurs so infrequently, wind is probably a more important disturbance agent in this type.

Coast redwood, the tallest tree the world, is the principal species of the coast redwood type (Waring and Major 1964, Zinke 1977, Noss 1999). Other important trees within the type's broad moisture gradient include western hemlock, western red cedar, Sitka spruce, tan oak (*Lithocarpus densiflorus*), and Douglas-fir. The type's fire regime is moderate severity, with fires mostly occurring as low-to-moderate severity with infrequent high-severity events (Veirs 1982; Jacobs et al. 1985; Stuart 1987; Finney and Martin 1989, 1992). Fire-return intervals may exceed 500 yr in moist areas, and be as low as 20–50 yr at drier inland locations. Structurally diverse, multi-aged forests maintained by fires are characteristic of the type.

Lower Montane Ecosystem

This is the most diverse ecosystem type and covers the largest geographic area. We recognize five major vegetation types within the lower montane ecosystem: western hemlock; Douglas-fir; mixed evergreen hardwood; white fir; and grand fir. In the western hemlock type, which encompasses the largest area, western hemlock and Douglas-fir commonly co-occur, with the hemlock more prevalent at the wet end and the fir at the dry end of the moisture gradient (Franklin and Dyrness 1973, Zobel et al. 1976, Topik et al. 1987, Topik 1989, Henderson et al. 1989, Hemstrom and Logan 1986, Ruggiero et al. 1991, Atzet et al. 1996). Other important species are Sitka spruce in the coastal areas and Pacific silver fir (*Abies amabilis*) within a wide band of vegetation that transitions into the upper montane. Fire frequency is variable across the type, reflecting diverse moisture and temperatures conditions found within the type. In the areas of high moisture fires burn very infrequently, with 250 to >500-yr return intervals (Hemstrom and Franklin 1982, Agee and Flewelling 1983, Agee and Huff 1987, Henderson et al. 1989, Agee 1991a, Huff 1995, Impara 1997, Agee and Krusemark 2001). A high-severity fire regime predominates throughout much of the type. Fires often create large patches of killed trees with a few very widely scattered Douglas-firs that survived, as well as, an occasional small island of trees that was spared. Trees may reestablish slowly in these burned patches, and may remain a forest opening for many decades. In the drier southern portion of the type, average fire-return interval drops quickly from about 200 yr down to 100–30 yr as the fire regime grades into a moderate severity (Means 1982, Teensma 1987, Morrison and Swanson 1990, Cissel et al. 1998, Weisberg 1998, Van Norman 1998, Olson 2000, Weisberg and Swanson 2003). In these more fire-prone areas, overlapping fires of varying severities create a complex age-class structure. Western hemlock (in wetter environments), Douglas-fir, and western red cedar are early seral species that attain great sizes, dominate for centuries, and, in late-successional conditions, show substantial structural diversity. In mesic and drier environments, Douglas-fir dominates after fire, sometimes in pure stands.

The Douglas-fir type usually occupies the warmest and driest environments in the lower montane (Franklin and Dyrness 1973; Atzet et al. 1992, 1996). It is often found upslope and adjacent to types dominated by ponderosa pine (*Pinus ponderosa*) or oaks. Douglas-fir is commonly an early and late-successional species throughout the Douglas-fir type,

with ponderosa pine distributed widely in dry and warm environments. Other locally important species are lodgepole pine (*Pinus contorta*), western larch (*Larix occidentalis*), Jeffrey pine (*Pinus jeffreyi*), and incense-cedar (*Libocedrus decurrens*). A low-severity fire regime is common throughout the type; characteristics of a moderate-severity regime emerge as moisture increases (Agee 1990, 1991b; Chang 1996; Skinner and Chang 1996; Taylor and Skinner 1997; Everett *et al.* 2000). Historically, fires probably burned often over relatively small areas and were confined mostly to the understory. Infrequent, larger fires that burned with varied severity eclipsed effects of these smaller fires. Fire history studies from across the region indicate that frequencies in dry locations averaged <10 yr but ranged up to 50 yr with increasing moisture. In this type, fire severity varied considerably by topographic position. Here, the most severe fires occur on upper slopes, ridge tops, and south- and west-facing slopes. These fires resulted in simple forest structures, whereas lower slope positions and riparian areas with less severe fires created complex forest structures, including the largest trees (e.g., Taylor and Skinner 1998).

The mixed evergreen hardwood type has the most restricted distribution in the lower montane ecosystem, but the most diverse tree composition (Whittaker 1960, 1961; Sawyer *et al.* 1977; Atzet *et al.* 1992, 1996). The type is most prominent in the coastal mountains and at mid slope and elevation, and moderate aspects. Important tree species are Douglas-fir, usually an overstory dominant, and tan oak. Tan oak has increased its density and cover throughout the type, presumably a result of fire suppression (Atzet *et al.* 1992). Other major species are Pacific madrone (*Arbutus menziesii*) and ponderosa pine.

The few fire-related studies in the mixed evergreen hardwood type show a wide range of fire severities, frequencies, and sizes (Thornburgh 1982, Wills and Stuart 1994, Agee 1991b), indicating that the type largely belongs in a moderate-severity fire regime. The mean fire-return intervals in pre-settlement times cluster around 15–35 yr, but range from 3 to >70, and the size of some fires have been quite large. Complex successional patterns have developed that parallel the variable nature of fire in this type. In general, after low- to moderate-severity fires, tanoak regenerates beneath a canopy of Douglas-fir. After more severe fires, Douglas-fir regenerates beneath a canopy of tanoak, a species that can sprout profusely after fire and have rapid early growth. Older forests usually have one to three age classes resulting from past fires, with Douglas-fir as an emergent canopy above various hardwoods, mostly evergreen.

The white fir type occurs in a range of environments wider than all other types in the Lower Montane except western hemlock (Rundel *et al.* 1977; Sawyer *et al.* 1977; McNeil and Zobel 1980; Atzet *et al.* 1992, 1996). White fir (*Abies concolor*), the most dominant tree species, can form pure stands in the coolest environments. The importance of Douglas-fir, relative to white fir, increases with increasing dryness and temperature. Ponderosa pine becomes a major species in drier environments. The white fir type typically exhibits a diverse and lush understory; yet, the fire regime likely fits a moderate or moderate-to-low severity. Documented mean fire-return intervals of ~10–65 yr and up to 160 yr (McNeil and Zobel 1980, Bork 1985, Agee 1991b, Stuart and Salazar 2000), broadly overlap other lower montane vegetation types in southwest Oregon and northwestern California, yet the type has not been studied extensively considering its breadth of environments. Fires are less frequent at higher elevations where white fir forms pure or nearly-pure stands. White fir has increased in cover and density with decades of fire suppression. In the white fir type, fires typically overlap to create a patchy mosaic of multiple structures and age classes, providing conditions for white fir to persist.

The grand fir type is found at mid-slopes in eastern Washington and Oregon on moist to dry sites (Hopkins 1979, Topik *et al.* 1988, Topik 1989, Lillybridge *et al.* 1995). In addition to grand fir, major tree species include ponderosa pine in warm and dry locations, lodgepole pine in cool and dry, Douglas-fir in environments broadly across the type, and western larch in the northern reaches of the type. Information on fire frequency and effects is scarce from this type (Agee 1993). Behavior of natural fires can be quite unpredictable, varying quickly from intense crown fires to surface fires. Higher severity fires may be an important part of forest development in the grand fir type, but to what degree is unclear. Preliminary indications are that a moderate-severity fire regime might be more likely than a low- or high-severity fire regime. Forest succession after fire should reflect fire-severity patterns and species present at the time of the fire: frequent low-to-moderate severities would favor, if present, Douglas-fir, ponderosa pine, and western larch survival and regeneration, while high severities would favor establishment of lodgepole pine in areas that are cool and dry. Grand fir can establish in open to partially shaded environments with moderate moisture, probably over a wide range of fire severities with low to moderate frequencies (Hall 1983).

Upper Montane Ecosystem

The vegetation types of the upper montane are found upslope from the lower montane, as conditions become cooler and wetter at higher elevations. There are two major upper montane vegetation types in the Pacific Northwest, Shasta red fir (*Abies magnifica*) and Pacific silver fir. The Shasta red fir type occupies a narrow ~500 m elevation band, typically above the white fir type and below the mountain hemlock type where Shasta red fir is common in the overstory and understory in a broad range of environments (Rundel et al. 1977; Sawyer et al. 1977; Atzet et al. 1992, 1996). Other important tree species include lodgepole pine and western white pine (*Pinus monticola*). Studies from Shasta red fir type and California red fir type (adjacent to the Pacific Northwest region), indicate that mean fire-return intervals are about 20–50 yr, with ranges of ~5–65 yr and fire-free periods spanning ~150 yr (McNeil and Zobel 1980, Pitcher 1987, Taylor and Halpern 1991, Agee 1993, Taylor 1993, Chappell and Agee 1996, Bekker and Taylor 2001, Taylor and Solem 2001). The type has a moderate-severity fire regime that creates a very patchy environment of diverse patch sizes and stand structures in close proximity that range from closed- and open-canopied late-successional forests to young, regenerating stands and open meadows.

In the Pacific silver fir type, Pacific silver fir is the dominant species; western and mountain hemlocks (*Tsuga mertensiana*) co-dominate at the lower and upper limits of the type, respectively (Fonda and Bliss 1969, Franklin and Dyrness 1973, Hemstrom et al. 1982, Packee et al. 1983, Brockway et al. 1983, Franklin et al. 1988, Henderson et al. 1989). Other important tree species are Douglas-fir and noble fir (*Abies procera*) in relatively dry and warm climates, Alaska-cedar (*Chamaecyparis nootkatensis*) in cooler and wetter sites, and western white pine in moderate environments. Subalpine fir (*Abies lasiocarpa*) is an important species in the eastern Cascades where the Pacific silver fir type transitions into a subalpine ecosystem. The type has a high-severity fire regime, in which fires burn infrequently (Agee 1993). Although fire history data are scant for this type, known fire-return intervals were about 100–200 yr along the drier eastern edge of the type that is influenced by a continental climate (Agee et al. 1990), and about 300–550 yr in the moister westside of the Cascades (Hemstrom and Franklin 1982). Pre-settlement fires in this type were usually associated with large fires that swept through adjacent types, creating large patches where most of the

vegetation in the Pacific silver fir type was killed. This type, however, can act as a barrier to fire spread, unless extreme conditions associated with prolonged drought and dry east winds are met. Pacific silver fir, a fire-sensitive species, rarely survives where fires have burned and postfire conditions are often too harsh for it to establish, except in very cool and wet locations. Stand establishment after fire can last for decades, and be very prolonged if seed sources are absent for species that function as pioneers, such as Douglas-fir, western white pine, noble fir, subalpine fir, and lodgepole pine. Pacific silver fir is more prominent as forests mature.

Subalpine Ecosystem

Two major vegetation types occur in subalpine ecosystems: subalpine fir and mountain hemlock. Subalpine fir dominates throughout the subalpine fir type and forms nearly pure stands (Franklin and Mitchell 1967; Fonda and Bliss 1969; Henderson 1973, 1982; Agee and Kertis 1987; Franklin et al. 1988; Henderson et al. 1989; Lillybridge et al. 1995). Other species that typically co-occur with subalpine fir are Pacific silver fir in transition upslope into the type; whitebark pine (*Pinus albicaulis*) at higher elevations; Engelmann spruce (*Picea engelmannii*) as the type transitions to the east and lodgepole pine in harsh environments and locations with higher fire frequencies. A high-severity fire regime characterizes the subalpine fir type (Fahnestock 1977, Agee and Smith 1984, Agee et al. 1990, Taylor and Fonda 1990, Huff and Agee 1991, Agee 1993). Fires in this type typically burn infrequently, though more often than in the mountain hemlock type because precipitation and snow pack are lower and summer months are warmer. The fire-return interval ranges about 100–250 yr from relatively dry to wet environments. Tree reestablishment can be slow and inconsistent (except where lodgepole pine is prevalent) due to severe climatic and site conditions for establishment and insufficient distribution of seed, leaving the postfire environment in a park-like setting for decades.

In the mountain hemlock type, mountain hemlock tends to occur in mixed stands with Pacific silver fir as a co-dominant, and with subalpine fir, lodgepole pine, and Alaska-cedar (Fonda and Bliss 1969; Henderson 1973, 1982; Sawyer et al. 1977; Agee and Kertis 1987; Franklin et al. 1988; Henderson et al. 1989; Atzet et al. 1992, 1996). Fire history is poorly understood in the mountain hemlock type; however, it is likely that fires burn very infrequently, suggesting a high-severity fire regime (Agee 1993, Dickman

and Cook 1989). Fire-return interval for similar mountain hemlock forests in nearby Canada were found to be >1,000 yr (Lertzman and Krebs 1991). All major tree species in this type are easily killed by fire, although older mountain hemlocks can survive with fire scars where fires burned at lower intensities. Stand establishment after fire is slow, inhibited by harsh climatic conditions and seeds that are poorly dispersed (Agee and Smith 1984).

FIRE AND BIRD COMMUNITIES IN THE PACIFIC NORTHWEST

Computer simulations of Pacific Northwest forests have demonstrated that over the last 3,000 yr, historical fire regimes maintained highly variable forest age-class distributions. Wimberly *et al.* (2000) estimated that the proportion of older forest age classes varied from 25–75% at various times during this period. Such fire-related fluctuations in habitat availability have important implications for understanding the dynamics of bird populations across the Pacific Northwest. We took a two-step approach to describe interactions between fire and birds. First, we identified bird species that were common and indicative of each vegetation type. When literature was not available for a specific vegetation type, we used general sources that synthesize bird distribution and habitat relations in the Pacific Northwest (e.g., Johnson and O'Neil 2001). Second, we characterized the responses of birds to short- and long-term effects of fire using existing literature from the Pacific Northwest; however bird-related information was not available for many of the vegetation types.

Lowlands and Foothills Ecosystem

Bird communities of the interior valley vegetation type are difficult to characterize because vegetation prior to European settlement has been dramatically altered over the last 150 yr. Contemporary species' habitat relations in natural dry grasslands or prairies and oak savannas are based on reports from agriculturally dominated landscapes, where only small, scattered remnants of original habitat remain. Using natural history information and historical accounts, Altman *et al.* (2000) developed a list of bird species historically associated with dry grassland and oak savanna in this region. They list six species as highly associated (abundance is likely to be significantly higher in this habitat during the breeding season than elsewhere in the Pacific Northwest) with dry grasslands and oak savannas, including the: Burrowing Owl (*Athene cunicularia*) (probably

extirpated from the region), Western Meadowlark (*Sturnella neglecta*), Vesper (*Pooecetes gramineus*) and Grasshopper Sparrows (*Ammodramus sarnanarum*), Horned Lark (*Eremophila alpestris*), and Western Bluebird (*Sialia mexicana*). Most of these species are not widely distributed elsewhere within the Pacific Northwest. No information was available regarding fire effects on birds inhabiting native grassland and savanna habitats.

To document possible long-term changes in oak woodland bird communities where fire was excluded, Hagar and Stern (2001) re-sampled three of Anderson's (1970) oak woodland bird study sites about three decades later. Results of both studies were expressed in terms of density (birds 100 acres⁻¹). Hagar and Stern (2001) found 14 species that were rare or absent in 1967–1968 but more abundant in 1994–1996, and three species detected in 1967–1968 that were absent in 1994–1996. These authors hypothesized that fire suppression had allowed more coniferous trees and a thicker understory to develop, favoring birds that the authors characterized as forest-dwelling species (e.g., Swainson's Thrush [*Catharus ustulatus*], Purple Finch [*Carpodacus purpureus*], House Wren [*Troglodytes aedon*], and Pacific-slope Flycatcher [*Empidonax difficilis*]) over birds favoring partially open habitat (e.g., Chipping Sparrow [*Spizella passerina*], Bushtit [*Psaltriparus minimus*], and Yellow Warbler [*Dendroica petechia*]).

Bird populations closely linked to oaks could be in jeopardy if fire exclusion continues. Oak-dominated woodlands of the Pacific Northwest are maintained by fire (Agee 1993), which has been controlled tenaciously because oaks are in close proximity to rural communities. Because fire exclusion in this vegetation has caused major structural and compositional changes (e.g., a shift to dominance by conifers) (Tveten and Fonda 1999), thus potentially jeopardizing bird populations closely linked to oaks. Bird species most likely to be affected by declining oak habitat resulting from fire suppression are White-breasted Nuthatch (*Sitta carolinensis*), Western Scrub-Jay (*Aphelocoma californica*), and Acorn Woodpecker (*Melanerpes formicivorus*), and in southern Oregon and northern California the Ash-throated Flycatcher (*Myiarchus cinerascens*) and Oak Titmouse (*Baeolophus inornatus*; Altman *et al.* 2000).

Western Scrub-Jay, California Towhee (*Pipilo crissalis*), and Lesser Goldfinch (*Carduelis psaltria*) appear to be the most common species in fire-dependent sclerophyllous shrublands of the interior valley type, based on informal field observations (Altman *et al.* 2000). These shrublands support a unique assem-

blage of bird species that seldom breed elsewhere in the Pacific Northwest, including Wrentit (*Chamaea fasciata*), Green-tailed Towhee (*Pipilo chlorurus*), and Fox Sparrow (*Passerella iliaca*). The effects of fire on this vegetation type probably are similar to other high-severity fire regimes where most of the aboveground vegetation is killed, although the amount and distribution of mortality can vary considerably within and among fires. Studies from burned and unburned shrublands suggest that after fire, bird occupancy shifts to species that appear to favor open environments, and that shrub-dwelling species persist at moderately to substantially lower populations or leave and re-colonize later when the shrub cover is suitable (e.g., Moriarty et al. 1985).

Coastal Forests Ecosystem

The bird community of coastal forest types are similar, and include: Olive-sided Flycatcher in early seral with residual trees; Hutton's Vireo in shrub with small tree regeneration; Winter Wren (*Troglodytes troglodytes*), Wilson's Warbler (*Wilsonia pusilla*), Pacific-slope Flycatcher, and Hermit Warbler (*Dendroica occidentalis*) in young and mature forest; Varied Thrush (*Ixoreus naevius*) and Pileated Woodpecker (*Dryocopus pileatus*) in mature forest; Red Crossbill (*Loxia curvirostra*) in mature and old-growth forest; and Vaux's Swift (*Chaetura vauxi*) in old-growth forests with snags (Oregon-Washington Partners in Flight (PIF) <http://community.gorge.net/natres/pif.html>). Also, the federally listed Northern Spotted Owl breed in these types, as well as federal species of concern including Olive-sided Flycatcher (USDI Fish and Wildlife Service 2002).

Historically, fire played an important role in maintaining a mosaic of seral stages or habitat structures (e.g., snags) that have facilitated the persistence of these species. The Sitka spruce and wetter portions of the western hemlock types have high-severity fires and birds probably respond to fire similarly in these types (see section on *Lower Montane Ecosystem*). Fire suppression probably has had little effect on bird communities in the Sitka spruce type because the fire-return interval is so long. The direct effects of fire on coast redwood forest bird communities are not well documented. Where fires are infrequent in the redwoods, fire is probably less important than other disturbances, such as wind-throw. In areas with frequent low to moderate severity fires, such as drier coastal forest sites, fire played a role in maintaining forest structure by limiting the amount of downed woody debris, or reducing understory vegetation.

Lower Montane Ecosystem

Bird communities have been studied extensively in the western hemlock type (e.g., Morrison and Meslow 1983a and 1983b, Manuwal and Huff 1987, Carey et al. 1991, Gilbert and Allwine 1991, Huff and Raley 1991, Manuwal 1991, McGarigal and McComb 1995, Bettinger 1996, Chambers et al. 1999). The most widespread and abundant bird species of older forests (>200 yr after fire) are Chestnut-backed Chickadee (*Poecile rufescens*), Pacific-slope Flycatcher, Winter Wren, Hermit Warbler, and Golden-crowned Kinglet (*Regulus satrapa*). Other important species are Varied Thrush (wet environments), Wilson's Warbler (with increasing importance of deciduous trees and shrubs), Red-breasted Nuthatch (*Sitta canadensis*; drier environments), and Northern Spotted Owl—for which a large proportion of its habitat lies within the western hemlock type (Bart et al. 1992).

In the wet Olympic Mountains where fires are rare, breeding bird communities were examined at two sites covering a long stand initiation stage, 1–3 and 19 yr after fire, and compared to surveys from the other successional stages (Huff 1984, Huff et al. 1985). Winter Wren and Dark-eyed Junco (*Junco hyemalis*) were the most abundant species during the first three years after fire. At year 19 after fire, Winter Wren abundance was 3–4 times below the three most abundant species, Dark-eyed Junco, Rufous Hummingbird (*Selasphorus rufus*), and American Robin (*Turdus migratorius*). Year 19 had the highest species richness and diversity, including the highest amount of woodpecker species. In general, the stand initiation stage was most favorable for ground- and brush-foraging species, while unfavorable for canopy-feeding species. A high proportion of species bred only in the stand initiation stage after fire (30%), as observed in other regions (e.g., Taylor and Barmore 1980). Pacific-slope Flycatcher, abundant in older forests, was negatively affected by fire during the stand initiation stage; more so than any other species (Huff et al. 1985).

In a high severity fire regime, wildfires and timber harvest followed by site preparation with fire have a few similar effects on bird habitat: loss of live tree overstory followed by re-colonization of herbs, shrubs, and trees. After logging followed by broadcast burning in the western hemlock type in Oregon, the most common birds in open managed stands (<15 yr-old and planted with conifer regeneration) were mostly ground- and brush-foraging species that included White-crowned Sparrow (*Zonotrichia leucophrys*) and Song Sparrow (*Melospiza melodia*), Swainson's

Thrush, Rufous Hummingbird, Spotted Towhee (*Pipilo maculatus*), American Goldfinch (*Carduelis tristis*), Dark-eyed Junco, and MacGillivray's (*Oporornis tolmiei*) and Orange-crowned Warbler (*Vermivora celata*; Morrison and Meslow 1983b, Bettinger 1996). In contrast to the stand initiation stage after a fire (e.g., Wyoming, Taylor and Barmore 1980; Washington, Huff 1984), primary and secondary cavity nesting birds were nearly absent in logged areas because past practices in this region typically removed most or all standing dead trees. Hermit Warbler, Chestnut-backed Chickadee, Golden-crowned Kinglet, Swainson's Thrush and Dark-eyed Junco that are common in young-tree thickets of rapidly maturing managed forests ~15–35 yr old (Bettinger 1996), are probably associated with similar conditions after wildfires but take longer to develop. About 80 yr after wildfires, bird species composition in the western hemlock type tends to stabilize, (i.e., Huff and Raley 1991), that is, generally not changing until the next wildfire, which could be centuries. During this extended period between fires, relative abundance among species appears to be regionally distinct and varies as forests grow older (Huff and Raley 1991). Bark foragers, such as Brown Creeper (*Certhia americana*), Hairy Woodpecker (*Picoides villosus*), Pileated Woodpecker, and Red-breasted Nuthatch tend to increase overtime as forests develop old-growth characteristics.

Relative to other lower montane forest types, the bird community of mixed evergreen hardwood type had more insectivorous species, foliage gleaning species, and snag nesting species (Alexander 1999). The mixed evergreen hardwood type also provides habitat for the federally listed Northern Spotted Owl and federal species of concern such as Olive-sided Flycatcher (USDI Fish and Wildlife Service 2002).

No studies have directly measured the response of mixed evergreen hardwood type bird communities to fire. Like the coastal forests, bird communities in this type vary among forest age classes (Raphael *et al.* 1988, Ralph *et al.* 1991). However, unlike the coastal forests, fire was probably far more widespread than any other disturbance. As a result, most authors agree that the historical fire regime has created and maintained high spatial and biological heterogeneity of vegetation in this type (Wills and Stuart 1994, Agee 1991b, 1993). Therefore, fire likely played an important role in maintaining the richness and diversity of these bird communities. Specifically, fires may influence communities by maintaining a heterogeneous seral composition, creating snags for foraging and nesting, and increasing availability of limiting food resources. The relative importance

of these mechanisms is likely to be highly variable among species.

Upper Montane and Subalpine Ecosystems

The bird community of upper montane Shasta red Fir type is less species rich than those in Lower Montane Ecosystem types (Alexander 1999). The federally listed Northern Spotted Owl use this type, as do species of concern such as Olive-sided Flycatcher (USDI Fish and Wildlife Service 2002).

In the Shasta red fir type, the relatively mixed effects of fire may be important for understanding the composition of the bird community. Red-breasted Nuthatch and Golden-crowned Kinglet, both strongly associated with this vegetation type, decrease in response to fire (Kreisel and Stein 1999) and management practices that reduce canopy cover (Chambers *et al.* 1999). However, Alexander (1999) also documented a high proportion (relative to lower montane conifer forests) of canopy seed-eating, ground-foraging, and ground-nesting species in this type. The persistence of both canopy dependent and shrub-dependent species may be facilitated by landscape-scale patchiness created by fire.

Bird communities of the Pacific silver fir and mountain hemlock types of the Pacific Northwest are likely comparable to those in the same vegetation types in nearby areas of southern British Columbia, Canada, studied by Waterhouse *et al.* (2002). The most common species found at >900 m were Red Crossbill and Pine Siskin (*Carduelis pinus*); other relatively common species were Dark-eyed Junco, Winter Wren, Golden-crowned Kinglet, Chestnut-backed Chickadee, Townsend's Warbler (*Dendroica townsendi*), and Varied Thrush, which also were important species at lower elevations. Birds characteristic of the subalpine fir type in Washington include Pine Siskin, Mountain Chickadee (*Poecile gambeli*), Yellow-rumped Warbler (*Dendroica coronata*), and Clark's Nutcracker (*Nucifraga columbiana*; Manuwal *et al.* 1987).

Bird response to fire in the high-severity regime of the mountain hemlock, Pacific silver, and subalpine fir types probably parallels high-severity regimes of the western hemlock type. After fire, forest reestablishment is slower in these types than the western hemlock type due to short growing seasons, favoring ground- and brush-dwelling species over canopy feeder for up to a century or more. In subalpine environments, species that use edges and forest openings, such as Olive-sided Flycatcher, may benefit from fires (Altman and Sallabanks 2000, but see Meehan and George 2003).

ALTERATIONS TO FIRE REGIMES

Southwest Oregon and northwest California have a long history of anthropogenic influence on fire regimes (Frost and Sweeney 2000). This influence began with burning by American Indians (Boyd 1999), continued with fires set by Euro-American settlers, and then shifted to policies of fire suppression during the 20th century. However, establishing how and to what extent these activities have changed the natural fire regime is difficult. Such alterations in fire regimes can influence the local and landscape patterns of vegetation structure and community composition that determine habitat availability and quality for many forest birds.

The use of fire by American Indians in low- to moderate-severity fire regimes of the Pacific Northwest is accepted (Frost and Sweeney 2000), but the extent of these fires is not well known. Most burning by Indians probably occurred in oak woodland and pine forests (LaLande and Pullen 1999), possibly leading to a pattern where ignition of fires by American Indians was greatest at low elevations and decreased at higher elevations; there is little evidence that coniferous forests at higher elevations were significantly affected by aboriginal burning (Frost and Sweeney 2000).

Although burning by American Indians declined after the initiation of European settlement in the early 1800s (Frost and Sweeney 2000), a number of anecdotal accounts suggest that Euro-American settlement led to large and severe fires throughout many areas of the West (Biswell 1989). Although more frequent fires may have been the case in some areas, quantitative evidence has not shown this effect to be ubiquitous. In Douglas-fir dominated forests of the California Klamath Mountains, Taylor and Skinner (1998) found no difference between pre-settlement (1626–1849) and post-settlement (1850–1904) fire-return intervals. Similarly, at Kinney Creek in the eastern Oregon Klamath Mountains, Agee (1991b) documented a pre-settlement (1760–1860) fire frequency of 16 yr that was not substantially different from a nearby post-settlement (1850–1920) estimate of 12 yr.

As in most of the West, fire suppression became the policy towards forest fires in the Pacific Northwest in the early 1900s (Atzet and Wheeler 1982, Biswell 1989, Agee 1993). However, most evidence suggests that this policy did not become highly effective until after World War II, when fire fighting became more mechanized (Pyne 1982). Additionally, increases in cattle and sheep grazing in the California and Oregon Klamath Mountains (Atzet and Wheeler 1982) may have facilitated effective

fire suppression if it reduced fuels that carried fires (Biswell 1989).

Given the inherent variability within and among the different forest types, drawing generalizations about the effects of fire suppression on low- to moderate-severity fire regimes is difficult. At Oregon Caves National Monument in the Oregon Klamath Mountains, Agee (1991b) found that the fire-free period between 1920 and 1989 was the longest in over 300 yr. Perhaps more convincingly, in mixed conifer forests of the California Klamath Mountains, fire-return intervals increased from 12.5 yr during 1850–1904 (during the pre-suppression period) to 21.8 yr from 1905–1992 (during suppression) (Taylor and Skinner 1998), and fire rotation (the time required for the entire study area to burn) increased 10 fold from 20–238 yr (Taylor and Skinner 2003). Although increases in fire severity are often ascribed to policies of suppression (Biswell 1989), the evidence to support this hypothesis is limited and requires more information. Because fire suppression has reduced fire frequency, fuel levels are likely to be greater than they were historically. In areas where fire-return intervals were short, such build-ups may result in fires that are more severe and larger than they would have been historically (Agee 1993). However, in an analysis of fires in the Klamath Mountains between 1909 and 1997, Frost and Sweeney (2000) determined that although high severity fires were more common than they were prior to 1950, there was no conclusive evidence that this trend was outside of the historical range. More data are needed before we understand how fire suppression has affected fire severity in low- and moderate-severity fire regimes.

Effects of 20th century fire suppression may be responsible for changes in forest structure and landscape composition at several spatial scales in the Pacific Northwest (Kauffmann 1990). Generally, such effects can be more pronounced in forest types where historical fire-return intervals were shorter, low- and moderate-severity fire regimes, than where they were longer (Agee 1993). Fire suppression can alter the habitat potential for species associated with the composition and structure maintained by recurring fires by, for example, creating dense canopy layers that can substantially reduce herbaceous ground cover (e.g., Thilenius 1968, Hall 1977) and altering important roost characteristics used by birds of prey (e.g., Dellasala et al. 1998). At larger spatial scales, the effect of fire suppression on landscape composition may be more pronounced (Agee 1993). It has been hypothesized that in the mixed-conifer forests types with low- and (to a lesser extent) moderate-severity

fire regimes in the Pacific Northwest, fire suppression has decreased stand heterogeneity and promoted a forest landscape that is more even-aged than was historically present (Agee 1993). In the Klamath Mountains and California Cascade Range, this hypothesis is supported by a comparison of forest openings measured in 1945 and again in 1985 (Skinner 1995). He found that during this time period, the median distance from random points to the nearest opening doubled, suggesting that the spatial diversity of forest structure has declined since 1945. Such changes are likely to reduce variation in fire severity, an important source of structural diversity among stands in forested landscapes (Taylor and Skinner 2003), and may decrease biological richness.

In forest types that have experienced fire-return intervals >40 yr, structural and biological changes may be relatively minor because the period of fire suppression has been shorter than the typical fire-free intervals (Chappell and Agee 1996), though this is difficult to validate. Because large areas of Sitka spruce, western hemlock, subalpine fir, and mountain hemlock forest types have fire-return intervals that greatly exceed fire-free intervals fire suppression effects to date are probably minimal. In the Shasta red fir type, Chappell and Agee (1996) found little evidence that fuel loads or vegetation structure of stands were outside of the range of natural variation at the stand scale. They suggested, however, that at larger spatial scales fire suppression has allowed more stands to develop late-successional characteristics, and thus creating more structural homogeneity across the landscape.

EFFECTS OF FIRE SUPPRESSION ON BIRDS

No studies have directly addressed effects of fire suppression on bird abundance in the Pacific Northwest. However, we do know that many bird species vary in abundance across successional vegetation gradients (Marcot 1984, Raphael *et al.* 1988, Huff and Raley 1991, Ralph *et al.* 1991). Thus, if fire suppression has changed forest composition, regional patterns of bird abundance may have also changed (e.g., Raphael *et al.* 1988). If subtle demographic mechanisms are important in the birds and operate even in the absence of successional changes, then the effects of fire suppression may be more complex and difficult to quantify.

Fire suppression may be one factor contributing to changes in bird community composition in the interior valley type. In the absence of fire, coniferous trees and non-native shrubs encroach upon these habitats (Tveten and Fonda 1999). Such changes

have been implicated in changes in bird community composition between the late 1960s and early 1990s in oak woodlands (Hagar and Stern 2001). Fire has also been suppressed in sclerophyllous shrublands enabling trees to establish (Altman *et al.* 2000). Although the addition of trees to sclerophyllous shrublands could increase bird species diversity, converting the shrublands to forest is likely detrimental to shrubland bird populations.

Identifying general mechanisms by which fire suppression may influence changes in bird abundance may be difficult in coniferous forests. Information from nearby regions suggests that changes in forest structure caused by fire have important consequences for bird abundance. Studies in the Sierra Nevada and northeastern Washington reported canopy-dwelling and foraging species, including Golden-crowned Kinglet, Red-breasted Nuthatch, and Brown Creeper are consistently less abundant in burned areas (Bock and Lynch 1970, Kreisel and Stein 1999). These results are supported by similar responses of these species to commercial thinning (Chambers *et al.* 1999) or stand age (Marcot 1984, Raphael *et al.* 1988) that were studied in our region. To the extent that fire suppression has allowed denser forest canopies to develop over broad areas within the low-severity fire regime, species that use dense canopy characteristics should increase, while species strongly associated with fire-maintained vegetation composition and structure (e.g., canopy openings or diverse herb and shrub layers) should decrease. Olive-sided Flycatchers, for example, which typically increase in response to forest openings and open understory conditions created by fire (Hutto 1995; Altman, unpublished data) or commercial thinning (Chambers *et al.* 1999), declined broadly throughout the western North America from 1966 to 1996 (Altman, unpubl. data). These declines may be associated with habitat loss from fire suppression, but such an effect is difficult to verify over broad areas. Furthermore, changes in abundance may not be the best measure of habitat quality. In a study of Olive-sided Flycatchers in northern California, Meehan and George (2003) showed that although burned sites were more likely to be occupied than unburned sites, nest success was greater on the unburned sites.

ROLE OF PRESCRIBED FIRE AND FIRE MANAGEMENT

PRESCRIBED BURNING FOR HABITAT RESTORATION AND MAINTENANCE

Prescribed fire is increasingly recognized as a tool for restoring and maintaining forest health and

reducing fuels (Biswell 1989), yet the risks of escape and smoke production have limited its application. Estimates of how much area should be treated with fuel reduction activities were made during revisions to the Northwest Forest Plan (USDA and USDI 2000). The estimated area burned historically in this region ($\sim 190,000 \text{ ha yr}^{-1}$) was used as a starting point, and this was reduced by the average area currently burned by wildland fire and what is feasible to treat considering budget constraints to arrive at a goal of treating 76,000 ha annually.

In areas that burn frequently, fire may play an important role in maintaining species composition of some unique vegetation communities. Fire has been reintroduced to grasslands within the Pacific Northwest, for example, to maintain endangered plant species (Pendergrass et al. 1999) and to restore prairie-like conditions (Clark and Wilson 2001). Such applications of fire are considered experimental because so little is known about the natural fire regime and the desired results have not necessarily been achieved. Using prescribed fire to mimic natural fire regimes in these habitats has the potential to restore and maintain vegetation communities at watershed scales (e.g., Taylor and Skinner 2003) and associated bird communities.

BIRD CONSERVATION STRATEGIES USING PRESCRIBED FIRE

Bird conservation plans developed by Oregon-Washington Partners in Flight (2000) recommend prescribed burning in oak woodlands and savannas of the interior valley type in western Oregon and Washington. Burning in oak woodlands is being used as a conservation strategy for birds by reducing encroachment of Douglas-fir, stimulating oak seedling recruitment, and creating multi-aged stands. Prescribed fire is also being used to restore oak woodlands in other regions of the United States (e.g., Abrams 1992). Although prescribed fire may restore forest plant communities, the effects on bird communities are not well documented. Potential declines may occur as a result of changes in vegetation structure or in response to fires that take place during the breeding season (e.g., Artman et al. 2001). Although the timing of prescribed fire relative to the breeding season should be considered, these relatively short-term disturbances should not outweigh the long-term conservation benefits of restoring oak woodlands (Oregon-Washington Partners in Flight 2000).

Although the effects of prescribed burning on bird abundance in the Pacific Northwest have not

been documented, observations from the Sierra Nevada (e.g., Kilgore 1971) have been incorporated into the Oregon-Washington PIF conservation strategy for coniferous forests on the east slope of the Cascade Range (Oregon-Washington Partners in Flight 2000). This strategy recommends use of prescribed fire to reduce fuel loads and accelerate development of late-seral conditions. Such conditions are hypothesized to enhance habitat for Olive-sided Flycatchers as well as Cassin's Finch (*Carpodacus cassinii*), Western Wood-Pewee (*Contopus sordidulus*), Mountain Bluebird (*Sialia currucoides*), Northern Flicker (*Colaptes auratus*), American Kestrel (*Falco sparverius*), and American Robin (Oregon-Washington Partners in Flight 2000). Similarly, the California Partners in Flight conservation strategy for coniferous forests has identified the restoration of fire cycles as potential conservation measures for Black-backed Woodpeckers (*Picoides arcticus*), Fox Sparrows, and Olive-sided Flycatcher (California Partners in Flight 2002a).

Awareness is growing that prescribed fire and other forest management activities need to be considered within the context of natural wildfire on the landscape. The ability of birds to recruit to a site after fire depends on management activities that occur before and after fires. In particular, snag-nesting species, which typically increase following fire, are affected by postfire snag availability (e.g., Haggard and Gaines 2001). Snag-nesting bird communities were compared among three salvage-logged treatments (high, moderate, and low amounts of snags), 4–5 yr after a high-severity fire in the Douglas-fir type of the east-central Washington Cascades (Haggard and Gaines 2001). Distribution patterns of snags (e.g., clustering) and snag size affected cavity-nesting species response. Intermediate densities ($15\text{--}35 \text{ snags ha}^{-1}$) of snags $>25 \text{ cm}$ diameter at breast height (dbh) were associated with the highest abundance, species richness, and nesting densities of cavity nesters.

Another example of the interaction between fire and forest management is the potential effect of fire and fire management on habitat for endangered species. This is illustrated by the habitat requirements of the Northern Spotted Owl in areas of moderate-severity fire regimes throughout the Pacific Northwest. Historically, much of the habitat in this fire regime was spatially and temporally dynamic because of fires. More recently, decades of fire suppression and selective harvest of large, fire-resistant trees have created multi-canopied forests with thick understories of fire-intolerant species, which

possibly has accelerated a shift towards larger and more severe fires (Agee and Edmonds 1992). As a result, the amount of suitable Northern Spotted Owl habitat (USDI 1992, Agee 1993) may have increased in some forest types, particularly the grand fir type. Because these areas provide high quality owl habitat, they have been protected as late successional reserves designed to sustain owl populations. Consequently, high-severity fires are expected to burn a greater portion of the landscape than they did historically, increasing the probability that Northern Spotted Owl habitat will be altered for longer periods and at larger spatial scales. Some level of low-severity fire that maintains high canopy cover may have modest, or even beneficial effects for Northern Spotted Owls (Bond *et al.* 2002). The owl is more likely to be present and persist in areas of mixed successional stages and forest structures in northern California where fire is likely a major contributing factor to the mosaic of conditions that are favorable (Thome *et al.* 1999, Zabel *et al.* 2003). Severely-burned areas, however, are avoided and may reduce owl productivity (Bevis *et al.* 1997, Gaines *et al.* 1997, *cf.* Bond *et al.* 2002). If the extent of severely burn area increases substantially, the amount of Northern Spotted Owl habitat in reserves and managed areas may not be sufficient to sustain owl populations.

MANAGEMENT QUESTIONS AND FUTURE RESEARCH CONSIDERATIONS

Managing fire and fuels to meet the ecological, economic, and public safety concerns of society is challenging. From an ecological perspective, the ability to apply effective management is mostly limited by lack of relevant information to make informed decisions. In the Pacific Northwest, where fire has undoubtedly had a profound effect on bird communities, few studies have addressed how fire affects birds (Table 2), and basic information about bird species composition and relative abundance patterns was lacking for most vegetation types in this region. To effectively manage fire and fuels in this region while considering short- and long-term bird conservation will require substantial investments to accurately portray potentially affected bird communities across vegetation types, monitor response to varied management actions, and develop models to predict related bird response. Monitoring within the context of adaptive management (Holling 1978, Walters and Holling 1990) will be critical to assure that ecological goals of fire management are met (Tiedemann *et al.* 2000). In some cases, monitoring programs may be designed

in such a way to contribute information to important research questions, and vice-versa. With this potential in mind, we have outlined ten questions that, if answered, will, at least, provide critical information for the application of fire management toward effective bird conservation.

WHAT WERE CHARACTERISTICS OF NATURAL FIRE REGIMES IN THE PACIFIC NORTHWEST?

Clearly one of the most crucial steps to understanding the relationships between fire and bird abundance is to understand their interactions at a large spatial scale. This requires understanding, not just of the central tendency of fire regimes, but also their variability (Gill and McCarthy 1998). Given the complex nature of fire regimes in the Pacific Northwest, more data on the frequency, severity, and spatial distribution of fires is needed, especially in regions with low- to moderate-severity fire regimes.

HOW DO BIRD POPULATIONS CHANGE IN RESPONSE TO FIRE?

Even relatively basic information on the response of birds to fire is still lacking. Few studies have distinguished between changes in behavior (e.g., habitat use or nest-site selection) versus larger-scale changes in population density. Studies that measure behavioral responses to fire (e.g., foraging activity [Kreisel and Stein 1999]), and nesting density and demographics (e.g., Saab and Dudley 1998, Saab *et al.* 2002) could provide these data. Using geographic information system (GIS) data to link these patterns with landscape level patterns may provide useful insights (Dettmers and Bart 1999, Saab *et al.* 2002). If fire affects important demographic patterns (e.g., Saab and Vierling 2001), then we should consider how source/sink dynamics might be influenced at large spatial scales.

WHEN BIRD POPULATIONS CHANGE IN RESPONSE TO FIRE, WHAT ARE THE DRIVING FACTORS?

With a better understanding of how bird abundance changes in response to fire, it will be important to determine causal explanations. A number of mechanisms have been hypothesized to influence postfire changes in bird communities, including food availability (Apfelbaum and Haney 1981), nest-site availability (Hutto 1995), predator abundance (Altman and Sallabanks 2000, Saab and Vierling 2001), and vegetation structure (Bock and Lynch 1970). An evaluation of the relative importance of these factors

TABLE 2. SUMMARY OF BIRD RESPONSES TO FIRE BY SPECIES.

Species	State	Year after fire	Size (ha) and No. of fires ^a	No. of replicate sites	Response ^b	Reference ^c	Comments ^d
Blue Grouse (<i>Dendragapus obscurus</i>)	WA	4	4000 (2)	13 burned, 9 unburned	0	1	1
Spotted Owl (<i>Strix occidentalis</i>)	CA, AZ, NM	1	>540 (1)	11 territories	0	2	2, Postfire estimates not outside range of unburned estimates.
Rufous Hummingbird (<i>Selasphorus rufus</i>)	WA	1-515	no data (7)	7 plots	m	3	3, Most abundant 19 yr postfire.
Downy Woodpecker (<i>Picoides pubescens</i>)	WA	4	4000 (2)	13 burned, 9 unburned	0	1	1
Hairy Woodpecker (<i>Picoides villosus</i>)	WA	4	4000 (2)	13 burned, 9 unburned	+	1	1
American Three-toed Woodpecker (<i>Picoides dorsalis</i>)	WA	4	4000 (2)	13 burned, 9 unburned	+	1	1
Black-backed Woodpecker (<i>Picoides arcticus</i>)	WA	4	4000 (2)	13 burned, 9 unburned	+	1	1
Pileated Woodpecker (<i>Dryocopus pileatus</i>)	WA	4	4000 (2)	13 burned, 9 unburned stations	0	1	1
Pacific-slope Flycatcher (<i>Empidonax difficilis</i>)	WA	1-515	np (7)	7 plots	-	3	3
Gray Jay (<i>Perisoreus canadensis</i>)	WA	4	4000 (2)	13 burned, 9 unburned	0	1	1
Steller's Jay (<i>Cyanocitta stelleri</i>)	WA	4	4000 (2)	13 burned, 9 unburned	0	1	1
Clark's Nutcracker (<i>Nucifraga columbiana</i>)	WA	4	4000 (2)	13 burned, 9 unburned	0	1	1
Black-billed Magpie (<i>Pica hudsonia</i>)	WA	4	4000 (2)	13 burned, 9 unburned	0	1	1
Common Raven (<i>Corvus corax</i>)	WA	4	4000 (2)	13 burned, 9 unburned	0	1	1
Black-capped Chickadee (<i>Poecile atricapilla</i>)	WA	4	4000 (2)	13 burned, 9 unburned	0	1	1
Mountain Chickadee (<i>Poecile gambeli</i>)	WA	4	4000 (2)	13 burned, 9 unburned	0	1	1
Red-breasted Nuthatch (<i>Sitta canadensis</i>)	WA	1-515	np (7)	7 plots	-	3	3
Chestnut-backed Chickadee (<i>Poecile rufescens</i>)	WA	4	4000 (2)	13 burned, 9 unburned	-	1	1
White-breasted Nuthatch (<i>Sitta carolinensis</i>)	WA	4	4000 (2)	13 burned, 9 unburned	0	1	1
Brown Creeper (<i>Certhia americana</i>)	WA	4	4000 (2)	13 burned, 9 unburned	0	1	1
Brown Creeper	WA	1-515	np (7)	7 plots	m	3	3
Winter Wren (<i>Troglodytes troglodytes</i>)	WA	4	4000 (2)	13 burned, 9 unburned	0	1	1
Winter Wren	WA	1-515	np (7)	7 plots	+	3	3
Golden-crowned Kinglet (<i>Regulus satrapa</i>)	WA	4	4000 (2)	13 burned, 9 unburned	-	1	1
Golden-crowned Kinglet	WA	1-515	np (7)	7 plots	-	3	3
American Robin (<i>Turdus migratorius</i>)	WA	1-515	np (7)	7 plots	m	3	Most abundant 3-19 yr postfire.
Varied Thrush (<i>Ixoreus naevius</i>)	WA	4	4000 (2)	13 burned, 9 unburned	0	1	1
Townsend's Warbler (<i>Dendroica townsendi</i>)	WA	1-515	np (7)	7 plots	m	3	3
Dark-eyed Junco (<i>Junco hyemalis</i>)	WA	1-515	np (7)	7 plots	+	3	3
Red Crossbill (<i>Loxia curvirostra</i>)	WA	4	4000 (2)	13 burned, 9 unburned	0	1	1

^a Only wildland fires are reported in this table.

^b + = increase; - = decrease; 0 = no effect or study inconclusive; m = mixed response.

^c References: 1 = Kreisel and Stein 1999, compared 13 stations in two 80-ha stand replacement burns with nine unburned stations; 2 = Bond et al. 2002, compared survival, site fidelity, and reproductive success for owls 1 yr postfire to unburned estimates; 3 = Huff et al. 1985, chronosequence comparing seven sites ranging from 1-515 yr postfire, positive fire response if birds more abundant in years immediately after fire.

for different nesting and foraging guilds is needed. An effective evaluation of these hypotheses could include experimental manipulations or large sample sizes in many treatments. Understanding the mechanisms responsible for these changes will provide a better understanding of the unique ecological effects of fire-mediated habitat change.

HOW DO FIRE REGIMES INFLUENCE THE BIRD COMMUNITY STRUCTURE AT LOCAL AND LANDSCAPE LEVELS?

With data on the response of bird populations to fire, researchers and managers will be in a better position to understand how fire regimes structure bird communities. To date, most studies of bird communities and fires have been performed at the scale of single forest stands (Bock and Lynch 1970, Kreisel and Stein 2000, but see Saab et al. 2002). In order to understand the relationship of fires and birds, it is important not only to measure the effect of fire at burned and unburned points, but to quantify the effect of fire, and different spatial patterns of fire, at larger spatial scales, such as watersheds or regions. Fire likely influences abundance and distribution of birds throughout entire regions, not simply within the immediate area of disturbance.

HOW DO CHANGES TO FIRE-RETURN INTERVALS AFFECT BIRD POPULATIONS?

It is clear that fire suppression has lengthened fire-return intervals in the low- and to some degree moderate-severity fire regimes, and may have altered patterns of fire severity. The ecological effects of these changes are not well understood (Chappell and Agee 1996, Frost and Sweeney 2000), but changes in composition and spatial distribution of large-scale habitat characteristics are probably important (Skinner 1995). Effects of reducing fire frequencies on bird abundance and demographics should be evaluated if a policy of suppression continues, and at least outside of wilderness areas it probably will. Such evaluations should be conducted at multiple spatial scales by expanding analyses beyond comparisons of burned and unburned points to consider how spatial distribution of landscape characteristics (e.g., edge/patch ratios or habitat composition around points). Inferences drawn from microhabitat and landscape-scale characteristics may be substantially different and vary depending on life-history characteristics (e.g., migration strategy) of the birds in question (Mitchell et al. 2001).

HAVE THE EFFECTS OF WILDFIRES ON BIRD POPULATIONS CHANGED?

Fire suppression may have created fires that are less frequent, larger, and more severe than those occurring previously. The potential shift to large and more severe fires may affect bird abundance and community composition. Little is known about potential changes in bird populations as a result of changing fire regimes. Comparing long-term data on bird abundance (e.g., Breeding Bird Survey data) with demographic models that can evaluate the effect of landscape composition on bird populations may be one way to evaluate the effect of changes created by fire suppression. Such an approach may be especially useful in evaluating the degree to which long-term declines of species that respond positively to fire may be a result of fire suppression policies.

HOW DOES PRESCRIBED FIRE CHANGE CONDITIONS FOR BIRDS?

A return to more natural fire regimes is often advocated, but given the constraints imposed by concerns of public safety, economics, and air quality, it is more likely that this will be achieved through the application of prescribed fire in combination with manual and mechanical treatments (e.g., thinning of trees and removal of invasive shrubs). The application of these management tools may occur during seasons when fires did not normally burn. In the Pacific Northwest, fires burned naturally from June through September, with most occurring in late summer or early fall (Taylor and Skinner 1998, Brown et al. 1999). To maximize fire control, however, late winter or early spring is usually a better time to apply prescribed burns (Biswell 1989) to prevent escape. Effects of such burning schedules on bird abundance may be immediate, especially for ground- and shrub-nesting species, when they conflict with the breeding season. Such effects may be relatively short-lived and must be considered within the context of long-term benefits of prescribed fire (California Partners in Flight 2002). These changes are poorly understood in the Pacific Northwest, but are presumed to include vegetation changes that lead to longer-lasting changes in bird abundance (Kilgore 1971) or nest success (Jones et al. 2002). Prescribed fire can be applied in many different ways and have many different effects on vegetation and fuel and are most likely to be applied to the low- and moderate-severity fire regime vegetation types. Use and effects of prescribed fire in these types, which include Douglas-fir, grand fir, mixed evergreen hardwood, white fir, Shasta red fir, and

coast redwood, is complex and generalizations are not easily drawn from one type that can be applied easily to birds in another type.

HOW DO FIRE SURROGATES (FUELS TREATMENTS) AFFECT BIRD POPULATIONS?

An alternative to prescribed fire is the use of fire surrogates (usually mechanical methods) to reduce fuels and mimic other structural and possibly ecological effects of fire. Such management activities vary widely in intensity, from collection and removal of fuels by hand to large-scale mechanical processes that have high amounts of incidental disturbance. Evidence suggests that these activities do not create the same structural or ecological conditions as a natural or prescribed fire (Imbeau et al. 1999, and see Hannon and Drapeau, this volume). For example, when reproductive success was examined for Olive-sided Flycatchers associated with recent burns and logged areas in the western Oregon Cascade Mountains (Altman and Sallabanks 2000), nest success at burned sites was nearly twice that in logged area, suggesting that openings created by fire are better habitat for the flycatcher than logged areas. In contrast, a similar study by Meehan and George (2003) showed that the probability of nest-loss was greater in burned than unburned areas. Clearly even the effect of fire on bird reproduction, let alone the ability of mechanical fuels reduction to mimic these effects, is poorly understood. Mechanical methods reduce fuels and are certainly likely to reduce fire risk. After fire, however, standing dead trees remain. Trees removed by mechanical methods create habitat that is structurally and ecologically different from postfire conditions. These activities should be monitored with respect to their effect on bird populations in order to evaluate their ecological effects.

WHAT ARE EFFECTS OF POSTFIRE SALVAGE?

Postfire salvage logging is a controversial practice that removes some amount of dead and at least

apparently damaged live trees from burned areas. After fire, snags provide important nest sites and foraging opportunities for many cavity-nesting and bark-foraging birds (Hutto 1995, Kreisel and Stein 1999). Salvage logging has negative consequences for some bird species (Hutto 1995, Wales 2001). Studies from the Washington Cascades (Haggard and Gaines 2001) and ponderosa pine forests of southwestern Idaho (Saab and Dudley 1998) have both suggested that snag density and distribution are important factors influencing the abundance of cavity-nesting birds. As in many other situations, the effects of such treatments may interact with burn severity. Removing structure from severely burned areas may have different effects than the same level of extraction from less severely burned areas (Kotliar et al. 2002). Investigations of the effects of salvage should consider these types of interactions.

IF CLIMATE CHANGE ALTERS FIRE REGIMES, WHAT IS THE EFFECT ON BIRD COMMUNITIES?

The possibility that climate change may alter local fire regimes should be considered (Johnson and Larsen 1991). Such changes may affect bird abundance, community composition, distribution, and diversity. Predicting the effects of climate change on fire regimes and bird communities may provide us with the opportunity to test hypotheses about the relative importance of deterministic and stochastic processes in community assembly.

ACKNOWLEDGMENTS

We thank Rich Young for assistance on developing the map figure. Portions of this paper benefited from our discussions with Danny Lee and Steve Norman. Bob Altman, Vicki Saab, Hugh Powell, and two anonymous reviewers provided helpful editorial comments.

LITERATURE CITED

- ABRAHAMSON, W. G., AND D. C. HARTNETT. 1990. Pine flatwoods and dry prairies. Pp. 103–149 in R. L. Myers, and J. J. Ewel (editors). *Ecosystems of Florida*. University of Central Florida Press, Orlando, FL.
- ABRAMS, M. D. 1985. Fire history of oak gallery forests in a northeast Kansas tallgrass prairie. *American Midland Naturalist* 114:188–191.
- ABRAMS, M. D. 1992. Fire and the development of oak forests. *BioScience* 42:346–353.
- ABRAMS, M. D. 1998. The red maple paradox. *BioScience* 48:355–364.
- AGEE, J. K. 1981. Fire effects on Pacific Northwest forests: flora, fuels, and fauna. Pp. 54–66 in *Proceedings, Northwest Forest Fire Council proceedings*. Northwest Forest Fire Council, Portland, OR.
- AGEE, J. K. 1990. The historical role of fire in Pacific Northwest forests. Pp. 25–38 in J. D. Walstad, S. R. Radosevich, and D. V. Sandberg (editors). *Natural and prescribed fire in Pacific Northwest forests*. Oregon State University Press, Corvallis, OR.
- AGEE, J. K. 1991a. Fire history of Douglas-fir forests in the Pacific Northwest. Pp. 25–34 in L. Ruggiero, K. Aubry, A. B. Carey, and M. H. Huff (technical coordinators). *Wildlife and vegetation of unmanaged Douglas-fir forests*. USDA Forest Service General Technical Report PNW-GTR-285. USDA Forest Service, Pacific Northwest Research Station, Portland, OR.
- AGEE, J. K. 1991b. Fire history along an elevational gradient in the Siskiyou Mountains, Oregon. *Northwest Science* 65:188–199.
- AGEE, J. K. 1993. *Fire ecology of Pacific Northwest forests*. Island Press, Washington, DC.
- AGEE, J. K. 1998. The landscape ecology of western forest fire regimes. *Northwest Science* 72:24–34.
- AGEE, J. K. 2002. The fallacy of passive management: managing for firesafe forest reserves. *Conservation Biology In Practice* 3:19–25.
- AGEE, J. K., AND R. E. EDMONDS. 1992. Appendix F. Forest protection guidelines for the Northern Spotted Owl. Pp. 419–480 in J. Bart, R. G. Anthony, M. Berg, J. H. Beuter, W. Elmore, J. Fay, R. J. Gutiérrez, H. T. Heintz, Jr., R. S. Holthausen, K. Starkey, and J. C. Tappeiner (editors). *Recovery plan for the Northern Spotted Owl: final draft*. U.S. Department of the Interior, Washington DC.
- AGEE, J. K., AND R. FLEWELLING. 1983. A fire cycle model based on climate for the Olympic Mountains, Washington. *Fire and Forest Meteorology Conference* 7:32–37.
- AGEE, J. K., AND M. H. HUFF. 1987. Fuel succession in a western hemlock/Douglas-fir forest. *Canadian Journal of Forest Research* 17:697–704.
- AGEE, J. K., AND J. KERTIS. 1987. Forest types of the North Cascades National Park Service Complex. *Canadian Journal of Botany* 65:1520–1530.
- AGEE, J. K., AND F. KRUSEMARK. 2001. Forest fire regime of the Bull Run Watershed, Oregon. *Northwest Science* 75:292–306.
- AGEE, J. K., AND L. SMITH. 1984. Subalpine tree establishment after fire in the Olympic Mountains, Washington. *Ecology* 65:810–819.
- AGEE, J. K., M. FINNEY, AND R. DEGOUVENAIN. 1990. Fire history of Desolation Peak, Washington. *Canadian Journal of Forest Research* 20:350–356.
- AGEE, J. K., B. BAHRO, M. A. FINNEY, P. N. OMI, D. B. SAPSIS, C. N. SKINNER, J. W. VAN WAGTENDONK, AND C. P. WEATHERSPOON. 2000. The use of shaded fuelbreaks in landscape fire management. *Forest Ecology and Management* 127:55–66.
- ALEXANDER, J. D. 1999. Bird-habitat relationships in the Klamath/Siskiyou Mountains. M.S. thesis, Southern Oregon University, Ashland, OR.
- ALLEN, C. D., J. L. BETANCOURT, AND T. W. SWETNAM. 1998. Landscape changes in the southwestern United States: Techniques, long-term datasets, and trends. Pp. 71–84. in T. D. Sisk (editor). *Perspectives on the land use history of North America: a context for understanding our changing environment*. USDI Geological Survey Biological Science Report USGS/BRD/BSR-1998-0003. USDI Geological Survey, Biological Resources Division, Washington, DC.
- ALLEN, C. D., M. SAVAGE, D. A. FALK, K. F. SUCKLING, T. W. SWETNAM, T. SCHULKE, P. B. STACEY, P. MORGAN, M. HOFFMAN, AND J. KLINGEL. 2002. Ecological restoration of southwestern ponderosa pine ecosystems: A broad perspective. *Ecological Applications* 12:1418–1433.
- ALLEN, E. B. (EDITOR). 1988. *The reconstruction of disturbed arid lands: an ecological approach*. Westview Press, Boulder, CO.
- ALLEN-DIAZ, B. H. AND J. W. BARTOLOME. 1992. Survival of *Quercus douglasii* (Fagaceae) seedlings under the influence of fire and grazing. *Madroño* 39:47–53.
- ALLEN-DIAZ, B., AND J. W. BARTOLOME. 1998. Sagebrush-grass vegetation dynamics: comparing classical and state-transition models. *Ecological Applications* 8: 795–804.
- ALTMAN, B., AND R. SALLABANKS. 2000. Olive-sided Flycatcher (*Conopus cooperi*). In A. Poole, and F. Gill (editors). *The Birds of North America*, No. 502. The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists' Union, Washington, DC.
- ALTMAN, B., M. HAYES, S. JANES, AND R. FORBES. 2000. Wildlife of westside grassland and chaparral habitats. Pp. 261–291 in D. H. Johnson, and T. A. O'Neil (managing directors). *Wildlife-habitat relationships in Oregon and Washington*. Oregon State University Press, Corvallis, OR.
- ANDERSON, K. 1993. Indian fire-based management in the sequoia-mixed conifer forests of the central and

- southern Sierra Nevada. Final Report, Cooperative Agreement 8027-2-002, Yosemite Research Center, Yosemite National Park, CA.
- ANDERSON, M. K., AND M. J. MORATTO. 1996. Native American land-use practices and ecological impacts. Pp. 187–206 in *Sierra Nevada Ecosystem Project: Final report to Congress, Volume II, Assessments and scientific basis for management options*. Wildland Resources Center Report No. 37. University of California, Centers for Water and Wildland Resources, Davis, CA.
- ANDERSON, R. C., AND M. L. BOWLES. 1999. Deep-soil savannas and barrens of the midwestern United States. Pp. 155–170 in R. C. Anderson, J. S. Fralish, and J. M. Baskin (editors). *Savannas, barrens, and rock outcrop plant communities of North America*. Cambridge University Press, Cambridge, UK.
- ANDERSON, R. C., J. S. FRALISH, AND J. M. BASKIN (EDITORS). 1999. *Savannas, barrens, and rock outcrop plant communities of North America*. Cambridge University Press, Cambridge, UK.
- ANDERSON, S. H. 1970. The avifaunal composition of Oregon white oak stands. *Condor* 72:417–423.
- ANGELSTAM, P. K. 1998. Maintaining and restoring biodiversity in European boreal forests by developing natural disturbance regimes. *Journal of Vegetation Science* 9:593–602.
- ANNAND, E. M., AND F. R. THOMPSON III. 1997. Forest bird response to regeneration practices in central hardwood forests. *Journal of Wildlife Management* 61:159–171.
- APFELBAUM, S., AND A. HANEY. 1981. Bird populations before and after wildfire in a Great Lakes pine forest. *Condor* 83:347–354.
- AQUILANI, S. M., D. C. LEBLANC, AND T. E. MORRELL. 2000. Effects of prescribed surface fires on ground- and shrub-nesting Neotropical migratory birds in a mature Indiana oak forest, USA. *Natural Areas Journal* 20: 317–24.
- ARCHER, S. 1994. Woody plant encroachment into southwestern grasslands and savannas: rates, patterns, and proximate causes. Pp. 13–68 in M. Vavra, W. A. Laycock, and R. D. Pieper (editors). *Ecological implications of livestock herbivory in the West*. Society for Range Management, Denver, CO.
- ARCHER, S., D. S. SCHIMMEL, AND E. A. HOLLAND. 1995. Mechanisms of shrubland expansion: land use, climate or CO₂? *Climatic Change* 29:91–99.
- ARTMAN, V. L., AND J. F. DOWNHOWER. 2003. Wood Thrush (*Hylocichla mustelina*) nesting ecology in relation to prescribed burning of mixed-oak forest in Ohio. *Auk* 120:874–882.
- ARTMAN, V. L., E. K. SUTHERLAND, AND J. F. DOWNHOWER. 2001. Prescribed burning to restore mixed-oak communities in southern Ohio: effects on breeding bird populations. *Conservation Biology* 15:1423–1434.
- ARNO, S. F. 2000. Fire in western forest ecosystems. Chapter 5 in J. K. Brown, and J. K. Smith (editors). *Wildland fire in ecosystems: effects of fire on flora*. USDA Forest Service General Technical Report RMRS-GTR-42-vol. 2. USDA Forest Service, Intermountain Research Station, Ogden, UT.
- ARNO, S. F., AND J. K. BROWN. 1991. Overcoming the paradox in managing wildland fire. *Western Wildlands* 17:40–46.
- ARNO, S. F., AND G. E. GRUELL. 1983. Fire history at the forest-grassland ecotone in southwestern Montana. *Journal of Range Management* 36:332–336.
- ARNO, S. F., AND T. D. PETERSEN. 1983. Variation in estimates of fire intervals: a closer look at fire history on the Bitterroot National Forest. USDA Forest Service Research Paper INT-301. USDA Forest Service, Intermountain Research Station, Ogden, UT.
- ARNO, S. F., J. H. SCOTT, AND M. G. HARTWELL. 1995. Age-class structure of old growth ponderosa pine/Douglas-fir stands and its relationship to fire history. USDA Forest Service Research Paper INT-RP-481. USDA Forest Service, Intermountain Research Station, Ogden, UT.
- ASKINS, R. A. 1993. Population trends in grassland, shrubland, and forest birds in eastern North America. *Current Ornithology* 11:1–34.
- ASKINS, R. A. 1998. Restoring forest disturbances to sustain populations of shrubland birds. *Restoration and Management Notes* 16:166–173.
- ASKINS, R. A. 1999. History of grassland birds in north-eastern North America. *Studies in Avian Biology* 19: 60–71.
- ASKINS, R. A. 2000. *Restoring North America's birds: lessons from landscape ecology*. Yale University Press, New Haven, CT.
- ATZET, T., AND D. L. WHEELER. 1982. Historical and ecological perspectives on fire activity in the Klamath geological province of the Rogue River and Siskiyou National Forests. USDA Forest Service Technical Report R-6-Range-102. USDA Forest Service, Pacific Northwest Region, Portland, OR.
- ATZET T., D. E. WHITE, L. A. MCCRIMMON, P. MARTINEZ, P. FONG, AND V. RANDALL. 1996. Field guide to the forested plant associations of southwestern Oregon. USDA Forest Service R6-ECOL-TP-17-96. USDA Forest Service, Pacific Northwest Region, Portland, OR.
- ATZET, T., D. L. WHEELER, B. SMITH, J. FRANKLIN, G. RIEGEL AND D. THORNBURGH. 1992. Vegetation. Pp. 92–113 in S. D. Hobbs, S. D. Tesch, P. Owston, R. Stewart, J. C. Tappeiner, and G. A. Wells (editors). *Reforestation practices in southwest Oregon and Northern California*. Forest Research Laboratory, Oregon State University, Corvallis, OR.
- AXELROD, D. I. 1985. The rise of the grassland biome, central North America. *Botanical Review* 51:163–201.
- BAHRE, C. J. 1985. Wildfire in southeastern Arizona between 1859 and 1890. *Desert Plants* 7:190–194.
- BAHRE, C. J. 1995. Human impacts on the grasslands of southeastern Arizona. Pp. 230–264 in M. P. McClaran, and T. R. Van Devender (editors). *The desert grassland*. University of Arizona Press, Tucson, AZ.
- BAHRE, C. J., AND M. L. SHELTON. 1993. Historic vegetation change, mesquite increases, and climate in southeastern Arizona. *Journal of Biogeography* 20:489–504.

- BAKER, W. L., AND D. EHLE. 2001. Uncertainty in surface-fire history: the case of ponderosa pine forests in the western United States. *Canadian Journal of Forest Research* 31:1205–1226.
- BAKER, W. L., AND D. J. SHINNEMAN. 2004. Fire and restoration of piñon-juniper woodlands in the western United States: a review. *Forest Ecology and Management* 189: 1–21.
- BALDA, R. P., AND N. MASTERS. 1980. Avian communities in the pinyon-juniper woodland. Pp. 146–167 in R. M. DeGraff (technical coordinator). Workshop proceedings, management of western forests and grasslands for nongame birds. USDA Forest Service General Technical Report INT-86. USDA Forest Service, Intermountain Research Station, Ogden, UT.
- BALLING, R. C., G. A. MEYER, AND S. G. WELLS. 1992. Relation of surface climate and burned area in Yellowstone National Park. *Agricultural Forest Meteorology* 60:285–293.
- BARBOUR, H., T. SIMMONS, P. SWAIN, AND H. WOOLSEY. 1999. Our irreplaceable heritage: protecting biodiversity in Massachusetts. Natural Heritage and Endangered Species Program, Massachusetts Division of Fish and Wildlife and The Nature Conservancy, Boston, MA.
- BARBOUR, M. G., AND W. D. BILLINGS. 2000. North American terrestrial vegetation (second edition). Cambridge University Press, Cambridge, UK.
- BARBOUR, M. G., AND J. MAJOR (EDITORS). 1977. Terrestrial vegetation of California. Wiley-Interscience, New York, NY.
- BARNES, T. A., AND D. H. VAN LEAR. 1998. Prescribed fire effects on advanced regeneration in mixed hardwood stands. *Southern Journal of Applied Forestry* 22: 138–142.
- BARRETT, S. W., AND S. F. ARNO. 1982. Indian fires as an ecological influence in the northern Rockies. *Journal of Forestry* 80:647–651.
- BART, J., G. ANTHONY, M. BERG, J. H. BEUTER, W. ELMORE, J. FAY, R. J. GUTIÉRREZ, H. T. HEINTZ, JR., R. S. HOLTHAUSEN, K. STARKEY, AND J. C. TAPPEINER. 1992. Recovery plan for the Northern Spotted Owl: final draft. U.S. Department of the Interior, Washington DC.
- BARTON, A. M. 1995. Fire adaptations in pines and oaks: tree population response to fire suppression in Arizona's Madiran forests. Pp. 159–163 in J. K. Brown, R. W. Mutch, C. W. Spoon, and R. H. Wakimoto (technical coordinators). Proceedings: symposium on fire in wilderness and park management. USDA Forest Service General Technical Report INT-GTR-320. USDA Forest Service, Intermountain Research Station, Ogden, UT.
- BEKKER, M., AND A. H. TAYLOR. 2001. Gradient analysis of fire regimes in montane forests of the Thousand Lakes Wilderness, southern Cascades, California, USA. *Plant Ecology* 155:15–28.
- BENTLEY, J. R., AND R. L. FENNER. 1958. Soil temperatures during burning related to postfire seedbeds on woodland range. *Journal of Forestry* 56:737–740.
- BERGERON, Y. 2000. Species and stand dynamics in the mixed-woods of Quebec's southern boreal forest. *Ecology* 81:1500–1516.
- BERGERON, Y., AND A. LEDUC. 1998. Relationships between change in fire frequency and mortality due to spruce budworm out-breaks in the southern Canadian boreal forest. *Journal of Vegetation Science* 9:492–500.
- BERGERON, Y., B. HARVEY, A. LEDUC, AND S. GAUTHIER. 1999. Basing forest management on natural disturbance: stand- and landscape-level considerations. *Forestry Chronicle* 75:49–54.
- BERGERON, Y., A. LEDUC, B. D. HARVEY, AND S. GAUTHIER. 2002. Natural fire regime: a guide for sustainable management of the Canadian boreal forest. *Silva Fennica* 36:81–95.
- BERGERON, Y., S. GAUTHIER, V. KAFKA, P. LEFORT, AND D. LESIEUR. 2001. Natural fire frequency for the eastern Canadian boreal forest: consequences for sustainable forestry. *Canadian Journal of Forest Research* 31: 384–391.
- BESSIE, W. C., AND E. A. JOHNSON. 1995. The relative importance of fuels and weather on fire behavior in subalpine forests in the southern Canadian Rockies. *Ecology* 26:747–762.
- BEST, L. B. 1972. First year effects of sagebrush control on two sparrows. *Journal of Wildlife Management* 36: 534–544.
- BETTINGER, K. A. 1996. Bird communities in 5 to 34-year-old managed Douglas-fir stands on the Willamette National Forest, Oregon Cascades. M.S. thesis, Oregon State University, Corvallis, OR.
- BEVIS, K. R., G. M. KING, AND E. E. HANSEN. 1997. Spotted Owls and 1994 fires on the Yakama Indian Reservation. Pp. 117–122 in J. M. Greenlee (editor). Proceedings, First conference on fire effects on rare and endangered species and habitats. International Association of Wildland Fire, Fairfield, WA.
- BIDWELL, T. (EDITOR). 2003. Ecology and management of the Greater Prairie-Chicken. Oklahoma Cooperative Extension Unit, Publication E-969. Oklahoma State University, Stillwater, OK.
- BILLINGS, W. D. 1994. Ecological impacts of cheatgrass and resultant fire on ecosystems in the Great Basin. Pp. 22–30 in S. B. Monsen, and S. G. Kitchen (compilers). Proceedings—ecology and management of annual grasslands. USDA Forest Service General Technical Report INT-GTR-313. USDA Forest Service, Intermountain Research Station, Ogden, UT.
- BISWELL, H. H. 1967. The use of fire in wildland management in California. Pp. 71–86 in S. V. Ciriacy-Wantrup, and J. J. Parsons (editors). Natural resources: quality and quantity. University of California Press, Berkeley, CA.
- BISWELL, H. 1989. Prescribed burning in California wildlands vegetation management. University of California Press, Berkeley, CA.
- BLAISDELL, J. P., AND R. C. HOLMGREN. 1984. Managing intermountain rangelands—salt desert shrub ranges. USDA Forest Service General Technical Report INT-

163. USDA Forest Service, Intermountain Forest and Range Experiment Station, Ogden, UT.
- BLAKE, J. G. 1982. Influence of fire and logging on non-breeding bird communities of ponderosa pine forests. *Journal of Wildlife Management* 46:404–415.
- BLAKE, J. G., AND B. SCHUETTE. 2000. Restoration of an oak forest in east-central Missouri: early effects of prescribed burning on woody vegetation. *Forest Ecology and Management* 139:109–126.
- BLOCK, W. M., AND M. L. MORRISON. 1990. Wildlife diversity of the central Sierra foothills. *California Agriculture* 44:19–22.
- BOCK, C. E., AND W. M. BLOCK. In press. Response of birds to fire in the American Southwest. Proceedings: Bird conservation implementation and integration in the Americas. USDA Forest Service General Technical Report. USDA Forest Service, Pacific Southwest Research Station, Berkeley, CA.
- BOCK, C. E., AND J. H. BOCK. 1978. Response of birds, small mammals, and vegetation to burning sacaton grasslands in southeastern Arizona. *Journal of Range Management* 31:296–300.
- BOCK, C. E., AND J. H. BOCK. 1992b. Response of birds to wildfire in native versus exotic Arizona grassland. *Southwestern Naturalist* 37:73–81.
- BOCK, C. E., AND J. H. BOCK. 1984. Importance of sycamores to riparian birds in southeastern Arizona. *Journal of Field Ornithology* 55:97–103.
- BOCK, C. E. AND J. H. BOCK. 1987. Avian habitat occupancy following fire in a Montana shrubsteppe. *Prairie Naturalist* 19:153–158.
- BOCK, C. E., AND Z. F. JONES. 2004. Avian habitat evaluation: should counting birds count? *Frontiers in Ecology and the Environment* 2:403–410.
- BOCK, C. E., AND J. F. LYNCH. 1970. Breeding bird populations of burned and unburned conifer forest in the Sierra Nevada. *Condor* 72:182–189.
- BOCK, C. E., AND B. WEBB. 1984. Birds as grazing indicator species in southeastern Arizona. *Journal of Wildlife Management* 48:1045–1049.
- BOCK, C. E., J. H. BOCK, AND H. M. SMITH. 1993. Proposal for a system of federal livestock exclosures on public rangelands in the western United States. *Conservation Biology* 7:731–733.
- BOCK, J. H., AND C. E. BOCK. 1992a. Vegetation responses to wildfire in native vs. exotic Arizona grassland. *Journal of Vegetation Science* 3:439–446.
- BOCK, J. H., AND C. E. BOCK. 1998. Tallgrass prairie: remnants and relicts. *Great Plains Research* 8:213–230.
- BOCK, J. H., C. E. BOCK, AND J. R. MCKNIGHT. 1976. A study of the effects of grassland fires at the Research Ranch in southeastern Arizona. *Journal of the Arizona Academy of Science* 11:49–57.
- BOERNER, R. E. J., E. K. SUTHERLAND, S. J. MORRIS, AND T. F. HUTCHINSON. 2000. Spatial variation in the effect of prescribed fire on N dynamics in a forested landscape. *Landscape Ecology* 15:425–439.
- BOLANDER, D. H. 1981. Chaparral in Arizona. Pp. 60–63 in C. E. Conrad, and W. C. Oechel (technical coordinators). Dynamics and management of Mediterranean-type ecosystems. USDA Forest Service General Technical Report PSW-58. USDA Forest Service, Pacific Southwest Forest and Range Experiment Station, Berkeley, CA.
- BOND, M. L., R. J. GUTIÉRREZ, A. B. FRANKLIN, W. S. LAHAYE, C. A. MAY, AND M. E. SEAMANS. 2002. Short-term effects of wildfires on Spotted Owl survival, site fidelity, mate fidelity, and reproductive success. *Wildlife Society Bulletin* 30:1022–1028.
- BONNICKSON, T. M. 2000. America's ancient forests. From the ice age to the age of discovery. John Wiley and Sons, New York, NY.
- BOREN, J. C., D. M. ENGLE, M. W. PALMER, R. M. MASTERS, AND T. CRINER. 1999. Land use change effects on breeding bird community composition. *Journal of Range Manage.* 52:420–430.
- BORK, J. 1985. Fire history in three vegetation types on the eastern side of the Oregon Cascades. Ph.D. dissertation, Oregon State University, Corvallis, OR.
- BORMANN, F. H., AND G. E. LIKENS. 1979. Catastrophic disturbance and the steady state in northern hardwood forests. *American Scientist* 67:660–669.
- BOWEN, D. E., JR. 1976. Coloniality, reproductive success, and habitat interactions in Upland Sandpiper (*Bartramia longicauda*). Ph.D. dissertation, Kansas State University, Mahattan, KS.
- BOYCE, S. G., AND W. H. MARTIN. 1993. The future of the terrestrial communities of the southeastern United States. Pp. 339–366 in W. H. Martin, S. G. Boyce, and A. C. Echternacht (editors). Biodiversity of the southeastern United States: upland terrestrial communities. John Wiley and Sons, New York, NY.
- BOYD, R. (EDITOR). 1999. Indians, fire and the land in the Pacific Northwest. Oregon State University Press, Corvallis, OR.
- BRAGG, T. B., AND L. C. HULBERT. 1976. Woody plant invasion of unburned Kansas bluestem prairie. *Journal of Range Management* 29:19–24.
- BRAUN, C. E., M. F. BAKER, R. L. ENG, J. S. GASHWILER, AND M. H. SCHROEDER. 1976. Conservation committee report on effects of alteration of sagebrush communities on the associated avifauna. *Wilson Bulletin* 88: 165–171.
- BRAUN, E. L. 1950. Deciduous forests of eastern North America. Blakiston Co., Philadelphia, PA.
- BRAWN, J. D., AND S. K. ROBINSON. 1996. Source-sink population dynamics may complicate the interpretation of long-term census data. *Ecology* 77:3–12.
- BRAWN, J. D., AND D. F. STOTZ. 2000. The importance of the Chicago region and the "Chicago Wilderness" initiative for avian conservation. Pp. 509–522 in J. M. Marzluff, R. Bowman, and R. Donnelly (editors). Avian ecology and conservation in an urbanizing world. Kluwer Academic Publishers, Boston, MA.
- BRAWN, J. D., S. K. ROBINSON, AND F. R. THOMPSON III. 2001. The role of disturbance in the ecology and conservation of birds. *Annual Review of Ecology and Systematics* 32:251–276.

- BREJDA, J. J. 1997. Soil changes following 18 years of protection from grazing in Arizona chaparral. *Southwestern Naturalist* 42:478–487.
- BRENNAN, L. A., J. L. COOPER, K. E. LUCAS, B. D. LEOPOLD, AND G. A. HURST. 1995. Assessing the influence of Red-cockaded Woodpecker colony site management on non-target forest vertebrates in loblolly pine forests of Mississippi: study design and preliminary results. Pp. 309–319 in D. L. Kulhavy, R. G. Hooper, and R. Costa (editors). *Red-cockaded Woodpecker: recovery, ecology, and management*. Stephen F. Austin State University, Nacogdoches, TX.
- BRENNAN, L. A., J. M. LEE, E. L. STALLER, S. D. WELLENDOFF, AND R. S. FULLER. 2000. Effects of seasonal fire applications on Northern Bobwhites brood habitat and hunting success. *National Quail Symposium* 4:66–69.
- BRENNER, J. 1991. Southern oscillation anomalies and their relationship to wildfire activity in Florida. *International Journal of Wildland Fire* 1:73–78.
- BRIGGS, J. M., AND D. J. GIBSON. 1992. Effect of fire on tree spatial patterns in a tallgrass prairie landscape. *Bulletin of the Torrey Botanical Club* 119:300–307.
- BRIGGS, J. M., G. A. HOCH, AND L. C. JOHNSON. 2002. Assessing the rate, mechanisms, and consequences of the conversion of tallgrass prairie to *Juniperus virginiana* forest. *Ecosystems* 5:578–586.
- BRIGGS, J. M., T. R. SEASTEDT, AND D. J. GIBSON. 1989. Comparative analysis of temporal and spatial variability in above-ground production in a deciduous forest and prairie. *Holarctic Ecology* 12:130–136.
- BROCKWAY, D. G., C. TOPIK, M. A. HEMSTROM, AND W. H. EMMINGHAM. 1983. Plant association and management guide for the Pacific silver fir zone, Gifford Pinchot National Forest. USDA Forest Service Technical Report R6-ECOL-130a. USDA Forest Service, Pacific Northwest Region, Portland, OR.
- BROOKS, M. L., AND D. A. PYKE. 2001. Invasive plants and fire in the deserts of North America. Tall Timbers Research Station Miscellaneous Publication No. 11: 1–14.
- BROSE, P. H., AND D. H. VAN LEAR. 1998. Responses of hardwood advance regeneration to seasonal prescribed fires in oak-dominated shelterwood stands. *Canadian Journal of Forest Research* 28:331–339.
- BROSE, P. H., AND D. H. VAN LEAR. 1999. Effects of seasonal prescribed fires on residual overstory trees in oak-dominated shelterwood stands. *Southern Journal of Applied Forestry*. 23:88–93.
- BROSE, P. H., T. SCHULER, D. VAN LEAR, AND J. BERST. 2001. Bringing fire back: the changing regimes of the Appalachian mixed-oak forests. *Journal of Forestry* 99:30–35.
- BROWN, D. E. (EDITOR). 1982a. Biotic communities of the American southwest—United States and Mexico. *Desert Plants* 4:1–342.
- BROWN, D. E., AND R. DAVIS. 1998. Terrestrial bird and mammal distribution changes in the American Southwest, 1890–1990. Pp. 47–64 in B. Tellman, D. M. Finch, C. Edminster, and R. Hamre (editors). The future of arid grasslands: identifying issues, seeking solutions. USDA Forest Service Proceedings RMRS-P-3. USDA Forest Service, Rocky Mountain Research Station, Fort Collins, CO.
- BROWN, D. E., AND R. A. MINNICH. 1986. Fire and changes in creosote bush scrub of the western sonoran desert, California. *American Midland Naturalist* 116: 411–422.
- BROWN, J. K. 2000. Introduction and fire regimes. Pp. 1–8 in J. K. Brown and J. Kapler (editors). *Wildland fire in ecosystems: effects of fire on flora*. USDA Forest Service General Technical Report RMRS-GTR42, Volume 2. USDA Forest Service, Rocky Mountain Research Station Ogden, UT.
- BROWN, P. M. 2004. Fire, climate, and forest structure in ponderosa pine forests of the Black Hills. Ph.D. dissertation, Colorado State University, Fort Collins, CO.
- BROWN, P. M., AND C. SIEG. 1996. Fire history in interior ponderosa pine forests of the Black Hills, South Dakota, USA. *International Journal of Wildland Fire* 6:97–105.
- BROWN, P. M., AND C. SIEG. 1999. Historical variability in fire at the ponderosa pine-northern Great Plains prairie ecotone, southeastern Black Hills, South Dakota. *Ecoscience* 6:539–547.
- BROWN, P. M., M. R. KAUFMAN, AND W. D. SHEPPARD. 1999. Long-term, landscape patterns of past fire events in a montane ponderosa pine forest of central Colorado. *Landscape Ecology* 14:513–532.
- BROWN, P. M., M. W. KAYE, AND D. BUCKLEY. 1999. Fire history in Douglas-fir and coast redwood forests at Point Reyes National Seashore, California. *Northwest Science* 73:205–216.
- BROWN, R. L. 1982b. Effects of livestock grazing on Mearns Quail in southeastern Arizona. *Journal of Range Management* 35:727–732.
- BUFFINGTON, L. C., AND C. H. HERBEL. 1965. Vegetation changes on a semidesert grassland range from 1858 to 1963. *Ecological Monographs* 35:139–164.
- BUNTING, S. C., B. M. KILGORE, AND C. L. BUSHLEY. 1987. Guidelines for prescribed burning sagebrush-grass rangelands in the northern Great Basin. USDA Forest Service General Technical Report INT-231. USDA Forest Service, Intermountain Research Station, Ogden, UT.
- BURCHAM, L. T. 1957. California range land. State of California, Division of Forestry, Department of Natural Resources, Sacramento, CA.
- BURGER, L. W., C. HARDY, AND J. BEIN. 1998. Effects of prescribed fire and midstory removal on breeding bird communities in mixed pine-hardwood ecosystems of southern Mississippi. Tall Timbers Fire Ecology Conference 20:107–113.
- BURGESS, T. L. 1995. Desert grassland, mixed shrub savanna, shrub steppe, or semidesert shrub? The dilemma of coexisting growth forms. Pp. 31–67 in M. P. McClaran, and T. R. Van Devender (editors). *The desert grassland*. University of Arizona Press, Tucson, AZ.
- BURGESS, T. L., J. E. BOWERS, AND R. M. TURNER. 1991.

- Exotic plants at the Desert Laboratory, Tucson, Arizona. *Madroño* 38:96–114.
- BUSCH, D. E. 1995. Effects of fire on southwestern riparian plant community structure. *Southwestern Naturalist* 40:259–267.
- BUSCH, D. E., AND S. D. SMITH. 1995. Mechanisms associated with decline of woody species in riparian ecosystems of the southwestern U. S. *Ecological Monographs* 65:347–370.
- BUSS, I. O., AND A. S. HAWKINS. 1939. The Upland Plover at Faville Grove, Wisconsin. *Wilson Bulletin* 51: 202–220.
- BYRNE, M. W. 2002. Habitat use by female Greater Sage Grouse in relation to fire at Hart Mountain National Antelope Refuge, Oregon. M.S. thesis, Oregon State University, Corvallis, OR.
- BYRNE, R., E. EDLUND, AND S. MENSING. 1991. Holocene changes in the distribution and abundance of oaks in California. Albany, CA. Pp. 182–188 in R. B. Standiford (editor). *Proceedings of the symposium on oak woodlands and hardwood rangeland management*. USDA Forest Service General Technical Report PSW-126. USDA Forest Service, Pacific Southwest Research Station, Berkeley, CA.
- CALIFORNIA PARTNERS IN FLIGHT. 2002a Version 1.0. The draft coniferous forest bird conservation plan: a strategy for protecting and managing coniferous forest habitats and associated birds in California. (J. Robinson, and J. Alexander, [lead authors]). Point Reyes Bird Observatory, Stinson Beach, CA.
- CALIFORNIA PARTNERS IN FLIGHT. 2002b Version 2.0. The oak woodland bird conservation plan: a strategy for protecting and managing oak woodland habitats and associated birds in California. Point Reyes Bird Observatory, Stinson Beach, CA. <<http://www.prbo.org/calpif/>> (12 June 2004).
- CAREY, A. B., M. HARDT, S. P. HORTON, AND B. L. BISWELL. 1991. Spring bird communities in the Oregon Coast Range. Pp. 127–137 in L. F. Ruggiero, K. B. Aubry, A. B. Carey, and M. H. Huff (technical coordinators). *Wildlife and vegetation of unmanaged Douglas-fir forests*. USDA Forest Service General Technical Report PNW-GTR-285. USDA Forest Service, Pacific Northwest Research Station, Portland, OR.
- CARLE, D. 2002. *Burning questions: America's fight with nature's fire*. Praeger, Westport, CT.
- CARMICHAEL, R. S., O. D. KNIFE, C. P. PASE, AND W. W. BRADY. 1978. Arizona chaparral: plant associations and ecology. USDA Forest Service Research Paper RM-202. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- CARTHERS, S. W., R. R. JOHNSON, AND S. W. AITCHISON. 1974. Population structure and social organization of southwestern riparian birds. *American Zoologist* 14: 97–108.
- CARTRON, J. E., AND D. M. FINCH (TECHNICAL EDITORS). 2000. Ecology and conservation of the Cactus Ferruginous Pygmy-Owl in Arizona. USDA Forest Service General Technical Report RMRS-GTR-43. USDA Forest Service, Rocky Mountain Research Station, Ogden, UT.
- CARTRON, J. E., S. H. STOLESON, P. L. L. STOLESON, AND D. W. SHAW. 2000. Riparian areas. Pp. 281–327 in R. Jemison, and C. Raish (editors). *Livestock management in the American southwest*. Elsevier, Amsterdam, The Netherlands.
- CASTRALE, J. J. 1982. Effect of two sagebrush control methods on nongame birds. *Journal of Wildlife Management* 46:945–952.
- CASWELL, H. 2001. *Matrix population models: construction, analysis, and interpretation*. Second Edition. Sinauer Associates, Inc., Sunderland, MA.
- CATON, E. L. 1996. Effects of fire and salvage logging on the cavity-nesting bird community in northwestern Montana. Ph.D. dissertation, University of Montana, Missoula, MT.
- CHABOT, M., R. GAGNON, L. GABORIAULT, AND R. GIGUERE. 1997. Analyse du régime de protection des forêts. Technical Report, Ministry of Natural Resources, Quebec. Quebec, Canada.
- CHAMBERS, C. L., W. C. MCCOMB, AND J. C. TAPPEINER II. 1999. Breeding bird responses to three silvicultural treatments in the Oregon Coast Range. *Ecological Applications* 9:171–185.
- CHANDLER, C. R., AND M. S. WOODREY. 1995. Status of Henslow's Sparrows during winter in coastal Mississippi. *Mississippi Kite* 25:20–24.
- CHANG, C. R. 1996. Ecosystem responses to fire and variations in fire regimes. Pp. 1071–1099 in *Status of the Sierra Nevada: Sierra Nevada ecosystem project final report to Congress Volume II*. Wildland Resources Center Report No. 37. Center for Water and Wildland Resources, University of California, Davis, CA.
- CHAPMAN, H. H. 1932. Is the longleaf type a climax? *Ecology* 13:328–334.
- CHAPPELL, C. B., AND J. K. AGEE. 1996. Fire severity and tree seedling establishment in *Abies magnifica* forests, southern Cascades, Oregon. *Ecological Applications* 6:628–640.
- CHAPPELL, C. B., AND J. KAGAN. 2000. Ceanothus-Manzanita shrublands. Pp. 43–44 in D. H. Johnson, and T. A. O'Neil (managing directors). *Wildlife-habitat relationships in Oregon and Washington*. University of Oregon Press, Corvallis, OR.
- CHRISTENSEN, N. L., JR. 1981. Fire regimes in southeastern ecosystems. Pp. 112–136 in H. A. Mooney, T. M. Bonnicksen, N. L. Christensen, Jr., J. E. Lotan, and W. A. Reiners (editors). *Proceedings of the Conference Fire Regimes and Ecosystem Properties*. General Technical Report, WO-26. USDA Forest Service, Washington, DC.
- CISSEL, J. H., F. J. SWANSON, G. E. GRANT, D. H. OLSON, S. V. GREGORY, S. L. GARMAN, L. H. ASHKENAS, M. G. HUNTER, J. A. KERTIS, J. H. MAYO, M. D. McSWAIN, S. G. SWETLAND, K. A. SWINDLE, AND D. O. WALLIN. 1998. A landscape plan based on historical fire regimes for a managed forest ecosystem: the Augusta Creek study. USDA Forest Service General Technical Report PNW-

- GTR-42. USDA Forest Service, Pacific Northwest Research Station, Portland, OR.
- CLARK, D. L., AND M. V. WILSON. 2001. Fire, mowing, and removal of woody species in restoring a native prairie in the Willamette Valley of Oregon. *Wetlands* 21: 135–144.
- CLARK, J. S., AND P. D. ROYALL. 1995. Transformation of a northern hardwood forest by aboriginal (Iroquois) fire: charcoal evidence from Crawford Lake, Ontario, Canada. *Holocene* 5:1–9.
- CLARK, R. G., K. A. HOBSON, J. D. NICHOLS, AND S. BEARHOP. 2004. Avian dispersal and demography: scaling up to the landscape and beyond. *Condor* 106:717–719.
- CLEWELL, A. F. 1989. Natural history of wiregrass (*Aristida stricta* Michx., Gramineae). *Natural Areas Journal* 9: 223–233.
- COLLINS, S. L. 1987. Interaction of disturbances in tallgrass prairie: A field experiment. *Ecology* 68:1243–1250.
- COLLINS, S. L. 1990. Introduction: Fire as a natural disturbance in tallgrass prairie ecosystems. Pp. 3–7 in S. L. Collins, and L. L. Wallace (editors). *North American tallgrass prairies*. University of Oklahoma Press, Norman, OK.
- COMER, P., J. KAGAN, M. HEINER, AND C. TOBALSKE. 2002. Current distribution of sagebrush and associated vegetation in the western United States. <<http://SAGEMAP.wr.usgs.gov>> (28 June 2004).
- CONNELLY, J. W., K. P. REESE, R. A. FISCHER, AND W. L. WAKKINEN. 2000. Response of a Sage Grouse breeding population to fire in southeastern Idaho. *Wildlife Society Bulletin* 28:90–96.
- CONNER, R. C., J. D. BORN, A. W. GREEN, AND R. A. O'BRIEN. 1990. Forest resources of Arizona. USDA Forest Service Resource Bulletin INT-69. USDA Forest Service, Intermountain Forest and Range Experiment Station, Ogden, UT.
- CONNER, R. N., D. C. RUDOLPH, AND J. R. WALTERS. 2001. *The Red-cockaded Woodpecker*. University of Texas Press, Austin, TX.
- COOPER, C. F. 1960. Changes in vegetation, structure, and growth of southwestern ponderosa pine forests since white settlement. *Ecological Monographs* 30: 129–164.
- COOPER, C. F. 1961. Pattern in ponderosa pine forests. *Ecology* 42:493–499.
- COOPER, W. S. 1922. The broad-sclerophyll vegetation of California: An ecological study of the chaparral and its related communities. Technical Press, Publication No. 319, Carnegie Institution of Washington, Washington, DC.
- COPPEDGE, B. R., D. M. ENGLE, C. S. TOEPPER, AND J. H. SHAW. 1998a. Effects of seasonal fire, bison grazing and climatic variation on tallgrass prairie vegetation. *Plant Ecology* 139:235–246.
- COPPEDGE, B. R., D. M. LESLIE, JR., AND J. H. SHAW. 1998b. Botanical composition of bison diets on tallgrass prairie in Oklahoma. *Journal of Range Management* 51: 379–382.
- COVINGTON, W. W., AND M. M. MOORE. 1994. Southwestern ponderosa pine forest structure: changes since Euro-American settlement. *Journal of Forestry* 92:39–47.
- COVINGTON, W. W., P. Z. FULÉ, M. M. MOORE, S. C. HART, T. E. KOLB, J. N. MAST, S. S. SACKET, AND M. R. WAGNER. 1997. Restoring ecosystem health in ponderosa pine forests of the Southwest. *Journal of Forestry* 95:23–29.
- CRONON, W. 1983. *Changes in the land: Indians, colonists, and the ecology of New England*. Hill and Wang, New York, NY.
- CROW, T. R. 1988. Reproductive mode and mechanisms for self-replacement of northern red oak (*Quercus rubra*)—a review. *Forest Science* 34:19–40.
- CROWLEY, C. M., AND J. W. CONNELLY. 1996. Sage-Grouse population and habitat trends in southeastern Idaho and southwestern Montana. Gamebird Research Group Report, Idaho Department of Fish and Game, Pocatello, ID.
- CULLY, J. F., JR., AND H. L. MICHAELS. 2000. Henslow's Sparrow habitat associations on Kansas tallgrass prairie. *Wilson Bulletin* 112:115–123.
- CUMMING, S. G. 2001. Forest type and wildfire in the Alberta boreal mixed woodland: what do fires burn? *Ecological Applications* 11:97–110.
- CUTTER, B. E., AND R. P. GUYETTE. 1994. Fire frequency on an oak-hickory ridgetop in the Missouri Ozarks. *American Midland Naturalist* 132:393–398.
- D'ANTONIO, C. M. 2000. Fire, plant invasions, and global changes. Pp. 65–93 in H. A. Mooney, and R. J. Hobbs (editors). *Invasive species in a changing world*. Island Press, Washington, DC.
- D'ANTONIO, C. M., AND P. M. VITOUSEK. 1992. Biological invasions by exotic grasses, the grass/fire cycle, and global change. *Annual Review of Ecology and Systematics* 23:63–87.
- DAUBENMIRE, R. F. 1943. Vegetational zonation in the Rocky Mountains. *Botanical Review* 9:325–393.
- DAUBENMIRE, R. 1968. Ecology of fire in grasslands. Pp. 209–266 in J. B. Cragg (editor). *Advances in ecological research* Vol. 5. Academic Press, London, UK.
- DAVIS, E. A. 1989. Prescribed fire in Arizona chaparral: effects on stream water quality. *Forest Ecology and Management* 26:189–206.
- DAVIS, F. W., D. M. STOMS, A. D. HOLLANDER, K. A. THOMAS, P. A. STINE, D. ODION, M. I. BORCHERT, J. H. THORNE, M. V. GRAY, R. E. WALKER, K. WARNER, AND J. GRAAE. 1998. The California gap analysis project—final report. University of California, Santa Barbara, CA. <http://www.biogeog.ucsb.edu/projects/gap/gap_home.html> (12 June 2004).
- DAVIS, J. N. 1999. Lawrence's Goldfinch (*Carduelis lawrencei*). In A. Poole, and F. Gill (editors). *The Birds of North America*, No. 480. The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists Union, Washington, DC.
- DAVIS, M. A., D. W. PETERSON, P. B. REICH, M. CROZIER, T. QUERY, E. MITCHELL, J. HUNTINGTON, AND P. BAZAKAS. 2000. Restoring savanna using fire: impact on the breeding bird community. *Restoration Ecology* 8:30–40.

- DAVIS, P. R. 1976. Response of vertebrate fauna to forest fire and clearcutting in south central Wyoming. Ph.D. dissertation, University of Wyoming, Laramie, WY.
- DAVIS, W. A., AND S. M. RUSSELL. 1990. Birds in southeastern Arizona. Tucson Audubon Society, Tucson, AZ.
- DAY, G. M. 1953. The Indian as an ecological factor in the northeastern forest. *Ecology* 34:329–346.
- DECHANT, J. A., M. L. SONDRAL, D. H. JOHNSON, L. D. IGL, C. M. GOLDADE, M. P. NENNEMAN, AND B. R. EULISS. 2001. Effects of management practices on grassland birds: Grasshopper Sparrow. Northern Prairie Wildlife Research Center, Jamestown, ND. <<http://www.npwrc.usgs.gov/resource/literatr/grasbird/grasshop/grasshop.htm>> (17 February 2000).
- DEGRAAF, R. M. 1991. Forest and rangeland birds of the United States. USDA Forest Service Agriculture Handbook 688. Washington, DC.
- DELCOURT, H. R., AND P. A. DELCOURT. 2000. Eastern deciduous forests. Pp. 358–395 in M. G. Barbour, and W. D. Billings (editors). *North American terrestrial vegetation*. Cambridge University Press, Cambridge, UK.
- DELCOURT, P. A., AND H. R. DELCOURT. 1998. The influence of prehistoric human-set fires on oak-chestnut forests in the southern Appalachians. *Castanea* 63:337–345.
- DELLASALA, D. A., R. G. ANTHONY, T. A. SPECIES, AND K. A. ENGEL. 1998. Management of Bald Eagle communal roosts in fire-adapted mixed-conifer forests. *Journal of Wildlife Management* 62:322–333.
- DELLASALLA, D. A., J. E. WILLIAMS, C. DEACON WILLIAMS, AND J. F. FRANKLIN. 2004. Beyond smoke and mirrors: a synthesis of fire policy and science. *Conservation Biology* 18:976–986.
- DENEVAN, W. M. (EDITOR). 1992. *The native population of the Americas in 1492*. University of Wisconsin Press, Madison, WI.
- DESELN, H. R., AND N. MURDOCK. 1993. Grass-dominated communities. Pp. 87–141 in W. H. Martin, S. G. Boyce, and A. C. Echternacht (editors). *Biodiversity of the southeastern United States: upland terrestrial communities*. John Wiley and Sons, Inc., New York, NY.
- DETTMERS, R. P., AND J. BART. 1999. A GIS modeling method applied to predicting forest songbird habitat. *Ecological Applications* 9:152–163.
- DEY, D. C., AND R. P. GUYETTE. 2000. Anthropogenic fire history and red oak forests in south-central Ontario. *Forestry Chronicle* 76:339–347.
- DICKMAN, A., AND S. COOK. 1989. Fire and fungus in a mountain hemlock forest. *Canadian Journal of Botany* 67:2005–2016.
- DICKSON, J. G. 1981. Effects of forest burning on songbirds. Pp. 67–72 in G. W. Wood (editor). *Prescribed fire and wildlife in southern forests*. Belle W. Baruch Forestry Science Institute, Clemson University, Georgetown, SC.
- DIEM, K. L., AND S. I. ZEVELOFF. 1980. Ponderosa pine bird communities. Pp. 170–197 in R. M. DeGraff, and N. G. Tilghman (editors). *Management of western forests and grasslands for nongame birds: workshop proceedings*. USDA Forest Service General Technical Report INT-86. USDA Forest Service, Intermountain Research Station, Ogden, UT.
- DIXON, R. D. 1995. Ecology of White-headed Woodpeckers in the central Oregon Cascades. M.S. thesis, University of Idaho, Moscow.
- DIXON, R. D., AND V. A. SAAB. 2000. Black-backed Woodpecker (*Picoides arcticus*). In A. Poole and F. Gill (editors). *The Birds of North America*, No. 509. The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists' Union, Washington, DC.
- DOBKIN, D. S., AND J. D. SAUDER. 2004. Shrubsteppe landscapes in jeopardy. Distributions, abundances, and the uncertain future of birds and small mammals in the intermountain West. High Desert Ecological Research Institute, Bend, OR.
- DOBLER, F. C., J. EBY, C. PERRY, S. RICHARDSON, AND M. VANDER HAEGEN. 1996. Status of Washington's shrubsteppe ecosystem: extent, ownership, and wildlife/vegetation relationships. Research Report. Washington Department of Fish and Wildlife, Olympia, WA.
- DODGE, J. M. 1975. Vegetational changes associated with land use and fire history in San Diego County. Ph.D. dissertation, University of California, Riverside, CA.
- DOMBECK, M. P., J. E. WILLIAMS, AND C. A. WOOD. 2004. Wildfire policy and public lands: integrating scientific understanding with social concerns across landscapes. *Conservation Biology* 18:883–889.
- DRAPEAU, P., A. NAPPI, J.-F. GIROUX, A. LEDUC, AND J.-P. SAVARD. 2002. Distribution patterns of birds associated with snags in natural and managed eastern boreal forests. Pp. 193–205. in B. Laudenslayer, W. F. Laudenslayer, Jr., P. J. Shea, B. E. Valentine, C. P. Weatherspoon, and T. E. Lisle (technical coordinators). *Ecology and management of dead wood in western forests*. USDA Forest Service General Technical Report PSW-GTR 181. USDA Forest Service, Pacific Southwest Research Station, Albany, CA.
- DREW, M. B., L. K. KIRKMAN, AND A. K. GHOLSON, JR. 1998. The vascular flora of Ichauway, Baker County, Georgia: a remnant longleaf/wiregrass ecosystem. *Castanea* 6:31–24.
- DREWA, P. B., AND K. M. HAVSTAD. 2001. Effects of fire, grazing, and the presence of shrubs on Chihuahuan desert grasslands. *Journal of Arid Environments* 48:429–443.
- DROEGE, S., AND J. R. SAUER. 1989. North American breeding bird survey annual summary 1988. USDI Fish Wildlife Service, Biological. Report 89:1–16.
- DUEBBERT, H. F., AND J. T. LOKEMOEN. 1977. Upland nesting of American Bitterns, Marsh Hawks, and Short-eared Owls. *Prairie Naturalist* 9:33–40.
- DUNCAN, D. A., AND W. J. CLAWSON. 1980. Livestock utilization of California's oak woodlands. Pp. 306–313 in T. R. Plumb (technical coordinator). *Proceedings of the symposium on the ecology, management, and utilization of California oaks*. USDA Forest Service General Technical Report PSW-44. USDA Forest Service, Pacific Southwest Research Station, Berkeley, CA.

- DUNNING, J. B. 1993. Bachman's Sparrow (*Aimophila aestivalis*). In A. Poole, P. Stettenheim, and F. Gill (editors). The Birds of North America, No. 38. The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists' Union, Washington, DC.
- DUNWIDDIE, P. W. 1989. Forest and heath: the shaping of vegetation on Nantucket Island. *Journal of Forest History* 33:126–133.
- DUNWIDDIE, P. W., AND C. CALJOUW. 1990. Prescribed burning and mowing of coastal heathlands and grasslands in Massachusetts. *New York State Museum Bulletin* 471:271–275.
- DWIRE, K. A., AND J. B. KAUFFMAN. 2003. Fire and riparian ecosystems in landscapes of the western USA. *Forest Ecology and Management* 178:61–74.
- DWYER, J. K. 2000. Response of secondary cavity-nesting birds to wildfire in ponderosa pine forests of northern Arizona. M. S. thesis, Northern Arizona University, Flagstaff, AZ.
- DWYER, J. K., AND W. M. BLOCK. 2000. Effects of wildfire on densities of secondary cavity nesting birds in ponderosa pine forests of northern Arizona. *Tall Timbers Fire Ecology Conference Proceedings* 21:151–156.
- EDWARDS, M. B. 1957. California black oak—its management and economic possibilities. *Journal of Forestry* 55:506–510.
- EHLE, D. S., AND W. L. BAKER. 2003. Disturbance and stand dynamics in ponderosa pine forests in Rocky Mountain National Park, USA. *Ecological Monographs* 73:543–566.
- EHRlich, P. R., D. S. DOBKIN, AND D. WHEYE. 1988. The birder's handbook: A field guide to the natural history of North American birds. Simon and Schuster/Fireside Books, New York, NY.
- ELLIOTT, K. J., R. L. HENDRICK, A. E. MAJOR, J. M. VOSE, AND W. T. SWANK. 1999. Vegetation dynamics after prescribed fire in the southern Appalachians. *Forest Ecology and Management* 114:199–213.
- ELLIS, L. M. 2001. Short-term response of woody plants to fire in a Rio Grande forest, central New Mexico, USA. *Biological Conservation* 97:159–170.
- ELLIS, L. M., C. S. CRAWFORD, AND M. C. MOLLES. 1998. Comparison of litter dynamics in native and exotic riparian vegetation along the Middle Rio Grande of central New Mexico, USA. *Journal of Arid Environments* 38:283–296.
- ENGLE, D. M., AND T. G. BIDWELL. 2001. Viewpoint: the response of central North American prairies to seasonal fire. *Journal of Range Management* 54:2–10.
- ENGLE, D. M., AND J. D. KULBETH. 1992. Growth dynamics of crowns of eastern redcedar at 3 locations in Oklahoma. *Journal of Range Management* 45:301–305.
- ENGLE, D. M., T. G. BIDWELL, AND M. E. MOSELEY. 1996. Invasion of Oklahoma rangelands and forests by eastern redcedar and ashe juniper. Oklahoma Cooperative Extension Service, Circular E-947. Oklahoma State University, Stillwater, OK.
- ENGLE, D. M., R. L. MITCHELL, AND R. L. STEVENS. 1998. Late growing-season fire effects in mid-successional tallgrass prairie. *Journal of Range Management* 51:115–121.
- ENGLE, D. M., J. F. STRITZKE, AND P. L. CLAYPOOL. 1987. Herbage standing crop around eastern redcedar trees. *Journal of Range Management* 40:237–239.
- ENGLE, D. M., J. F. STRITZKE, T. G. BIDWELL, AND P. L. CLAYPOOL. 1993. Late-summer fire and follow-up herbicide treatments in tallgrass prairie. *Journal of Range Management* 46:542–547.
- ENGSTROM, R. T. 1993. Characteristic mammals and birds of longleaf pine forests. The longleaf pine ecosystem: ecology, restoration, and management. *Tall Timbers Fire Ecology Conference* 18:127–138.
- ENGSTROM, R. T., AND W. E. PALMER. In press. Two species in one ecosystem: management of Northern Bobwhite and Red-cockaded Woodpecker in the Red Hills. In C. J. Ralph and T. D. Rich (editors). Bird conservation implementation and integration in the Americas. USDA Forest Service General Technical Report PSW-GTR-191. USDA Forest Service, Pacific Southwest Research Station, Albany, CA.
- ENGSTROM, R. T., R. L. CRAWFORD, AND W. W. BAKER. 1984. Breeding bird populations in relation to changing forest structure following fire exclusion: a 15-year study. *Wilson Bulletin* 96:437–450.
- ENGSTROM, R. T., D. B. MCNAIR, L. A. BRENNAN, C. L. HARDY, AND L. W. BURGER. 1996. Influence on birds of dormant versus lightning-season prescribed fire in longleaf pine forests: experimental design and preliminary results. *North American Wildlife and Natural Resources Conference* 61:200–207.
- ENGSTROM, R. T., S. GILBERT JR., M. L. HUNTER, D. MERRIWETHER, G. J. NOWACKI, AND P. SPENCER. 1999. Practical applications of disturbance ecology to natural resource management. Pp. 313–330 in N. C. Johnson, A. J. Malk, W. T. Sexton, and R. C. Szaro (editors). *Ecological stewardship: a common reference for ecosystem management*. Elsevier Science Ltd., Oxford, UK.
- EVANS, R. A., AND J. A. YOUNG. 1978. Effectiveness of rehabilitation practices following wildfire in degraded big sagebrush-downy brome communities. *Journal of Range Management* 31:185–188.
- EVERETT, R. L., R. SCHELLHAAS, D. KEENUM, D. SPURBECK, AND P. OHLSON. 2000. Fire history in the ponderosa pine/Douglas-fir forests on the east slope of the Washington Cascades. *Forest Ecology and Management* 129:207–225.
- EWING, A. L., AND D. M. ENGLE. 1988. Effects of late summer fire on tallgrass prairie microclimate and community composition. *American Midland Naturalist* 120:212–223.
- FABER-LANGENDOEN D., AND M. A. DAVIS. 1995. Effects of fire frequency on tree canopy cover at Allison Savanna, east-central Minnesota. *Natural Areas Journal* 15:319–328.
- FAHEY, T. J., AND W. A. REINERS. 1981. Fire in the forests of Maine and New Hampshire. *Bulletin of the Torrey Botanical Club* 108:362–373.

- FAHNESTOCK, G. R. 1977. Interaction of forest fires, flora, and fuels in two Cascade Range wildernesses. Ph.D. dissertation, University of Washington, Seattle, WA.
- FERGUSON, J. P. 1998. Prescribed fire on the Apalachicola Ranger District: the shift from dormant season to growing season and effects on wildfire suppression. Tall Timbers Fire Ecology Conference 20:120–126.
- FFOLIOTT, P. F., L. F. DEBANO, M. B. BAKER JR., G. J. GOTTFRIED, G. SOLIS-GARZA, C. B. EDMINSTER, D. G. NEARY, L. S. ALLEN, AND R. H. HAMRE (TECHNICAL COORDINATORS). 1996. Effects of fire on Madrean province ecosystems—a symposium proceedings. USDA Forest Service General Technical Report RM-GTR-289. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- FINCH, D. M., AND R. T. REYNOLDS. 1987. Bird response to understory variation and conifer succession in aspen forests. Pp. 87–96 in J. Emerick, S. Q. Foster, L. Hayden-Wing, J. Hodgson, J. W. Monarch, A. Smith, O. Thorne II, and J. Todd (editors). Issues and technology in the management of impacted wildlife. Thorne Ecological Institute, Boulder, CO.
- FINNEY, M. A., AND R. E. MARTIN. 1989. Fire history in a *Sequoia sempervirens* forest at Salt Point State Park, California. Canadian Journal of Forest Research 19: 1451–1457.
- FINNEY, M. A., AND R. E. MARTIN. 1992. Short fire interval recorded at Annadale State Park, California. Madroño 39:251–262.
- FISCHER, R. A., W. L. WAKKINEN, K. P. REESE, AND J. W. CONNELLY. 1997. Effects of prescribed fire on movements of female Sage Grouse from breeding to summer ranges. Wilson Bulletin 109:82–91.
- FISCHER, R. A., A. D. APA, W. L. WAKKINEN, K. P. REESE, AND J. W. CONNELLY. 1996. An investigation on fire effects with xeric Sage-Grouse brood habitat. Journal of Range Management 49:194–198.
- FLANNIGAN, M., Y. BERGERON, O. ENGELMARK, AND M. WOTTON. 1998. Future wildfire in circumboreal forests in relation to global warming. Journal of Vegetation Science. 9:469–476.
- FLOYD, M. L., W. H. ROMME, AND D. D. HANNA. 2000. Fire history and vegetation pattern in Mesa Verde National Park, Colorado, USA. Ecological Applications 10: 1666–1680.
- FONDA, R. W. 1974. Forest succession in relation to river terrace development in Olympic National Park, Washington. Ecology 55:927–942.
- FONDA, R. W., AND L. C. BLISS. 1969. Forest vegetation of the montane and subalpine zones, Olympic Mountains, Washington. Ecological Monographs 39:271–301.
- FOREST ECOSYSTEM MANAGEMENT ASSESSMENT TEAM. 1993. Forest ecosystem management: an ecological, economic, and social assessment. Report of the Forest Ecosystem Management Assessment Team. U.S. Government Printing Office 1993-793-071. USDA, Forest Service; USDI Fish and Wildlife Service, Bureau of Land Management, and National park Service; USDC National Oceanic and Atmospheric Administration and National Marine Fisheries Service; and the U.S. Environmental Protection Agency. U.S. Government Printing Office, Washington, DC.
- FOSTER, D. R. 1983. The history and pattern of fire in the boreal forest of southeastern Labrador. Canadian Journal of Botany 61:2459–2471.
- FRANKLIN, J. F., AND C. T. DYRNESS. 1973. Natural vegetation of Oregon and Washington. USDA Forest Service General Technical Report PNW-GTR-8. USDA Forest Service, Pacific Northwest Research Station, Portland, OR.
- FRANKLIN, J. F., AND C. T. DYRNESS. 1988. Natural Vegetation of Oregon and Washington. Oregon State University Press, Corvallis OR.
- FRANKLIN, J. F., AND R. G. MITCHELL. 1967. Successional status of subalpine fir in the Cascade Range. USDA Forest Service Research Paper PNW-RP-46. USDA Forest Service, Pacific Northwest Research Station, Portland, OR.
- FRANKLIN, J. F., W. H. MOIR, M. A. HEMSTROM, S. E. GREENE, AND B. G. SMITH. 1988. The forest communities of Mount Rainier National Park. Scientific Monograph Series 19. USDI National Park Service, Washington DC.
- FRETWELL, S. 1977. Is the Dickcissel a threatened species? American Birds 31:923–932.
- FROST, E. J., AND R. SWEENEY. 2000. Fire regimes, fire history, and forest conditions in the Klamath-Siskiyou region: An overview and synthesis of knowledge. World Wildlife Fund, Ashland, OR.
- FROST, C. C. 1998. Presettlement fire frequency regimes of the United States: a first approximation. Tall Timbers Fire Ecology Conference 20:70–81.
- FROST, C. C., J. WALKER, AND R. K. PEET. 1986. Fire-dependent savannas and prairies of the southeast: original extent, preservation status and management problems. Pp. 348–357 in D. L. Kulhavy, and R. N. Conner (editors). Wilderness and natural areas management in the eastern United States: a management challenge. Stephen F. Austin State University, School of Forestry, Center for Applied Studies, Nacogdoches, TX.
- FRY, D. L. 2002. Effects of a prescribed fire on oak woodland stand structure. Pp. 235–242 in R. Standiford, D. McCreary, and K. L. Purcell (technical coordinators), Proceedings of the fifth symposium on oak woodlands: oaks in California's changing landscape. USDA Forest Service General Technical Report PSW-184. USDA Forest Service, Pacific Southwest Research Station, Albany, CA.
- FUHLENDORF, S. D., AND D. M. ENGLE. 2001. Restoring heterogeneity on rangelands: Ecosystem management based on evolutionary grazing patterns. BioScience 51: 625–632.
- FULÉ, P. Z., W. W. COVINGTON, M. M. MOORE, T. A. HEINLEIN, AND A. E. M. WALTZ. 2002. Natural variability in forests of Grand Canyon, USA. Journal of Biogeography 29:31–47.
- GAINES, W. L., R. A. STRAND, AND S. D. PIPER. 1997. Effects of the Hatchery Complex Fires on the Northern

- Spotted Owls in the eastern Washington Cascades. Pp. 123–130 in J. M. Greenlee (editor). Proceedings, First conference on fire effects on rare and endangered species and habitats. International Association of Wildland Fire, Fairfield, WA.
- GARRETT, K. L., M. G. RAPHAEL, AND R. D. DIXON. 1996. White-headed Woodpecker (*Picoides albolarvatus*). In A. Poole, and F. Gill (editors). The Birds of North America, No. 252. The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists' Union, Washington, DC.
- GAYTON, A. H. 1948. Yokuts and western Mono ethnography II: northern foothill Yokuts and western Mono. Anthropological Records Vol. 10, No. 2. University of California Press, Berkeley, CA.
- GEDNEY, D. R., D. L. AZUMA, C. L. BOLSINGER, AND N. MCKAY. 1999. Western juniper in eastern Oregon. USDA Forest Service General Technical Report PNW-GTR-464. USDA Forest Service, Pacific Northwest Research Station, Portland, OR.
- GERMAINE, H. L., AND S. S. GERMAINE. 2002. Forest restoration treatment effects on the nesting success of Western Bluebirds (*Sialia mexicana*). Restoration Ecology 10:362–367.
- GIBSON, D. J. 1988. Regeneration and fluctuation of tallgrass prairie vegetation in response to burning frequency. Bulletin of the Torrey Botanical Club 115:1–12.
- GIBSON, D. J., AND L. C. HULBERT. 1987. Effects of fire, topography and year-to-year climatic variation on species composition in tallgrass prairie. Vegetatio 72: 175–185.
- GILBERT, F. F., AND R. ALLWINE. 1991. Spring bird communities in the Oregon Cascade Range. Pp. 145–158 in L. F. Ruggiero, K. B. Aubry, A. B. Carey, and M. H. Huff (technical coordinators). Wildlife and vegetation of unmanaged Douglas-fir forests. USDA Forest Service General Technical Report PNW-GTR-285. USDA Forest Service, Pacific Northwest Research Station, Portland, OR.
- GILL, M. A., AND M. A. MCCARTHY. 1998. Intervals between prescribed fires in Australia: what intrinsic variation should apply? Biological Conservation 85:161–169.
- GLITZENSTEIN, J. S., W. J. PLATT, AND D. R. STRENG. 1995. Effects of fire regime and habitat on tree dynamics in north Florida longleaf pine savannas. Ecological Monographs 65:441–476.
- GOGGANS, R., R. D. DIXON, AND L. C. SEMINARAQ. 1988. Habitat use by Three-toed and Black-backed woodpeckers, Deschutes National Forest, Oregon Technical Report 87-3-02, Oregon Department of Fish and Wildlife, Portland, OR.
- GORDON, C. E. 2000. Fire and cattle grazing on wintering sparrows in Arizona grasslands. Journal of Range Management 53:384–389.
- GOSZ, R. J., AND J. R. GOSZ. 1996. Species interactions on the biome transition zone in New Mexico: response of blue grama (*Bouteloua gracilis*) and black grama (*Bouteloua eripoda*) to fire and herbivory. Journal of Arid Environments 34:101–114.
- GRAHAM, R. T., S. MCCAFFREY, AND T. B. JAIN. 2004. Science basis for changing forest structure to modify wildfire behavior and severity. USDA Forest Service General Technical Report RMRS-GTR-120. USDA Forest Service, Rocky Mountain Research Station, Fort Collins, CO.
- GREEN, G. A., AND R. G. ANTHONY. 1989. Nesting success and habitat relationships of Burrowing Owls in the Columbia Basin, Oregon. Condor 91:347–354.
- GREENE, S. W. 1931. The forest that fire made. American Forests 37:583–584.
- GRIFFIN, J. R. 1976. Regeneration in *Quercus lobata* savannas, Santa Lucia Mountains, California. American Midland Naturalist 95:422–435.
- GRIFFIN, J. R. 1987. Landscape disturbance and success of natural oak regeneration. Pp. 38–43 in P. Bowler, and S. Brown (editors). Proceedings of the California oak heritage conservation conference. Sea and Sage Audubon Society, Santa Ana, CA.
- GRIFFIN, J. R. 1988. Oak woodland. Pp. 383–415 in M. G. Barbour, and J. Major (editors). Terrestrial vegetation of California. California Native Plant Society, Special Publication Number 9, Davis, CA.
- GRIFFIS, K. 1999. Bird use of small aspen stands in a northern Arizona ponderosa pine forest. M. S. thesis, Northern Arizona University, Flagstaff, AZ.
- GRIFFITH, D. M., D. M. DIGIOVANNI, T. L. WITZEL, AND E. H. WARTON. 1993. Forest statistics for Ohio, 1991. USDA Forest Service Resource Bulletin NE-128. USDA Forest Service, Northeastern Forest Experiment Station, Radnor, PA.
- GRISSINO-MAYER, H. D., AND T. W. SWETNAM. 2000. Century scale changes in fire regimes and climate in the Southwest. The Holocene 10:207–214.
- GRISSINO-MAYER, H. D., C. H. BAISAN, AND T. W. SWETNAM. 1995. Fire history of the Pinaleno Mountains of southeastern Arizona. Pp. 399–407 in L. F. DeBano, P. F. Ffolliott, A. Ortega-Rubio, G. J. Gottfried, R. H. Hamre, and C. B. Edminster (technical coordinators). Biodiversity and management of Madrean Archipelago: the sky islands of southwestern United States and northwestern Mexico. USDA Forest Service General Technical Report RM-GTR-264. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- GROSS, A. O. 1932. Heath Hen. Pp. 264–280 in A. C. Bent (editor). Life histories of North American gallinaceous birds. U.S. National Museum Bulletin 162, Washington, DC.
- GRUELL, G. E. 1983. Fire and vegetative trends in the northern Rockies: Interpretations from 1871–1982. USDA Forest Service General Technical Report INT-158. USDA Forest Service, Intermountain Research Station, Ogden, UT.
- GUYETTE, R. P., AND B. E. CUTTER. 1991. Tree-ring analysis of fire history of a post oak savanna in the Missouri Ozarks. Natural Areas Journal 11:93–99.
- GUYETTE, R. P., D. C. DEY, AND M. C. STAMBAUGH. 2003. Fire and human history of a barren-forest mosaic in

- southern Indiana. *American Midland Naturalist* 149: 21–34.
- GUYETTE, R. P., R. M. MUZIKA, AND D. C. DEY. 2002. Dynamics of an anthropogenic fire regime. *Ecosystems* 5:472–486.
- HABECK, J. R. 1961. The original vegetation of the mid-Willamette Valley, Oregon. *Northwest Science* 35: 65–77.
- HAGAR, J. C., AND M. A. STERN. 2001. Avifauna in oak woodlands of the Willamette Valley, Oregon. *Northwestern Naturalist* 82:12–25.
- HAGGARD, M., AND W. L. GAINES. 2001. Effects of stand-replacement fire and salvage logging on a cavity-nesting bird community in eastern Cascades, Washington. *Northwest Science* 75:387–396.
- HAGGERTY, P. K. 1994. Damage and recovery in southern Sierra Nevada foothill oak woodland after a severe ground fire. *Madroño* 41:185–198.
- HALL, F. C. 1977. Ecology of natural underburning in the Blue Mountains of Oregon. USDA Forest Service Technical Report R6-ECOL-79-001. USDA Forest Service, Pacific Northwest Region, Portland, OR.
- HALL, F. C. 1983. Ecology of grand fir. Pp. 43–52 in C. D. Oliver, and R. M. Kenady, (editors). *Proceedings of the biology and management of true fir in the Pacific Northwest symposium*. University of Washington, Seattle, WA.
- HALL, L. S., M. L. MORRISON, AND W. M. BLOCK. 1997. Songbird status and roles. Pp. 69–88 in W. M. Block, and D. M. Finch (technical editors). *Songbird ecology in southwestern ponderosa pine forests: a literature review*. USDA Forest Service General Technical Report RM-GTR-292. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- HAMILTON, R. G. 1996. Using fire and bison to restore a functional tallgrass prairie landscape. *Transactions of the North American Wildlife and Natural Resources Conference* 61:208–214.
- HANN, W. J., AND D. L. BUNNELL. 2001. Fire and land management planning and implementation across multiple scales. *International Journal of Wildland Fire* 10:389–403.
- HANN, W. J., M. J. WISDOM, AND M. M. ROWLAND. 2002. Disturbance departure and fragmentation of natural systems in the interior Columbia Basin. USDA Forest Service Research Paper PNW-RP-545. USDA Forest Service, Pacific Northwest Station, Portland, OR.
- HANN, W. J., J. L. JONES, M. G. KARL, P. F. HESSBURG, R. E. KEAN, D. G. LONG, J. P. MENAKIS, C. H. MCNICOLL, S. G. LEONARD, R. A. GRAVENMIER, AND B. G. SMITH. 1997. An assessment of ecosystem components in the interior Columbia Basin and portions of the Klamath and Great basins, Vol. II. Landscape dynamics of the basin. USDA Forest Service General Technical Report PNW-GTR-405. USDA Forest Service, Pacific Northwest Research Station, Portland, OR.
- HARCOMBE, P. A., J. S. GLITZENSTEIN, R. G. KNOX, S. L. ORZELL, AND E. L. BRIDGES. 1995. Vegetation of the longleaf pine region of the west gulf coastal plain. *Tall Timbers Fire Ecology Conference* 18:83–104.
- HARDY, P. C., AND M. L. MORRISON. 2001. Nest site selection by Elf Owls in the Sonoran desert. *Wilson Bulletin* 113:23–32.
- HARNISS, R. O., AND R. B. MURRAY. 1973. Thirty years of vegetal change following burning of sagebrush-grass range. *Journal of Range Management* 26:322–325.
- HARPER, R. M. 1962. Historical notes on the relation of fires to forests. *Tall Timbers Fire Ecology Conference* 1: 11–29.
- HARRINGTON, C. A., AND M. A. KALLAS. 2002. A bibliography for *Quercus garyana* and other geographically associated and botanically related oaks. USDA Forest Service General Technical Report PNW-GTR-554. USDA Forest Service, Pacific Northwest Research Station, Portland, OR.
- HARRIS, J. A. 1998. The effects of prescribed burning of grasslands on the nesting habitat of the Grasshopper Sparrow on Martha's Vineyard, Massachusetts. M.A. thesis, Harvard University, Cambridge, MA.
- HARRIS, M. A. 1982. Habitat use among woodpeckers in forest burns. M.S. thesis, University of Montana, Missoula, MT.
- HARTNETT, D. C., K. R. HICKMAN, AND L. E. FISCHER WALTER. 1996. Effects of bison grazing, fire, and topography on floristic diversity in tallgrass prairie. *Journal of Range Management* 49:413–420.
- HASSAN, M. A., AND N. E. WEST. 1986. Dynamics of soil seed pools in burned and unburned sagebrush semi-deserts. *Ecology* 67:269–272.
- HEALY, W. H., AND W. J. MCSHEA. 2002. Goals and guidelines for managing oak ecosystems for wildlife. Pp. 333–341 in W. J. McShea, and W. H. Healy (editors). *Oak forest ecosystems: ecology and management for wildlife*. Johns Hopkins University Press, Baltimore, MD.
- HEIN, D. 1980. Management of lodgepole pine for birds. Pp. 238–246 in R. M. DeGraff, and N. G. Tilghman (editors). *Management of western forests and grasslands for nongame birds: workshop proceedings*. USDA Forest Service General Technical Report INT-86. USDA Forest Service, Intermountain Research Station, Ogden, UT.
- HEMSTROM, M. A., AND J. A. FRANKLIN. 1982. Fire and other disturbances of the forests in Mount Rainier National Park. *Quaternary Research* 18:32–51.
- HEMSTROM, M. A., AND S. E. LOGAN. 1986. Plant association and management guide, Suislaw National Forest. USDA Forest Service Technical Report R6-ECOL-220-1986a. USDA Forest, Service Pacific Northwest Region, Portland, OR.
- HEMSTROM, M. A., W. H. EMMINGHAM, N. M. HALVERSON, S. E. LOGAN, AND C. T. TOPIK. 1982. Plant association and management guide for the Pacific silver fir zone, Mt. Hood and Willamette National Forests. USDA Forest Service Technical Report R6-ECOL-100-1982a. USDA Forest Service, Pacific Northwest Region, Portland, OR.
- HEMSTROM, M. A., M. J. WISDOM, W. J. HANN, M. M. ROWLAND, B. C. WALES, AND R. A. GRAVENMIER. 2002.

- Sagebrush-steppe vegetation dynamics and restoration potential in the interior Columbia Basin, U.S.A. *Conservation Biology* 16:1243–1255.
- HENDERSON, J. A. 1973. Composition, distribution, and succession of subalpine meadows in Mount Rainier National Park, Washington. Ph.D. dissertation, Oregon State University, Corvallis, OR.
- HENDERSON, J. A. 1982. Ecology of subalpine fir. Pp. 53–58. *in* C. D. Oliver, and R. M. Kenady (editors). *Proceedings of the biology and management of true fir in the Pacific Northwest symposium*. University of Washington, Seattle, WA.
- HENDERSON, J. A., D. H. PETER, R. D. LESHNER, AND D. C. SHAW. 1989. Forested plant associations of the Olympic National Forest. USDA Forest Service Technical Report R6 ECOL TP-001-88. USDA Forest Service, Pacific Northwest Region, Portland, OR.
- HENGST, G. E., AND J. O. DAWSON. 1994. Bark properties and fire resistance of selected tree species from the central hardwood region of North America. *Canadian Journal of Forest Research* 24:688–696.
- HERKERT, J. R. 1994a. Breeding bird communities of midwestern prairie fragments: The effects of prescribed burning and habitat-area. *Natural Areas Journal* 14: 128–135.
- HERKERT, J. R. 1994b. Status and habitat selection of the Henslow's Sparrow in Illinois. *Wilson Bulletin* 106: 35–45.
- HERKERT, J. R. 1994c. The effects of habitat fragmentation on midwestern grassland bird communities. *Ecological Applications* 4:461–471.
- HERKERT, J. R. 2001. Effects of management practices on grassland birds: Henslow's Sparrow. Northern Prairie Wildlife Research Center, Jamestown, ND. <<http://www.npwrc.usgs.gov/resource/literatr/grasbird/henslows/henslows.htm>> (17 February 2000).
- HERKERT, J. R., AND W. D. GLASS. 1999. Henslow's Sparrow response to prescribed fire in an Illinois prairie remnant. *Studies in Avian Biology* 19:160–164.
- HERKERT, J. R., P. D. VICKERY, AND D. E. KROODSMA. 2002. Henslow's Sparrow (*Ammodramus henslowii*). *In* A. Poole, and F. Gill (editors). *The Birds of North America*, No. 672. The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists' Union, Washington, DC.
- HERKERT, J. R., S. A. SIMPSON, R. L. WESTMEIER, T. L. ESKER, AND J. W. WALK. 1999. Response of Northern Harriers and Short-eared Owls to grassland management in Illinois. *Journal of Wildlife Management* 63: 517–523.
- HIGGINS, K. F. 1986. Interpretation and compendium of historical fire accounts in the northern Great Plains. USDI Fish and Wildlife Service Research Publication 161, Washington, DC.
- HIRTH, D. H., L. D. HARRIS, AND R. F. NOSS. 1991. Avian community dynamics in a peninsular Florida longleaf pine forest. *Florida Field Naturalist* 19:38–48.
- HITCHCOX, S. M. 1996. Abundance and nesting success of cavity-nesting birds in unlogged and salvage-logged burned forest in northwestern Montana. M. S. thesis, University of Montana, Missoula, MT.
- HOBSON, K. A., AND J. SCHIECK. 1999. Changes in bird communities in boreal mixed woodland forest: harvest and wildfire effects over 30 years. *Ecological Applications* 9:849–863.
- HOFFMAN, N. J. 1997. Distribution of *Picoides* woodpeckers in relation to habitat disturbance within the Yellowstone area. M.S. thesis, Montana State University, Bozeman, MT.
- HOLLAND, V. L. 1988. Coastal oak woodland. Pp. 78–79 *in* K. E. Mayer, and W. F. Laudenslayer (editors). *A guide to wildlife habitats of California*. California Department of Forestry, Sacramento, CA.
- HOLLING, C. S. (EDITOR). 1978. *Adaptive environmental assessment and management*. John Wiley, New York, NY.
- HOLTHUIZJEN, A. M. A., AND T. L. SHARIK. 1985. Colonization of abandoned pastures by eastern red cedar (*Juniperus virginiana* L.). *Canadian Journal of Forest Research* 15:1065–1068.
- HOLZMAN, B. A., AND B. H. ALLEN-DIAZ. 1991. Vegetation change in blue oak woodlands in California. Pp. 189–193 *in* R. B. Standiford (editor). *Proceedings of the symposium on oak woodlands and hardwood rangeland management*. USDA Forest Service General Technical Report PSW-126. USDA Forest Service, Pacific Southwest Research Station, Berkeley, CA.
- HOPKINS, W. E. 1979. Plant associations of South Chiloquin and Klamath Ranger Districts—Winema National Forest. USDA Forest Service Technical Report R6-ECOL-79-005. USDA Forest Service, Pacific Northwest Region, Portland, OR.
- HORTON, S. P. 1987. Effects of prescribed burning on breeding birds in a ponderosa pine forest, southeastern Arizona. M.S. thesis, University of Arizona, Tucson, AZ.
- HORTON, S. P., AND R. W. MANNAN. 1988. Effects of prescribed fire on snags and cavity-nesting birds in southeastern Arizona pine forests. *Wildlife Society Bulletin* 16:37–44.
- HOWE, H. F. 1994. Managing species diversity in tallgrass prairie: assumptions and implications. *Conservation Biology* 8:691–704.
- HOYT, J. S. 2000. Habitat associations of Black-backed and Three-toed woodpeckers in the northeastern boreal forest of Alberta. M.S. thesis, University of Alberta, Edmonton, AB, Canada.
- HOYT, J. S., AND HANNON S. J. 2002. Habitat associations of Black-backed and Three-toed woodpeckers in the boreal forest of Alberta. *Canadian Journal of Forest Research* 32:1881–1888.
- HUDDLE, J. A., AND S. G. PALLARDY. 1996. Effects of long-term annual and periodic burning on tree survival and growth in a Missouri Ozark oak-hickory forest. *Forest Ecology and Management* 82:1–9.
- HUFF, M. H. 1984. Post-fire succession in the Olympic Mountains, Washington: forest vegetation, fuels, and avifauna. Ph.D. dissertation, University of Washington, Seattle, WA.

- HUFF, M. H. 1995. Forest age structure and development following wildfires in the western Olympic Mountains, Washington. *Ecological Applications* 5:471–483.
- HUFF, M. H., AND J. K. AGE. 1991. Subalpine forest dynamics after fire in the Pacific Northwest national parks. National Park Service Cooperative Park Studies Unit, College of Forest Resource, University of Washington, Seattle, WA.
- HUFF, M. H., AND C. M. RALEY. 1991. Regional patterns of diurnal breeding bird communities in Oregon and Washington. Pp. 177–205 in L. F. Ruggiero, K. B. Aubry, A. B. Carey, and M. H. Huff (technical coordinators). *Wildlife and vegetation of unmanaged Douglas-fir forests*. USDA Forest Service General Technical Report PNW-GTR-285. USDA Forest Service, Pacific Northwest Research Station, Portland, OR.
- HUFF, M. H., J. K. AGE, AND D. A. MANUWAL. 1985. Postfire succession of avifauna in the Olympic Mountains, Washington. Pp. 8–15 in J. E. Lotan, and J. K. Brown (editors). *Fire's effects on wildlife habitat: Symposium proceedings*, USDA Forest Service General Technical Report INT-186. USDA Forest Service, Intermountain Research Station, Ogden, UT.
- HULBERT, L. C. 1988. The causes of fire effects in tallgrass prairie. *Ecology* 69:46–58.
- HULL, A. C. 1974. Species for seeding arid rangelands in southern Idaho. *Journal of Range Management* 27: 216–218.
- HUMPHREY, R. R. 1958. The desert grassland. *Botanical Review* 24:193–253.
- HUMPHREY, R. R. 1974. Fire in the deserts and desert grassland of North America. Pp. 365–400 in T. T. Kozlowski, and C. E. Ahlgren (editors). *Fire and ecosystems*. Academic Press, New York, NY.
- HUMPHREY, R. R. 1987. Ninety years and 535 miles. *Vegetation changes along the Mexican border*. University of New Mexico Press, Albuquerque, NM.
- HUNTER, M. L. JR. 1993. Natural fire regimes as spatial models for managing boreal forests. *Biological Conservation* 65:115–120.
- HUNTER, W. C., D. A. BUEHLER, R. A. CANTERBURY, J. L. CONFER, AND P. B. HAMEL. 2001. Conservation of disturbance-dependent birds in eastern North America. *Wildlife Society Bulletin* 29:440–455.
- HUTCHINSON, T. F., AND S. SUTHERLAND. 2000. Fire and understory vegetation: a large-scale study in Ohio and a search for general response patterns in central hardwood forests. Pp. 64–74 in D. Yaussy (editor). *Proceedings: workshop on fire, people, and the central hardwoods landscape*. USDA Forest Service General Technical Report NE-265. USDA Forest Service, Northeastern Research Station, Newton Square, PA.
- HUTTO, R. L. 1995. Composition of bird communities following stand-replacement fires in Northern Rocky Mountain (U.S.A.) conifer forests. *Conservation Biology* 9:1041–1058.
- HUTTO, R. L., AND J. S. YOUNG. 1999. Habitat relationships of landbirds in the Northern Region, USDA Forest Service. General Technical Report RMRS-GTR-32. USDA Forest Service, Rocky Mountain Research Station, Ogden, UT.
- IMBEAU, L., J.-P. L. SAVARD, AND R. GAGNON. 1999. Comparing bird assemblages in successional black spruce stands originating from fire and logging. *Canadian Journal of Zoology* 77:1850–1860.
- IMPORA, P. C. 1997. Spatial and temporal patterns of fire in the forests of the central Oregon Coast Range. Ph.D. dissertation, Oregon State University, Corvallis, OR.
- JACKSON, J. A. 1988. The southeastern pine forest ecosystem and its birds: past, present, and future. *Bird Conservation* 3:119–159.
- JACOBS, D., F., D. W. COLE, AND J. R. MCBRIDE. 1985. Fire history and perpetuation of natural coast redwood ecosystems. *Journal of Forestry* 83:494–497.
- JAMES, F. C., C. A. HESS, AND D. KUFRIN. 1997. Species-centered environmental analysis: indirect effects of fire history on Red-cockaded Woodpeckers. *Ecological Applications* 7:118–129.
- JANES, S. W. 1983. Status, distribution, and habitat selection of the grasshopper sparrow in Morrow County, Oregon. *Murrelet* 64:51–54.
- JEHLE, G., A. A. YACKEL ADAMS, J. A. SAVIDGE, AND S. K. SKAGEN. 2004. Nest survival estimation: a review of alternatives to the Mayfield estimator. *Condor* 106:472–484.
- JENSEN, M. E. 1990. Interpretation of environmental gradients which influence sagebrush community distribution in northeastern Nevada. *Journal of Range Management* 43:161–166.
- JEPSON, W. L. 1910. *Memoirs of the University of California Volume 2. The silva of California*. The University Press, Berkeley, CA.
- JEPSON, W. L. 1923. *The trees of California*. Second edition. Independent Pressroom and Williams Printing Company, San Francisco, CA.
- JOHANNESSEN, C. L., W. A. DAVENPORT, A. MILLET, AND S. MCWILLIAMS. 1971. The vegetation of the Willamette Valley. *Annals of the Association of American Geographers* 61:286–302.
- JOHNSON, A. S., AND P. E. HALE. 2002. The historical foundations of prescribed burning for wildlife: a southeastern perspective. Pp. 11–23 in W. Ford, K. R. Russell, and C. E. Moorman (editors). *The role of fire in nongame wildlife management and community restoration: traditional uses and new directions*. USDA Forest Service General Technical Report NE-288. USDA Forest Service, Northeastern Research Station, Newtown Square, PA.
- JOHNSON, D. E., H. A. M. MUKHTAR, R. MAPSON, AND R. R. HUMPHREY. 1962. The mortality of oak-juniper woodland species following wildfire. *Journal of Range Management* 15:201–205.
- JOHNSON, D. H., AND L. D. IGL. 2001. Area requirements of grassland birds: a regional perspective. *Auk* 118: 24–34.
- JOHNSON, D. H. AND T. A. O'NEIL (MANAGING DIRECTORS). 2001. *Wildlife-habitat relationships in Oregon and Washington*. Oregon State University Press, Corvallis, OR.

- JOHNSON, E. A. 1992. Fire and vegetation dynamics: studies from the North American boreal forest. Cambridge University Press, Cambridge, UK.
- JOHNSON, E. A., AND C. P. S. LARSEN. 1991. Climatically induced change in the fire frequency in the southern Canadian Rockies. *Ecology* 72:194–201.
- JOHNSON, E. A., K. MIYANISHI, AND S. R. J. BRIDGE. 2001. Wildfire regime in the boreal forest and the idea of suppression and fuel buildup. *Conservation Biology* 15:1554–1557.
- JOHNSON, E. A., K. MIYANISHI, AND J. M. H. WEIR. 1998. Wildfires in the western Canadian boreal forest: landscape patterns and ecosystem management. *Journal of Vegetation Science* 9:603–610.
- JOHNSON, P. S. 1993. Perspectives on the ecology and silviculture of oak-dominated forests in the central and eastern states. USDA Forest Service General Technical Report NC-153. USDA Forest Service, North Central Experiment Station, St. Paul, MN.
- JOHNSON, R. G., AND S. A. TEMPLE. 1986. Assessing habitat quality for birds nesting in fragmented tallgrass prairies. Pp. 245–249 in J. Verner, M. Morrison, and C. J. Ralph (editors). *Wildlife 2000: modeling habitat relationships of terrestrial vertebrates*. University of Wisconsin Press, Madison, WI.
- JOHNSON, R. G., AND S. A. TEMPLE. 1990. Nest predation and brood parasitism of tallgrass prairie birds. *Journal of Wildlife Management* 54:106–111.
- JOHNSON, R. R., AND D. E. JONES (TECHNICAL COORDINATORS). 1977. Importance, preservation, and management of riparian habitat: a symposium. USDA Forest Service General Technical Report RM-43. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- JOHNSON, T. H., AND R. H. WAUER. 1996. Avifaunal response to the 1977 La Mesa fire. Pp. 70–94 in C. D. Allen (technical editor). *Fire effects in southwestern forests: proceedings of the second La Mesa Fire symposium*. USDA Forest Service General Technical Report RM-GTR-286. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- JOHNSTON, V. R. 1970. The ecology of fire. *Audubon* 72: 76–119.
- JONES, D. D., L. M. CONNER, R. J. WARREN, AND G. O. WARE. 2002. The effect of supplemental prey and prescribed fire on success of artificial nests. *Journal of Wildlife Management* 66:1112–1117.
- KANTRUD, H. A., AND K. F. HIGGINS. 1992. Nest and nest site characteristics of some ground-nesting non-passerine birds of northern grasslands. *Prairie Naturalist* 24:67–84.
- KAUFFMANN, J. B. 1990. Ecological relationships of vegetation and fire in the Pacific Northwest. Pp. 39–52 in J. D. Walstad, S. R. Radosevich, and D. V. Sandberg (editors). *Natural and prescribed fire in Pacific Northwest forests*. Oregon State University Press, Corvallis, OR.
- KAUFMANN, M. R., C. M. REGAN, AND P. M. BROWN. 2000. Heterogeneity in ponderosa pine/Douglas-fir forests: Age and size structure in unlogged and logged landscapes of central Colorado. *Canadian Journal of Forest Research* 30:698–711.
- KAUTZ, R. S., D. T. GILBERT, AND G. M. MAULDIN. 1993. Vegetative cover in Florida based on 1985–1989 Landsat Thematic Mapper imagery. *Florida Scientist* 56:135–154.
- KAY, C. E. 1995. Aboriginal overkill and native burning: implications for modern ecosystem management. *Western Journal of Applied Forestry* 10:121–126.
- KEANE, R. E., K. C. RYAN, T. T. VEBLEN, C. D. ALLEN, J. LOGAN, AND B. HAWKES. 2002. Cascading effects of fire exclusion in Rocky Mountain ecosystems: a literature review. USDA Forest Service General Technical Report RMRS-GTR-91. USDA Forest Service, Rocky Mountain Research Station, Fort Collins, CO.
- KEELEY, J. E. 2000. Chaparral. Pp. 203–253 in M. G. Barbour, and W. D. Billings (editors). *North American terrestrial vegetation* (second edition). Cambridge University Press, Cambridge, UK.
- KERLEY, L. L., AND S. H. ANDERSON. 1995. Songbird responses to sagebrush removal in a high elevation sagebrush steppe ecosystem. *Prairie Naturalist* 27: 129–146
- KILGORE, B. M. 1971. Response of breeding bird populations to habitat changes in giant sequoia forest. *American Midland Naturalist* 85:135–152.
- KILGORE, B. M. 1981. Fire in ecosystem distribution and structure: western forests and scrublands. Pp. 58–89 in H. A. Mooney, T. M. Bonnicksen, N. L. Christensen, J. E. Lotan, and W. A. Reiners (technical coordinators). *Proceedings of the conference fire regimes and ecosystem properties*. USDA Forest Service General Technical Report WO-26. USDA Forest Service, Washington, DC.
- KING, D. I., AND B. E. BYERS. 2002. An evaluation of power line rights-of-way as habitat for early-successional shrubland birds. *Wildlife Society Bulletin* 30: 868–874.
- KING, D. I., R. M. DEGRAAF, AND C. R. GRIFFIN. 2001. Productivity of early successional birds in clearcuts and groupcuts in an eastern deciduous forest. *Journal of Wildlife Management* 65:345–350.
- KING, T. G., M. A. HOWELL, B. R. CHAPMAN, K. V. MILLER, AND R. A. SCHORR. 1998. Comparisons of wintering bird communities in mature pine stands managed by prescribed burning. *Wilson Bulletin* 110:570–574.
- KIPFMUELLER, K. F., AND W. L. BAKER. 2000. A fire history of a subalpine forest in south-eastern Wyoming, USA. *Journal of Biogeography* 27:71–85.
- KIRKMAN, L. K., M. B. DREW, AND D. EDWARDS. 1998. Effects of experimental fire regimes on the population dynamics of *Schwalbea americana* L. *Plant Ecology* 137:115–137.
- KIRKPATRICK, C., S. DESTEFANO, R. W. MANNAN, AND J. LLOYD. 2002. Trends in abundance of grassland birds following a spring prescribed burn in southern Arizona. *Southwestern Naturalist* 47:282–292.
- KIRKPATRICK, R. L., AND P. J. PEKINS. 2002. Nutritional

- value of acorns for wildlife. Pp. 173–181 in W. J. McShea, and W. M. Healy (editors). Oak forest ecosystems: ecology and management for wildlife. Johns Hopkins University Press, Baltimore, MD.
- KLEMMEDSON, J. O., AND J. G. SMITH. 1964. Cheatgrass (*Bromus tectorum* L.). Botanical Review 30:226–262.
- KLUTE, D. S., R. J. ROBEL, AND K. E. KEMP. 1997. Will conversion of Conservation Reserve Program (CRP) lands to pasture be detrimental for grassland birds in Kansas? American Midland Naturalist 137:206–212.
- KNAPP, A. K., AND T. R. SEASTEDT. 1998. Introduction: grasslands, Konza Prairie, and long-term ecological research. Pp. 3–15 in A. K. Knapp, J. M. Briggs, D. C. Hartnett, and S. L. Collins (editors). Grassland dynamics, Oxford University Press, New York, NY.
- KNICK, S. T. 1999. Forum: requiem for a sagebrush ecosystem? Northwest Science 73:47–51.
- KNICK, S. T., AND J. T. ROTENBERRY. 1995. Landscape characteristics of fragmented shrubsteppe landscapes and breeding passerine birds. Conservation Biology 9: 1059–1071.
- KNICK, S. T., AND J. T. ROTENBERRY. 1997. Landscape characteristics of disturbed shrubsteppe habitats in southwestern Idaho (U.S.A.). Landscape Ecology 12: 287–297.
- KNICK, S. T., AND J. T. ROTENBERRY. 1999. Spatial distribution of breeding passerine bird habitats in southwestern Idaho. Studies in Avian Biology 19:104–111.
- KNICK, S. T., AND J. T. ROTENBERRY. 2000. Ghosts of habitats past: contribution of landscape change to current habitats used by shrubland birds. Ecology 81:220–227.
- KNICK, S. T., AND J. T. ROTENBERRY. 2002. Effects of fragmentation on passerine birds breeding in intermountain shrubsteppe. Studies in Avian Biology 25:131–141.
- KNICK, S. T., D. S. DOBKIN, J. T. ROTENBERRY, M. A. SCHROEDER, W. M. VANDER HAEGEN, AND C. VAN RIPER III. 2003. Teetering on the edge or lost? Conservation and research issues for the avifauna of sagebrush habitats. Condor 105:611–634.
- KNOPF, F. L. 1988. Conservation of steppe birds in North America. Pp. 27–41 in P. D. Goriup (editor). Ecology and conservation of grassland birds. ICBP Technical Publication No. 7. International Council for Bird Preservation, Cambridge, UK.
- KNOPF, F. L. 1994. Avian assemblages on altered grasslands. Studies in Avian Biology 15:245–257.
- KOMAREK, E. V. 1965. Fire ecology—grasslands and man. Tall Timbers Fire Ecology Conference 4:169–220.
- KOMAREK, E. V. 1968. Lightning and lightning fires as ecological forces. Tall Timbers Fire Ecology Conference 8:169–197.
- KOMAREK, E. V. 1974. Effects of fire on temperate forests and related ecosystems: southeastern United States. Pp. 251–277 in T. T. Kozlowski, and C. E. Ahlgren (editors). Fire and ecosystems. Physiological ecology: a series of monographs, texts, and treatises Academic Press, New York, NY.
- KOMAREK, E. V. 1981. History of prescribed fire and controlled burning in wildlife management in the South. Pp. 1–14 in G. W. Wood (editor). Prescribed fire and wildlife in southern forests. Belle W. Baruch Forest Science Institute, Clemson University, Georgetown, SC.
- KOTLIAR, N. B., S. J. HEIL, R. L. HUTTO, V. A. SAAB, C. P. MELLCHER, AND M. E. MCFADZEN. 2002. Effects of fire and post-fire salvage logging on avian communities in conifer-dominated forests of the western United States. Studies in Avian Biology 25:49–64.
- KRAMMES, J. S. (TECHNICAL COORDINATOR). 1990. Effects of fire management of southwestern natural resources. USDA Forest Service General Technical Report RM-191. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO
- KREISEL, K. J., AND S. J. STEIN. 1999. Bird use of burned and unburned coniferous forests during winter. Wilson Bulletin 111:243–250.
- KROHN, W. B., R. B. BOONE, S. A. SADER, J. A. HEPINSTALL, S. M. SCHAEFER, AND S. L. PAINTON. 1998. Maine GAP analysis: a geographic analysis of biodiversity. Maine Cooperative Fish and Wildlife Research Unit, Maine Image Analysis Laboratory, U.S. Geological Survey, University of Maine, Orono, ME.
- KRUGER, E. L., AND P. B. REICH. 1997. Responses of hardwood regeneration to fire in mesic forest openings. I. Post-fire community dynamics. Canadian Journal of Forest Research 27:1822–1831.
- KÜCHLER, A. W. 1964. Potential natural vegetation of the conterminous United States 15(1:3,168,000 scale map). American Geographical Society, Special Publication Number 36. New York, NY.
- KÜCHLER, A. W. 1976. Potential natural vegetation of California (map 1:1,000,000). American Geographical Society, New York, NY.
- KUDDER-FISCHER, L. M., AND M. A. ARTHUR. 2002. Response of understory vegetation and tree regeneration to a single prescribed fire in oak-pine forests. Natural Areas Journal 22:43–52.
- KUSH, J., R. MELDAHL, AND W. BOYER. 1999. Understory plant community response after 23 years of hardwood control treatments in natural longleaf pine (*Pinus palustris*) forests. Canadian Journal of Forest Research 29:1047–1054.
- LALANDE, J., AND R. PULLEN. 1999. Burning for a “Fine and beautiful open country:” native uses of fire in southwestern Oregon. Pp. 255–276 in R. Boyd (editor). Indians, fire, and the land in the Pacific Northwest. Oregon State University Press, Corvallis, OR.
- LANDERS, J. L., AND W. D. BOYER. 1999. An old-growth definition for upland longleaf and south Florida slash pine forests, woodlands, and savannas. USDA Forest Service General Technical Report No.SRS-29. USDA Forest Service, Southern Research Station, Asheville, NC.
- LANDRES, P., P. MORGAN, AND F. J. SWANSON. 1999. Overview of the use of natural variability concepts in managing ecological systems. Ecological Applications 9:1179–1188.
- LANHAM, J. D., P. D. KEYSER, P. H. BROSE, AND D. H. VAN LEAR. 2002. Oak regeneration using the shelterwood-burn technique: management options and implications

- for songbird conservation in the southeastern United States. *Forest Ecology and Management* 155: 143–152.
- LARSEN, C. P. S. 1997. Spatial and temporal variations in boreal forest fire frequency in northern Alberta. *Journal of Biogeography* 24:663–673.
- LATHROP, E. W., AND C. D. OSBORNE. 1991. Influence of fire on oak seedlings and saplings in southern oak woodland on the Santa Rosa Plateau Preserve, Riverside County, California. Pp. 366–370 in R. B. Standiford (editor). Proceedings of the symposium on oak woodlands and hardwood rangeland management. USDA Forest Service General Technical Report PSW-126. USDA Forest Service, Pacific Southwest Research Station, Berkeley, CA.
- LAUNCHBAUGH, J. L., AND C. E. OWENSBY. 1978. Kansas rangelands: their management based on a half century of research. Kansas Agriculture Experiment Station. Bulletin 662. Manhattan, KS.
- LAVERTY, L., AND J. WILLIAMS. 2000. Protecting people and sustaining resources in fire-adapted ecosystems—a cohesive strategy. USDA Forest Service Report, Dillon, MT.
- LAWRENCE, G. E. 1966. Ecology of vertebrate animals in relation to chaparral fire in the Sierra Nevada foothills. *Ecology* 47:278–291.
- LAYCOCK, W. A. 1991. Stable states and thresholds of range condition on North American rangelands: a viewpoint. *Journal of Range Management* 44:427–433.
- LEE, P. 2002. Forest structure after wildfire and harvesting. Chapter 5 in S. J. Song (editor). Ecological basis for stand management: a synthesis of ecological responses to wildfire and harvesting. Alberta Research Council Inc., Vegreville, AB, Canada.
- LEE, P., AND S. BRADBURY. 2002. Coarse filter management paradigms for timber harvest. Pp. 2–12 in S. J. Song (editor). Ecological basis for stand management: a synthesis of ecological responses to wildfire and harvesting. Alberta Research Council Inc., Vegreville, AB, Canada.
- LEETE, B. E. 1938. Forest fires in Ohio 1923 to 1935. Ohio Agricultural Experiment Station, Bulletin 598, Wooster, OH.
- LEFORT, P. 2003. Ecology and management of claybelt forests: a knowledge synthesis. Natural Resources Canada, Lake Abitibi Model Forest. Cochrane, ON, Canada.
- LEGLER, J. M. 1960. Natural history of the ornate box turtle, *Terrapene ornata ornata* Agassiz. University of Kansas Museum of Natural History. Publication 11:527–669.
- LEONARD, D. L. JR. 2001. Three-toed Woodpecker (*Picoides tridactylus*). In A. Poole, and F. Gill (editors). The Birds of North America, No. 588. The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists' Union, Washington, DC.
- LEOPOLD, A. 1924. Grass, brush, and timber fire in southern Arizona. *Journal of Forestry* 22:1–10.
- LERTZMAN, K. P., AND C. J. KREBS. 1991. Gap-phase structure of a subalpine old-growth forest. *Canadian Journal of Forest Research* 12:1730–1741.
- LEWIS, H. T. 1973. Patterns of Indian burning in California: ecology and ethno-history. Anthropological Paper No. 1. Ballena Press, Ramona, CA.
- LEWIS, H. T. 1977. Maskuta: The ecology of Indian fires in northern Alberta. *Western Canadian Journal of Anthropology* 7:15–52.
- LEWIS, H. T. 1980. Hunters-gatherers and problems for fire history. Pp. 115–119 in M. A. Stokes, and J. H. Dieterich (editors). Fire history workshop. USDA Forest Service General Technical Report RM-81. USDA Forest Service, Rocky Mountain Research Station, Fort Collins, CO.
- LEWIS, H. T. 1982. Fire technology and resource management in Aboriginal North America and Australia. Pp. 45–67 in N. M. Williams, and E. S. Hunn (editors). Resource managers: North American and Australian hunter-gatherers. AAAS Selected Symposium No. 67. Westview Press, Boulder, CO.
- LEWIS, H. T. 1985. Why Indians burned: specific versus general reasons. Pp. 75–80 in J. E. Loton, B. Kilgore, W. Fischer, and R. Mutch (editors). Symposium and workshop on wilderness fire. USDA Forest Service General Technical Report INT-182. USDA Forest Service, Intermountain Forest and Range Experiment Station, Missoula, MT.
- LEWIS, H. T. 1989. Ecological and technological knowledge of fire: aborigines versus park rangers in northern Australia. *American Anthropologist* 91:940–961.
- LEWIS, H. T. 1993. Patterns of Indian burning in California: ecology and ethno-history. Pp. 55–116 in T. C. Blackburn, and K. Anderson (editors), Before the wilderness: environmental management by Native Californians. Ballena Press, Menlo Park, CA.
- LI, P., AND T. E. MARTIN. 1991. Nest-site selection and nesting success of cavity-nesting birds in high elevation forest drainages. *Auk* 108:405–418.
- LILLYBRIDGE, T. R., B. L. KOVALCHIK, C. K. WILLIAMS, AND B. G. SMITH. 1995. Field guide for forested plant association of the Wenatchee National Forest. USDA Forest Service General Technical Report PNW-GTR-359. USDA Forest Service, Pacific Northwest Research Station, Portland, OR.
- LINDER, K. A., AND S. H. ANDERSON. 1998. Nesting habitat of Lewis' Woodpeckers in southeastern Wyoming. *Journal of Field Ornithology* 69:109–116.
- LITVAITIS, J. A., D. L. WAGNER, J. L. CONFER, M. D. TARR, E. J. SNYDER. 1999. Early-successional forests and shrub-dominated habitats: land-use artifact or critical community in the northeastern United States. *Northeast Wildlife* 54:101–117.
- LLOYD, J., R. W. MANNAN, S. DESTEFANO, AND C. KIRKPATRICK. 1998. The effects of mesquite invasion on a southeastern Arizona grassland bird community. *Wilson Bulletin* 110:403–408.
- LORIMER, C. G. 1985. The role of fire in the perpetuation of oak forests. Pp. 8–25 in J. E. Johnson (editor). Proceedings: challenges in oak management and

- utilization. Cooperative Extension Service, University of Wisconsin, Madison, WI.
- LORIMER, C. G. 1993. Causes of the oak regeneration problem. Pp. 14–39 in D. L. Loftis, and C. E. McGee (editors). Oak regeneration: serious problems, practical recommendations. USDA Forest Service General Technical Report SE-84. USDA Forest Service, Southeastern Forest Experiment Station, Asheville, NC.
- LORIMER, C. G., J. W. CHAPMAN, AND W. D. LAMBERT. 1994. Tall understory vegetation as a factor in the poor development of oak seedlings beneath mature stands. *Journal of Ecology* 82:227–237.
- LOTAN, J. E., AND J. K. BROWN (COMPILERS). 1985. Fire's effects on wildlife habitat—symposium proceedings. USDA Forest Service General Technical Report INT-186. USDA Forest Service, Intermountain Research Station, Ogden, UT.
- LOWE, C. H., AND D. E. BROWN. 1982. Introduction to biotic communities of the American Southwest. *Desert Plants* 4:8–16.
- LOWE, P. O., P. F. FOLLIOTT, J. H. DEITERICH, AND D. R. PATTON. 1978. Determining potential benefits from wildlife in Arizona ponderosa pine forests. USDA Forest Service Gen. Tech. Rep. RM-GTR-52. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- MACK, R. N. 1981. Invasion of *Bromus tectorum* L. into western North America: an ecological chronicle. *Agro-Ecosystems* 7:145–165.
- MACLEAN, D. A. 1980. Vulnerability of fir–spruce stands during uncontrolled spruce budworm outbreaks: a review and discussion. *Forestry Chronicle* 56:213–221.
- MACMAHON, J. A. 2000. Warm deserts. Pp. 285–322 in M. G. Barbour, and W. D. Billings (editors). *North American terrestrial vegetation* (second edition). Cambridge University Press, Cambridge, UK.
- MAINE FOREST SERVICE. 2001. The 2001 biennial report on the state forest and progress report on forest sustainability standards. <<http://www.maine.gov/doc/mfs/pubs/pdf/2001sof.pdf>> (12 June 2004).
- MANUWAL, D. A. 1991. Spring bird communities in the southern Washington Cascade Range. Pp. 161–174 in L. F. Ruggiero, K. B. Aubry, A. B. Carey, and M. H. Huff (technical coordinators). *Wildlife and vegetation of unmanaged Douglas-fir forests*. USDA Forest Service General Technical Report PNW-GTR-285. USDA Forest Service, Pacific Northwest Research Station, Portland, OR.
- MANUWAL, D. A., AND M. H. HUFF. 1987. Spring and winter bird populations in a Douglas-fir forest sere. *Journal of Wildlife Management* 51:586–595.
- MANUWAL, D. A., M. H. HUFF, M. R. BAUER, C. B. CHAPPELL, AND K. HEGSTAD. 1987. Summer birds of the upper subalpine zone of Mount Adams, Mount Rainier, and Mount Saint Helens, Washington. *Northwest Science* 61:82–92.
- MARCOT, B. G. 1984. Habitat relationships of birds and young-growth Douglas-fir in northwestern California. Ph.D. dissertation, Oregon State University, Corvallis, OR.
- MARKS, J. S., D. L. EVANS, AND D. W. HOLT. 1994. Long-eared Owl (*Asio otus*). In A. Poole, and F. Gill (editors), *The Birds of North America*, No. 133. The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists' Union, Washington, DC.
- MARSHALL, J. T., JR. 1963. Fire and birds in the mountains of southern Arizona. *Proceedings of the Tall Timbers Fire Ecology Conference* 2:134–141.
- MARTIN, A. C., H. S. ZIM, AND A. L. NELSON. 1961. *American wildlife and plants: a guide to wildlife food habits*. Dover Publishers, New York.
- MARTIN, K., AND J. M. EADIE. 1999. Nest webs: a community-wide approach to the management and conservation of cavity-nesting forest birds. *Forest Ecology and Management* 115:243–257.
- MARTIN, K. K., E. H. AITKEN, AND K. L. WIEBE. 2004. Nest sites and nest webs for cavity-nesting communities in interior British Columbia: nest characteristics and niche partitioning. *Condor* 106:5–19.
- MARTIN, R. C. 1990. Sage Grouse responses to wildfire in spring and summer habitats. M.S. thesis, University of Idaho, Moscow, ID.
- MARTIN, R. E., AND D. B. SAPSIS. 1992. Fires as agents of biodiversity: pyrodiversity promotes biodiversity. Pp. 150–157 in *Proceedings of the symposium on biodiversity of northwestern California*. University of California, Berkeley, Extension, Department of Forestry and Range Management, Berkeley, CA.
- MARTIN, T. E., AND P. LI. 1992. Life history traits of open-vs. cavity-nesting birds. *Ecology* 73:579–592.
- MAST, J. N., P. Z. FULÉ, M. M. MOORE, W. W. COVINGTON, AND A. WALTZ. 1999. Restoration of presettlement age structure of an Arizona ponderosa pine forest. *Ecological Applications* 9:228–239.
- MASTERS, R. E., C. W. WILSON, D. S. CRAM, AND G. A. BUKENHOFER. 2002. Influence of ecosystem restoration for Red-cockaded Woodpeckers on breeding bird and small mammal communities. Pp. 73–90 in W. Ford, K. R. Russell, and C. E. Moorman (editors). *The role of fire in nongame wildlife management and community restoration: traditional uses and new directions*. USDA Forest Service General Technical Report NE-288. USDA Forest Service, Northeastern Research Station, Newtown Square, PA.
- MAURER, B. A. 1986. Predicting habitat quality for grassland birds using density-habitat correlations. *Journal of Wildlife Management* 50:556–566.
- MAY, C. A., AND R. J. GUTIERREZ. 2002. Habitat associations of Mexican Spotted Owl nest and roost sites in central Arizona. *Wilson Bulletin* 114:457–466.
- MAYFIELD, H. 1961. Nesting success calculated from exposure. *Wilson Bulletin* 73:255–261.
- MCCARTHUR, E. D. 1994. Ecology, distribution, and values of sagebrush within the Intermountain region. Pp. 347–351 in S. B. Monsen, and S. G. Kitchen (compilers). *Proceedings—ecology and management of annual rangelands*. USDA Forest Service General Technical

- MILLER, R. F., AND R. J. TAUSCH. 2001. The role of fire in pinyon and juniper woodlands: a descriptive analysis. Tall Timbers Research Station Miscellaneous Publication No. 11:15–30.
- MILLER, R. F., AND P. F. WIGAND. 1994. Holocene changes in semiarid pinyon-juniper woodlands. *BioScience* 44: 465–474.
- MILLER, R. F., T. SVEJCAR, AND J. A. ROSE. 2000. Impacts of western juniper on plant community composition and structure. *Journal of Range Management* 53:574–585.
- MILLS, G. S., J. B. DUNNING, AND J. M. BATES. 1991. The relationship between breeding bird density and vegetation volume. *Wilson Bulletin* 103:468–479.
- MINNICH, R. A., M. G. BARBOUR, J. H. BURK, AND R. F. FERNAU. 1995. Sixty years of change in California conifer forests of the San Bernardino Mountains. *Conservation Biology* 9:902–914.
- MITCHELL, M. S., R. A. LANCIA, AND J. A. GERWIN. 2001. Using landscape-level data to predict the distribution of birds on managed forest: effects of scale. *Ecological Applications* 11:1692–1708.
- MITCHELL, R. B., R. A. MASTERS, S. S. WALLER, K. J. MOORE, AND L. J. YOUNG. 1996. Tallgrass prairie vegetation response to spring burning dates, fertilizer, and atrazine. *Journal of Range Management* 49:131–136.
- MITTON, J. B., AND K. B. STURGEON (EDITORS). 1982. Bark beetles in North American conifers: a system for the study of evolutionary biology. University of Texas Press, Austin, TX.
- MOIR, W. H. 1993. Alpine tundra and coniferous forest. Pp. 47–84 in W. A. Dick-Peddie (editor). *New Mexico vegetation: past, present, and future*. University of New Mexico Press, Albuquerque, NM.
- MOIR, W. H., AND J. O. CARLETON. 1986. Classification of pinyon-juniper sites on National Forests in the Southwest. Pp. 216–226 in R. L. Everett (compiler). *Proceedings—pinyon-juniper conference*. USDA Forest Service General Technical Report INT-GTR-215. USDA Forest Service, Intermountain Research Station, Ogden, UT.
- MOIR, W. H., AND J. H. DIETERICH. 1988. Old-growth ponderosa pine from succession on pine-bunchgrass habitat types in Arizona and New Mexico. *Natural Areas Journal* 8:17–24.
- MOIR, W. H., B. W. GEILS, M. A. BENOIT, AND D. SCURLOCK. 1997. Ecology of Southwestern ponderosa pine forests. Pp. 3–27 in W. M. Block, and D. M. Finch (technical editors). *Songbird ecology in southwestern ponderosa pine forests: a literature review*. USDA Forest Service General Technical Report RM-GTR-292. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- MONSEN, S. B. 1994. The competitive influences of cheatgrass (*Bromus tectorum*) on site restoration. Pp. 43–50 in S. B. Monsen, and S. G. Kitchen (compilers). *Proceedings—ecology and management of annual rangelands*. USDA Forest Service General Technical Report INT-GTR-313. USDA Forest Service, Intermountain Research Station, Ogden, UT.
- MOORE, C. T. 1972. Man and fire in the central North American grasslands 1535–1890: A documentary historical geography. Ph.D. dissertation, University of California, Los Angeles.
- MOORE, M. M., W. W. COVINGTON, AND P. Z. FULÉ. 1999. Reference conditions and ecological restoration: a southwestern ponderosa pine perspective. *Ecological Applications* 9:1266–1277.
- MORGAN, P., C. C. HARDY, T. W. SWETNAM, M. G. ROLLINS, AND D. G. LONG. 2001. Mapping fire regimes across time and space: understanding coarse and fine-scale fire patterns. *International Journal of Wildland Fire* 10:329–342.
- MORIARTY, D. J., R. E. FARRIS, D. K. NODA, AND P. A. STANTON. 1985. Effects of fire on a coastal sage scrub bird community. *Southwestern Naturalist* 30:452–453.
- MORISSETTE, J. L., T. P. COBB, R. M. BRIGHAM, AND P. C. JAMES. 2002. The response of boreal forest songbird communities to fire and post-fire harvesting. *Canadian Journal of Forest Research* 32:2169–2183.
- MORRISON, M. L., AND E. C. MESLOW. 1983a. Bird community structure on early-growth clearcuts in western Oregon (USA). *American Midland Naturalist* 110: 129–137.
- MORRISON, M. L., AND E. C. MESLOW. 1983b. Bird community structure on early-growth clearcuts in the Oregon Coast ranges. USDA Forest Service Research Paper PNW-RP-305. USDA Forest Service, Pacific Northwest Forest and Range Experiment Station, Portland, OR.
- MORRISON, P., AND F. J. SWANSON. 1990. Fire history in two forest ecosystems of the central western Cascades of Oregon. USDA Forest Service General Technical Report PNW-GTR-254. USDA Forest Service, Pacific Northwest Research Station, Portland, OR.
- MOTZKIN, G., D. FOSTER, A. ALLEN, J. HARROD, AND R. BOONE. 1996. Controlling site to evaluate history: vegetation patterns of a New England sand plain. *Ecological Monographs* 66:345–365.
- MURPHY, E. C., AND W. A. LEHNHASEN. 1998. Density and foraging ecology of woodpeckers following a stand-replacement fire. *Journal of Wildlife Management* 62: 1359–1372.
- MURPHY, P. J. 1985. Methods for evaluating the effects of fire management in Alberta. Ph.D. dissertation, University of British Columbia, Vancouver, BC, Canada.
- MYERS, R. L. 1990. Scrub and high pine. Pp. 150–193 in R. L. Myers, and J. J. Ewel (editors). *Ecosystems of Florida*. University of Central Florida Press, Orlando, FL.
- MYERS, R. L., AND J. J. EWEL. 1990. Problems, prospects, and strategies for conservation. Pp. 574–632 in R. L. Myers, and J. J. Ewel (editors). *Ecosystems of Florida*. University of Central Florida Press, Orlando, FL.
- NABHAN, G. P. 2000. Biodiversity: the variety of life that sustains our own. Pp. 119–126 in S. J. Phillips, and P. W. Cornus (editors). *A natural history of the Sonoran desert*. University of California Press, Berkeley, CA.

- NAPPI, A. 2000. Distribution des pics et utilisation des arbres morts dans les forêts soumises aux perturbations naturelles en pessière à mousses. M.S. thesis, Université du Québec à Montréal, Montréal, Québec, Canada.
- NAPPI, A., P. DRAPEAU, J.-F. GIROUX, AND J.-P. SAVARD. 2003. Snag use by foraging Black-backed Woodpeckers in a recently burned eastern boreal forest. *Auk* 120: 505–511.
- NATIONAL AGRICULTURAL STATISTICS SERVICE. 2002. Annual bulletin: New England agricultural statistics, 2002. <<http://www.nass.usda.gov/nh/>> (12 June 2004).
- NATIONAL INTERAGENCY FIRE CENTER. 2003. Wildland fire statistics. <<http://www.nifc.gov/stats/>> (2 September 2003).
- NELLE, P. J., K. P. REESE, AND J. W. CONNELLY. 2000. Long-term effects of fire on Sage-Grouse habitat. *Journal of Range Management* 53:586–591.
- NIEMI, G. J. 1978. Breeding birds of burned and unburned areas in northern Minnesota. *Loon* 50:73–84.
- NORTON, M. R., AND S. J. HANNON. 1997. Songbird response to partial-cut logging in the boreal mixed-wood forest of Alberta. *Canadian Journal of Forest Research* 27: 44–53.
- NOSON, A. C. 2002. Avian communities in relation to habitat influenced by fire in a sagebrush steppe landscape. M.S. thesis, Oregon State University, Corvallis, OR.
- NOSS, R. F. 1989. Longleaf pine and wiregrass: keystone components of an endangered ecosystem. *Natural Areas Journal* 9:211–213.
- NOSS, R. F. (EDITOR). 1999. The redwood forest: history, ecology, and conservation of the coast redwoods. Island Press, Washington, DC.
- NOSS, R. F., E. T. LAROE, III, AND J. M. SCOTT. 1995. Endangered ecosystems of the United States: a preliminary assessment of loss and degradation. National Biological Service Biological Report 28, Washington, DC.
- NUZZO, V. 1986. Extent and status of midwest oak savanna: presettlement and 1985. *Natural Areas Journal* 6: 6–36.
- O'MEARA, T. E., J. B. HAUFLE, L. H. LAVERN, AND J. G. NAGY. 1981. Nongame wildlife responses to chaining of pinyon-juniper woodlands. *Journal of Wildlife Management* 45:381–389.
- OGG, A. G., JR. 1994. A review of the chemical control of downy brome. Pp. 194–196 in S. B. Monsen, and S. G. Kitchen (compilers). Proceedings—ecology and management of annual rangelands. USDA Forest Service General Technical Report INT-GTR-313, USDA Forest Service, Intermountain Research Station, Ogden, UT.
- OLSON, D. L. 2000. Fire in riparian zones: a comparison of historical fire occurrence in riparian and upland forests in the Blue Mountains and southern Cascades of Oregon. M.S. thesis, University of Washington, Seattle, WA.
- OREGON-WASHINGTON PARTNERS IN FLIGHT. 2000. Landbird conservation plans. <http://community.gorge.net/natres/pif/cons_page1_.html> (23 July 2004).
- ÖSTLUND, L., O. ZACKRISSON, AND A.-L. AXELSSON. 1997. The history and transformation of a Scandinavian boreal forest landscape since the 19th century. *Canadian Journal of Forest Research* 27:1198–1206.
- OVERBY, S. T., AND H. M. PERRY. 1996. Direct effects of prescribed fire on available nitrogen and phosphorus in an Arizona chaparral watershed. *Arid Soil Research and Rehabilitation* 10:347–357.
- OVERTURF, J. H. 1979. The effects of forest fire on breeding bird populations of ponderosa pine forests of northern Arizona. M.S. thesis, Northern Arizona University, Flagstaff, AZ.
- PACKEE, E. C., C. D. OLIVER, AND P. D. CRAWFORD. 1983. Ecology of Pacific silver fir. Pp. 19–34 in C. D. Oliver, and R. M. Kenady (editors). Proceedings of the biology and management of true fir in the Pacific Northwest symposium. University of Washington, Seattle, WA.
- PADEN, I. D., AND M. E. SCHLICHTMANN. 1959. The Big Oak Flat Road: an account of freighting from Stockton to Yosemite. Yosemite Natural History Association, Yosemite National Park, CA.
- PAIGE, C., AND S. A. RITTER. 1999. Birds in a sagebrush sea: managing sagebrush habitats for bird communities. Partners in Flight Western Working Group, Boise, ID.
- PALMER, W. E., K. D. GODWIN, G. A. HURST, AND D. A. MILLER. 1996. Effects of prescribed burning on Wild Turkeys. Transactions of the North American Wildlife and Natural Resources Conference 61:228–236.
- PAMPUSH, G. J., AND R. G. ANTHONY. 1993. Nest success, habitat utilization and nest-site selection of Long-billed Curlews in the Columbia Basin, Oregon. *Condor* 95:957–967.
- PARSHALL, T., AND D. R. FOSTER. 2002. Fire on the New England landscape: regional and temporal variation, cultural and environmental controls. *Journal of Biogeography* 29:1305–1317.
- PARTNERS IN FLIGHT. 2004. Partners in Flight—U.S. <<http://www.partnersinflight.org>> (2 February 2004).
- PASE, C. P., AND D. E. BROWN. 1982. Interior chaparral. *Desert Plants* 4:95–99.
- PASE, C. P., AND C. E. GRANFELT (TECHNICAL COORDINATORS). 1977. The use of fire on Arizona rangelands. Arizona Interagency Range Commission, Publication No. 4. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- PATTEN, D. T. 1998. Riparian ecosystems of semi-arid North America: diversity and human impacts. *Wetlands* 18:498–512.
- PATTERSON, M. P., AND L. B. BEST. 1996. Bird abundance and nesting success in Iowa CRP fields: the importance of vegetation structure and composition. *American Midland Naturalist* 135:153–167.
- PATTERSON, W. A., III, AND K. E. SASSAMAN. 1988. Indian fires in the prehistory of New England. Pp. 107–135 in G. P. Nichols (editor). *Holocene human ecology in northeastern North America*. Plenum, New York, NY.
- PAYSEN, T. E., R. J. ANSELY, J. K. BROWN, G. J. GOTTFRIED, S. M. HAASE, M. G. HARRINGTON, M. G. NAROG, S.

- S. SACKETT, AND R. C. WILSON. 2000. Fire in western shrubland, woodland, and grassland ecosystems. Pp. 121–159 in J. K. Brown, and J. K. Smith (editors). Wildland fire in ecosystems: effects of fire on flora. USDA Forest Service General Technical Report RMRS-GTR-42-Volume 2. USDA Forest Service, Rocky Mountain Research Station, Ogden, UT.
- PEET, R. K. 1981. Forest vegetation of the Colorado Front Range. *Vegetatio* 45:3–75.
- PEET, R. K. 2000. Forests of the Rocky Mountains. Pp. 76–121 in M. G. Barbour, and W. D. Billings (editors). North American terrestrial vegetation. Cambridge University Press, Cambridge, UK.
- PEET, R. K., AND D. J. ALLARD. 1995. Longleaf pine vegetation of the southern Atlantic and eastern Gulf Coast regions: a preliminary classification. Tall Timbers Fire Ecology Conference 18:45–81.
- PENDERGRASS, K. L., P. M. MILLER, J. B. KAUFFMAN, AND T. N. KAYE. 1999. The role of prescribed burning in the maintenance of an endangered plant species, *Lomatium bradshawii*. *Ecological Applications* 9:1420–1429.
- PERKINS, D. W., P. D. VICKERY, T. F. DEAN, AND M. D. SCHEUERELL. 1998. Florida Grasshopper Sparrow reproductive success based on nesting records. *Florida Field Naturalist* 26:7–17.
- PETERJOHN, B. G., AND J. R. SAUER. 1999. Population status of North American grassland birds. *Studies in Avian Biology* 19:27–44.
- PETERJOHN, B. G., J. R. SAUER, AND W. A. LINK. 1994. The 1992 and 1993 summary of the North American Breeding Bird Survey. *Bird Populations* 2:46–61.
- PETERS, E. F., AND S. C. BUNTING. 1994. Fire conditions pre- and post-occurrence of annual grasses on the Snake River Plain. Pp. 31–36 in S. B. Monsen, and S. G. Kitchen (compilers). Proceedings—ecology and management of annual grasslands. USDA Forest Service General Technical Report INT-GTR-313. USDA Forest Service, Intermountain Research Station, Ogden, UT.
- PETERSEN, K. L., AND L. B. BEST. 1987. Effects of prescribed burning on nongame birds in a sagebrush community. *Wildlife Society Bulletin* 15:317–329.
- PETERSEN, K. L., AND L. B. BEST. 1999. Design and duration of perturbation experiments: implications for data interpretation. *Studies in Avian Biology* 19:230–236.
- PETERSON, D. W., AND P. B. REICH. 2001. Prescribed fire in oak savanna: fire frequency effects on stand structure and dynamics. *Ecological Applications* 11:914–927.
- PETERSON, E. B., AND N. M. PETERSON. 1992. Ecology, management and use of aspen and balsam poplar in the prairie provinces. Special Report 1, Forestry Canada, Northern Forestry Centre, Edmonton, AB, Canada.
- PISTER, A. R. 1980. Postfire avian ecology in Yellowstone National Park. M.S. thesis, Washington State University, Pullman, WA.
- PICKETT, S. T. A., AND P. S. WHITE. 1985. The ecology of natural disturbance and patch dynamics. Academic Press, New York, NY.
- PIDGEON, A. M., N. E. MATHEWS, R. BENOIT, AND E. V. NORDHEIM. 2001. Response of avian communities to historic habitat change in the northern Chihuahuan desert. *Conservation Biology* 6:1772–1788.
- PIEMEISEL, R. L. 1951. Causes effecting change and rate of change in a vegetation of annuals in Idaho. *Ecology* 32:53–72.
- PIEPER, R. D., AND R. D. WITTIE. 1990. Fire effects in southwestern chaparral and pinyon-juniper vegetation. Pp. 87–93 in J. S. Krammes (technical coordinator). Effects of fire management of southwestern natural resources. USDA Forest Service General Technical Report RM-191. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- PITCHER, D. C. 1987. Fire history and age structure in red fir forests of Sequoia National Park, California. *Canadian Journal of Forest Research* 17:582–587.
- PLATT, W. J. 1999. Southeastern pine savannas. Pp. 23–51 in R. C. Anderson, J. S. Fralish, and J. M. Baskin (editors). Savannas, barrens, and rock outcrop plant communities of North America. Cambridge University Press, Cambridge, UK.
- PLENTOVICH, S., N. R. HOLLER, AND G. E. HILL. 1999. Habitat requirements of Henslow's Sparrows wintering in silvicultural lands of the Gulf coastal plain. *Auk* 116:109–115.
- PLUMB, T. R. 1980. Response of oaks to fire. Pp. 202–215 in T. R. Plumb (technical coordinator). Proceedings of the symposium on the ecology, management, and utilization of California oaks. USDA Forest Service General Technical Report PSW-44. USDA Forest Service, Pacific Southwest Research Station, Berkeley, CA.
- PLUMB, T. R., AND A. P. GOMEZ. 1983. Five southern oaks: identification and postfire management. USDA Forest Service General Technical Report PSW-71. USDA Forest Service, Pacific Southwest Forest and Range Experiment Station, Berkeley, CA.
- POOLE, A., AND F. GILL. (EDITORS). 2004. The Birds of North America. The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists' Union, Washington, DC.
- POWELL, B. F., AND R. J. STEIDL. 2000. Nesting habitat and reproductive success of southwestern riparian birds. *Condor* 102:823–831.
- POWELL, H. 2000. The influence of prey density on post-fire habitat use of the Black-backed Woodpecker. M.S. thesis, University of Montana, Missoula, MT.
- POWERS, R. F. 1990. Digger pine. Pp. 463–469 in R. M. Burns, and B. H. Honkala (technical coordinators). *Silvics of North America. Volume 1, conifers*. USDA Forest Service Agriculture Handbook 654, Washington, DC.
- PROVENCER, L., N. M. GOBRIS, L. A. BRENNAN. 2002a. Effects of hardwood reduction on winter birds in northwest Florida longleaf pine sandhill forests. *Auk* 119:71–87.
- PROVENCER, L., N. M. GOBRIS, L. A. BRENNAN, D. R. GORDON, AND J. L. HARDESTY. 2002b. Breeding bird response to midstory hardwood reduction in Florida sandhill longleaf pine forests. *Journal of Wildlife Management* 66:641–661.

- PRUITT, L. 1996. Henslow's Sparrow status assessment. USDI Fish and Wildlife Service, Bloomington Field Office, Bloomington, IN.
- PULLIAM, H. R., AND J. B. DUNNING. 1987. The influence of food supply on local density and diversity of sparrows. *Ecology* 68:1009–1014.
- PURCELL, K. L., AND J. VERNER. 1998. Density and reproductive success of California Towhees. *Conservation Biology* 12:442–450.
- PURCELL, K. L., J. VERNER, AND L. W. ORING. 1997. A comparison of the breeding ecology of birds nesting in boxes and tree cavities. *Auk* 114:646–656.
- PYKE, D. A. 1994. Ecological significance of seed banks with special reference to alien annuals. Pp. 197–201 in S. B. Monsen, and S. G. Kitchen (compilers). *Proceedings—ecology and management of annual rangelands*. USDA Forest Service General Technical Report INT-GTR-313. USDA Forest Service, Intermountain Research Station, Ogden, UT.
- PYKE, D. A. 2000. Invasive exotic plants of sagebrush ecosystems of the intermountain West. Pp. 43–54 in P. G. Entwistle, A. M. DeBolt, J. H. Kaltenecker, and K. Steenhof (compilers). *Proceedings: sagebrush steppe ecosystems symposium*. USDI Bureau of Land Management Publication No. BLM/ID/PT-001001+1150. Boise, ID.
- PYKE, D. A., AND S. J. NOVAK. 1994. Cheatgrass demography—establishment attributes, recruitment, ecotypes, and genetic variability. Pp. 12–21 in S. B. Monsen, and S. G. Kitchen (compilers). *Proceedings—ecology and management of annual rangelands*. USDA Forest Service General Technical Report INT-GTR-313. USDA Forest Service, Intermountain Research Station, Ogden, UT.
- PYLE, W. H., AND J. A. CRAWFORD. 1996. Availability of foods of Sage-Grouse chicks following prescribed fire in sagebrush-bitterbrush. *Journal of Range Management* 49:320–324.
- PYNE, S. J. 1982. *Fire in America: a cultural history of wildland and rural fire*. Princeton University Press, Princeton, NJ.
- PYNE, S. J. 1984. *Introduction to wildland fire*. John Wiley, New York, NY.
- PYNE, S. J. 1993. Keeper of the flame: a survey of anthropogenic fire. Pp. 245–266 in P. J. Crutzen, and J. G. Goldammer (editors). *Fire in the environment: the ecological, atmospheric, and climatic importance of vegetation fires*. John Wiley and Sons, New York, NY.
- PYNE, S. J. 1996. *Nouvelle Southwest*. Pp. 10–16 in W. W. Covington, and P. K. Wagner (technical coordinators). *Conference on adaptive ecosystem restoration and management: restoration of cordilleran conifer landscapes of North America*. USDA Forest Service General Technical Report RM-GTR-278. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- PYNE, S., P. ANDREWS, AND R. LAVEN. 1996. *Introduction to wildland fire (second edition)*. John Wiley and Sons, New York, NY.
- QUINN, R. D. 1994. Animals, fire, and vertebrate herbivory in Californian chaparral and other Mediterranean-type ecosystems. Pp. 46–78 in J. M. Moreno, and W. C. Oechel (editors). *The role of fire in Mediterranean-type ecosystems*. Springer-Verlag, New York, NY.
- RALPH, C. J., P. W. C. PATON, AND C. A. TAYLOR. 1991. Habitat association pattern of breeding birds and small mammals in Douglas-fir/hardwood stands in northwestern California and southwestern Oregon. Pp. 379–394 in L. F. Ruggiero, K. B. Aubry, A. B. Carey, and M. H. Huff (technical coordinators). *Wildlife and vegetation of unmanaged Douglas-fir forests*. USDA Forest Service General Technical Report PNW-GTR-285. USDA Forest Service, Pacific Northwest Research Station, Portland, OR.
- RAPHAEL, M. G., M. L. MORRISON, AND M. P. YODER-WILLIAMS. 1987. Breeding bird populations during twenty-five years of post-fire succession. *Condor* 89: 614–626.
- RAPHAEL, M. G., K. V. ROSENBERG, AND B. G. MARCOT. 1988. Large-scale changes in bird populations of Douglas-fir forests, northwestern California. Pp. 63–83 in J. A. Jackson (editor). *Bird conservation III*. University of Wisconsin Press, Madison, WI.
- REDMOND, R. L., AND D. A. JENNI. 1986. Population ecology of the Long-billed Curlew (*Numenius americanus*) in western Idaho. *Auk* 103:755–767.
- REICH, P. B., M. D. ABRAMS, D. S. ELLSWORTH, E. L. KRUGER, AND T. J. TABONE. 1990. Fire affects ecophysiology and community dynamics of central Wisconsin oak forest regeneration. *Ecology* 71:2179–2190.
- REICHMAN, O. J. 1987. *Konza Prairie*. University of Kansas Press, Lawrence, KS.
- REINKENSMeyer, D. P. 2000. *Habitat associations of bird communities in shrub-steppe and western juniper woodlands*. M.S. thesis, Oregon State University, Corvallis, OR.
- REINKING, D. L., AND D. P. HENDRICKS. 1993. Occurrence and nesting of Henslow's Sparrow in Oklahoma. *Bulletin of the Oklahoma Ornithological Society* 26: 33–36.
- REINKING, D. L., D. A. WIEDENFELD, D. H. WOLFE, AND R. W. ROHRBAUGH, JR. 2000. Distribution, habitat use, and nesting success of Henslow's Sparrow in Oklahoma. *Prairie Naturalist* 32:219–232.
- RENWALD, J. D. 1977. Effect of fire on Lark Sparrow nesting densities. *Journal of Range Management* 30:283–285.
- RIEGEL, G. M., B. G. SMITH, AND J. F. FRANKLIN. 1992. Foothill oak woodlands of the interior valleys of southwest Oregon. *Northwest Science* 66:66–76.
- RITTER, L. V. 1988a. Blue oak woodland. Pp. 74–75 in K. E. Mayer, and W. F. Laudenslayer (editors). *A guide to wildlife habitats of California*. California Department of Forestry, Sacramento, CA.
- RITTER, L. V. 1988b. Valley oak woodland. Pp. 76–77 in K. E. Mayer, and W. F. Laudenslayer (editors). *A guide to wildlife habitats of California*. California Department of Forestry, Sacramento, CA.
- ROBBINS, C. S., D. K. DAWSON, AND B. A. DOWELL. 1989.

- Habitat area requirements of breeding forest birds of the Middle Atlantic States. *Wildlife Monographs* 103: 1–34.
- ROBBINS, L. E., AND R. L. MYERS. 1992. Seasonal effects of prescribed burning in Florida: a review. Tall Timbers Research Miscellaneous Publication 8.
- ROBBINS, M. B., AND M. A. ORTEGA-HUERTA. 2002. Major negative impacts of early intensive cattle stocking on tallgrass prairies: the case of the Greater Prairie-Chicken (*Tympanuchus cupido*). *North American Birds* 56:239–244.
- ROBEL, R. J., J. P. HUGHES, S. D. HULL, K. E. KEMP, AND D. S. KLUTE. 1998. Spring burning: resulting avian abundance and nesting in Kansas CRP. *Journal of Range Management* 51:132–138.
- ROBERTSON, J. H. 1954. Half-century changes on northern Nevada ranges. *Journal of Range Management* 7: 117–121.
- ROBICHAUX, R. H. (EDITOR). 1999. *Ecology of Sonoran desert plants and plant communities*. University of Arizona Press, Tucson, AZ.
- ROGERS, G. F. 1985. Mortality of burned *Cereus giganteus*. *Ecology* 66:630–632.
- ROGERS, G. F., AND J. STEELE. 1980. Sonoran desert fire ecology. Pp. 15–19 in M. A. Stokes, and J. H. Dieterich (technical coordinators). *Proceedings of the fire history workshop*. USDA Forest Service General Technical Report RM-81. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- ROGERS, R. E., AND D. E. SAMUEL. 1984. Ruffed Grouse brood use of oak-hickory managed with prescribed burning. *Transactions of the Northeast Section of The Wildlife Society* 40:142–154.
- ROHRBAUGH, R. W., JR., D. L. REINKING, D. H. WOLFE, S. K. SHERROD, AND M. A. JENKINS. 1999. Effects of prescribed burning and grazing on nesting and reproductive success of three grassland passerine species in tallgrass prairie. *Studies in Avian Biology* 19:165–170.
- ROLAND, J. 1993. Large-scale forest fragmentation increases the duration of tent caterpillar outbreaks. *Oecologia* 93:25–30.
- ROLLINS, M. G., P. MORGAN, AND T. SWETNAM. 2002. Landscape-scale controls over 20th century fire occurrence in two large Rocky Mountain (USA) wilderness areas. *Landscape Ecology* 17:539–557.
- ROMME, W. H. 1982. Fire and landscape diversity in subalpine forests of Yellowstone National Park. *Ecological Monographs* 52:199–221.
- ROMME, W. H., AND D. G. DESPAIN. 1989. The long history of fire in the Greater Yellowstone Ecosystem. *Wildlands* 15:10–17.
- ROMME, W. H., AND D. H. KNIGHT. 1981. Fire frequency and subalpine forest succession along a topographic gradient in Wyoming. *Ecology* 62:319–326.
- ROMME, W. H., M. L. FLOYD-HANNA, AND D. D. HANNA. 2003. Ancient pinyon-juniper forests of Mesa Verde and the West: a cautionary note for forest restoration programs. Pp. 335–350 in *Fire, fuel treatments, and ecological restoration: conference proceedings*. USDA Forest Service Proceedings RMRS-P-29. USDA Forest Service, Rocky Mountain Research Station, Fort Collins, CO.
- ROSENBERG, K. V., R. D. OHMART, W. C. HUNTER, AND B. W. ANDERSON. 1991. *Birds of the lower Colorado River valley*. University of Arizona Press, Tucson, AZ.
- ROSSI, R. S. 1980. History of cultural influences on the distribution and reproduction of oaks in California. Pp. 7–18 in T. R. Plumb (technical coordinator). *Ecology and utilization of California oaks*. USDA Forest Service General Technical Report PSW-44. USDA Forest Service, Pacific Southwest Research Station, Berkeley, CA.
- ROTBERRY, J. T. 1985. The role of habitat in avian community composition: physiognomy or floristics? *Oecologia* 67:213–217.
- ROTBERRY, J. T. 1998. Avian conservation research needs in western shrublands: exotic invaders and the alteration of ecosystem processes. Pp. 262–272 in J. M. Marzluff, and R. Sallabanks (editors). *Avian conservation: research and management*. Island Press, Washington, DC.
- ROTBERRY, J. T., AND S. T. KNICK. 1999. Multiscale habitat associations of the Sage Sparrow: implications for conservation biology. *Studies in Avian Biology* 19: 95–103.
- ROTBERRY, J. T., AND J. A. WIENS. 1978. Nongame bird communities in northwestern rangelands. Pp. 32–46 in R. M. DeGraaf (technical coordinator). *Proceedings of a workshop on nongame bird habitat management in the coniferous forests of the western United States*. USDA Forest Service General Technical Report PNW-64. USDA Forest Service, Pacific Northwest Research Station, Portland, OR.
- ROTBERRY, J. T., AND J. A. WIENS. 1980. Habitat structure, patchiness, and avian communities in North American steppe vegetation: A multivariate analysis. *Ecology* 61:1228–1250.
- ROTBERRY, J. T., R. J. COOPER, J. M. WUNDERLE, AND K. G. SMITH. 1995. When and how are populations limited: the roles of insect outbreaks, fire, and other natural perturbations. Pp. 55–84 in T. E. Martin, and D. M. Finch (editors). *Ecology and management of Neotropical migratory birds: a synthesis and review of critical issues*. Oxford University Press, New York, NY.
- ROUNDY, B. A., E. D. MCARTHUR, J. S. HALEY, AND D. K. MANN (COMPILERS). 1995. *Proceedings: wildland shrub and arid land restoration symposium*. USDA Forest Service General Technical Report INT-GTR-315. USDA Forest Service, Intermountain Research Station, Ogden, UT.
- RUDNICKY, J. L., W. A. PATTERSON, AND R. P. COOK. 1999. Experimental use of prescribed fire for managing grassland bird habitat at Floyd Bennett Field, Brooklyn, New York. Pp. 99–118 in P. D. Vickery, and P. W. Dunwiddie (editors). *Grasslands of northeastern North America*. Massachusetts Audubon Society, Lincoln, MA.

- RUFFNER, C. M., AND M. D. ABRAMS. 1998. Lightning strikes and resultant fires from archival (1912–1917) and current (1960–1997) information in Pennsylvania. *Journal of the Torrey Botanical Society* 125:249–252.
- RUGGIERO, L. F., K. B. AUBRY, A. B. CAREY, AND M. H. HUFF (TECHNICAL COORDINATORS). 1991. Wildlife and vegetation of unmanaged douglas-fir forests. USDA Forest Service General Technical Report PNW-GTR-285. USDA Forest Service, Pacific Northwest Research Station, Portland, OR.
- RUNDEL, P. W., D. J. PARSONS, AND D. T. GORDON. 1977. Montane and subalpine vegetation of the Sierra Nevada and Cascade Ranges. Pp. 559–599 in M. G. Barbour, and J. Major (editors). *Terrestrial vegetation of California*. Wiley-Interscience, New York, NY.
- RUSSELL, E. W. B. 1983. Indian-set fires in the forests of the northeastern United States. *Ecology* 64:78–88.
- SAAB, V. A., AND J. G. DUDLEY. 1998. Responses of cavity nesting birds to stand-replacement fire and salvage logging in ponderosa pine/Douglas-fir forests of southwestern Idaho. USDA Forest Service Research Paper RMRS-RP-11. USDA Forest Service, Rocky Mountain Research Station, Fort Collins, CO.
- SAAB, V., AND K. VIERLING. 2001. Reproductive success of Lewis's Woodpecker in burned pine and cottonwood riparian forests. *Condor* 103:491–501.
- SAAB, V. A., J. DUDLEY, AND W. L. THOMPSON. 2004. Factors influencing occupancy of nest cavities in recently burned forests. *Condor* 106:20–36.
- SAAB, V. A., R. BRANNON, J. DUDLEY, L. DONOHOO, D. VANDERZANDEN, V. JOHNSON, AND H. LACHOWSKI. 2002. Selection of fire-created snags at two spatial scales by cavity-nesting birds. Pp. 835–848 in P. J. Shea, W. F. Laudenslayer, Jr., B. Valentine, C. P. Weatherspoon, and T. E. Lisle (technical coordinators). USDA Forest Service General Technical Report PSW-GTR-181. USDA Forest Service, Pacific Southwest Research Station, Albany, CA.
- SAENZ, D., R. N. CONNER, D. C. RUDOLPH AND R. T. ENGSTROM. 2001. Is a "hands-off" approach appropriate for Red-cockaded Woodpecker conservation in twenty-first-century landscapes? *Wildlife Society Bulletin* 29:956–966.
- SAMPSON, A. W. 1944. Plant succession on burned chaparral lands in Northern California. University of California. College of Agriculture, Agricultural Experiment Station, Bulletin 685, Berkeley, CA.
- SAMSON, F. B. 1980. Island biogeography and the conservation of nongame birds. *Transactions of the North American Wildlife and Natural Resources Conference* 45:245–251.
- SAMSON, F., AND F. KNOPF. 1994. Prairie conservation in North America. *BioScience* 44:418–421.
- SANDERSON, H. R., E. L. BULL, AND P. J. EDGERTON. 1980. Bird communities in mixed conifer forests of the interior Northwest. Pp. 224–237 in R. M. DeGraff, and N. G. Tilghman (editors). *Management of western forests and grasslands for nongame birds: workshop proceedings*. USDA Forest Service General Technical Report INT-86. USDA Forest Service, Intermountain Research Station, Ogden, UT.
- SAUER, J. R., J. E. HINES, AND J. FALLON. 2001. The North American breeding bird survey, results and analysis 1966–2000. Version 2001.2, USGS Patuxent Wildlife Research Center, Laurel, MD.
- SAVAGE, M., AND T. W. SWETNAM. 1990. Early 19th-century fire decline following sheep pasturing in a Navajo ponderosa pine forest. *Ecology* 71:2374–2378.
- SAWYER, J. O., D. A. THORNBURGH, AND J. R. GRIFFIN. 1977. Mixed evergreen forest. Pp. 359–381 in M. G. Barbour, and J. Major (editors). *Terrestrial vegetation of California*. Wiley-Interscience, New York, NY.
- SCHIECK, J., AND K. A. HOBSON. 2000. Bird communities associated with live residual tree patches within cut blocks and burn habitat in mixed woodland boreal forests. *Canadian Journal of Forest Research* 30:1281–1295.
- SCHIECK, J. AND S. J. SONG. 2002. Responses of boreal birds to wildfire and harvesting. Pp. 9–1–9–45 in S. J. Song (editor). *Ecological basis for stand management: a synthesis of ecological responses to wildfire and harvesting*. Alberta Research Council Inc., Vegreville, AB, Canada.
- SCHIECK, J., M. NIETTFELD, AND J. B. STELFOX. 1995. Differences in bird species richness and abundance among three successional stages of aspen-dominated boreal forests. *Canadian Journal of Zoology* 73:1417–1431.
- SCHIECK, J., K. STUART-SMITH, AND M. NORTON. 2000. Bird communities are affected by amount and dispersion of vegetation retained in mixed woodland boreal forest harvest areas. *Forest Ecology and Management* 126:239–254.
- SCHLESINGER, W. H., J. F. REYNOLDS, G. L. CUNNINGHAM, L. F. HUENNEKE, W. M. JARELL, R. A. VIRGINIA, AND W. G. WHITFORD. 1990. Biological feedbacks in global desertification. *Science* 247:1043–1048.
- SCHMID, M. K., AND G. F. ROGERS. 1988. Trends in fire occurrence in the Arizona upland subdivision of the Sonoran desert, 1955–1983. *Southwestern Naturalist* 33:437–444.
- SCHMIDT, K. M., J. P. MENAKIS, C. C. HARDY, W. J. HANN, AND D. L. BUNNELL. 2002. Development of coarse-scale spatial data for wildland fire and fuel management. USDA Forest Service General Technical Report RMRS-GTR-87. USDA Forest Service, Rocky Mountain Research Station, Fort Collins, CO.
- SCHMIEGELOW, F. K. A., D. STEPNIKY AND C. STAMBAUGH. 2001. Post-fire use of burned and salvage-logged areas by boreal birds in a mixed woodland forest. Report to Sustainable Ecosystems Research Users Group, Alberta Environment, Edmonton, AB, Canada.
- SCHOENNAGEL, T., T. T. VEBLEN, AND W. H. ROMME. 2004. The interaction of fire, fuels, and climate across Rocky Mountain forests. *BioScience* 54:661–676.
- SCHROEDER, M. A., AND L. A. ROBB. 1993. Greater Prairie-Chicken. In A. Poole, and F. Gill (editors). *The Birds of North America*, No. 36. The Academy of Natural

- Sciences, Philadelphia, PA, and The American Ornithologists' Union, Washington, DC.
- SCHROEDER, M. H., AND D. L. STURGES. 1975. The effect on the Brewer's Sparrow of spraying big sagebrush. *Journal of Range Management* 28:294–297.
- SCHULENBERG, J. H., G. L. HORAK, M. D. SCHWILLING, AND E. J. FINK. 1994. Nesting of Henslow's Sparrow in Osage County, Kansas. *Kansas Ornithological Society Bulletin* 44:25–28.
- SCHWARZ, G. F. 1907. *The longleaf pine in virgin forest*. John Wiley and Sons, New York, NY.
- SCHWEITZER, D. F., AND T. J. RAWINSKI. 1988. Northeastern pitch pine/scrub oak barrens. Eastern Heritage Task Force, The Nature Conservancy, Boston, MA.
- SEAMAN, B. D., AND D. G. KREMENTZ. 2000. Movements and survival of Bachman's Sparrow in response to prescribed summer burns in South Carolina. *Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies* 54:227–240.
- SEDGWICK, J. A. 1987. Avian habitat relationships in pinyon-juniper woodland. *Wilson Bulletin* 99:413–431.
- SEDGWICK, J. A. 2000. Willow Flycatcher (*Empidonax traillii*). In A. Poole, and F. Gill (editors). *The Birds of North America*, No. 533. The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists' Union, Washington, DC.
- SEDGWICK, J. A., AND R. A. RYDER. 1987. Effects of chaining on nongame wildlife. Pp. 541–551. in R. L. Everitt (compiler). *Proceedings of the pinyon-juniper conference*. USDA Forest Service General Technical Report INT-215. USDA Forest Service, Intermountain Research Station, Ogden, UT.
- SETTERINGTON, M. A., I. D. THOMPSON, AND W. A. MONTEVECCHI. 2000. Woodpecker abundance and habitat use in mature balsam fir forests in Newfoundland. *Journal of Wildlife Management* 64:335–345.
- SEVERSON, K. E., AND J. N. RINNE. 1990. Increasing habitat diversity in southwestern forests and woodlands via prescribed burning. Pp. 94–104 in J. S. Krammes (technical coordinator). *Effects of fire management of southwestern natural resources*. USDA Forest Service General Technical Report RM-191. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- SHAW, N. L., AND S. B. MONSEN. 2000. Controlling annual grasses with OUST herbicide. Pp. 66–69 in P. G. Entwistle, A. M. DeBolt, J. H. Kaltenecker, and K. Steenhof (compilers). *Proceedings: sagebrush steppe ecosystems symposium*. Bureau of Land Management Publication No. BLM/ID/PT-001001+1150. Boise, ID.
- SHAW, N. L., V. A. SAAB, S. B. MONSEN, AND T. D. RICH. 1999. *Bromus tectorum* expansion and biodiversity loss on the Snake River plain, southern Idaho, U.S.A. Pp. 586–588 in D. Eldridge, and D. Freudenberger (editors). *People and rangelands: building the future*. *Proceedings of the International Rangeland Congress*, Townsville, Australia.
- SHINNEMAN, D. J., AND W. L. BAKER. 1997. Nonequilibrium dynamics between catastrophic disturbances and old-growth forests in ponderosa pine landscapes of the Black Hills. *Conservation Biology* 11:1276–1288.
- SHRIVER, W. G., AND P. D. VICKERY. 2001. Response of breeding Florida Grasshopper and Bachman's sparrows to winter prescribed burning. *Journal of Wildlife Management* 65:470–475.
- SHRIVER, W. G., P. D. VICKERY, AND D. W. PERKINS. 1999. The effects of summer burns on breeding Florida Grasshopper and Bachman's sparrows. *Studies in Avian Biology* 19:144–148.
- SHRIVER, W. G., A. L. JONES, P. D. VICKERY, A. WEIK, AND J. V. WELLS. In press. The distribution and abundance of obligate grassland birds breeding in New England and New York. In C. J. Ralph, and T. D. Rich (editors). *Bird conservation implementation and integration in the Americas*. USDA Forest Service General Technical Report PSW-GTR-191. USDA Forest Service, Pacific Southwest Research Station, Albany, CA.
- SHUMAR, M. L., AND J. E. ANDERSON. 1986. Gradient analysis of vegetation dominated by two sub-species of big sagebrush. *Journal of Range Management* 39:156–160.
- SHUMWAY, D. L., M. D. ABRAMS, AND C. M. RUFFNER. 2001. A 400-year history of fire and oak recruitment in an old-growth oak forest in western Maryland. *Canadian Journal of Forest Research* 31:1437–1443.
- SIMBERLOFF, D. 1998. Flagships, umbrellas, and keystones: is single-species management passé in the landscape era? *Biological Conservation* 83:247–257.
- SIMBERLOFF, D. 2001. Management of boreal forest biodiversity—a view from the outside. *Scandinavian Journal of Forest Research*. Supplement 3:105–118.
- SKAGEN, S. K., C. P. MELCHER, W. H. HOWE, AND F. L. KNOPF. 1998. Comparative use of riparian corridors and oases by migrating birds in southeast Arizona. *Conservation Biology* 12:896–909.
- SKINNER, C. N. 1995. Change in spatial characteristics of forest openings in the Klamath Mountains of northwestern California, U.S.A. *Landscape Ecology* 10:219–228.
- SKINNER, C. N., AND C. CHANG. 1996. Fire regimes, past and present. Pp. 1041–1069 in *Status of the Sierra Nevada: Sierra Nevada Ecosystem Project Final Report to Congress Volume II. Wildland Resources Center Report No. 37*. Center for Water and Wildland Resources, University of California, Davis, CA.
- SMITH, C. R. 1997. Use of public grazing lands by Henslow's Sparrows, Grasshopper Sparrows, and associated grassland birds in New York state. Pp. 171–186 in P. D. Vickery, and P. W. Dunwiddie (editors). *Grasslands of northeastern North America*. Massachusetts Audubon Society, Lincoln, MA.
- SMITH, E. F., AND C. E. OWENSBY. 1978. Intensive early stocking and season-long stocking of Kansas Flint Hills range. *Journal of Range Management* 31:14–17.
- SMITH, J. K. (EDITOR). 2000. *Wildland fire in ecosystems:*

- effects of fire on fauna. USDA Forest Service General Technical Report RMRS-GTR-42-vol 1. USDA Forest Service, Rocky Mountain Research Station, Ogden, UT.
- SMITH, K. G. 1980. Nongame birds of the Rocky Mountain spruce-fir forests and their management. Pp. 258-279 in R. M. DeGraff, and N. G. Tilghman (editors). Management of western forests and grasslands for nongame birds: workshop proceedings. USDA Forest Service General Technical Report INT-86. USDA Forest Service, Intermountain Research Station, Ogden, UT.
- SMITH, K. T., AND E. K. SUTHERLAND. 1999. Fire-scar formation and compartmentalization in oak. Canadian Journal of Forest Research 29:166-171.
- SMITH, S. D., T. E. HUXMAN, S. F. ZITSER, T. N. CHARLET, D. C. HOUSMAN, J. S. COLEMAN, L. K. FENSTERMAKER, J. R. SEEMANN, AND R. S. NOWAK. 2000. Elevated CO₂ increases productivity and invasive species success in an arid ecosystem. Nature 408:79-82.
- SMITH, W. B., J. S. VISSAGE, D. R. DARR, AND R. M. SHEFFIELD. 2001. Forest resources of the United States, 1997. USDA Forest Service General Technical Report NC-219. USDA Forest Service, Northcentral Forest Experiment Station, St. Paul, MN.
- SMITH, W. P. 1985. Plant associations within the interior valleys of the Umpqua River Basin, Oregon. Journal of Range Management 38:526-530.
- SPARKS, J. C., R. E. MASTERS, D. M. ENGLE, M. E. PAYTON, AND G. A. BUKENHOFER. 1999. Influence of fire season and fire behavior on woody plants in Red-cockaded Woodpecker clusters. Wildlife Society Bulletin 27: 124-133.
- STEARNS, F. S. 1949. Ninety years of change in a northern hardwood forest in Wisconsin. Ecology 30:350-358.
- STEINAUER, E. M., AND S. L. COLLINS. 1996. Prairie ecology—the tallgrass prairie. Pp. 39-52 in F. B. Samson, and F. L. Knopf (editors). Prairie conservation. Island Press, Washington, DC.
- STELFOX, J. B. 1995. Relationships between stand age, stand structure, and biodiversity in aspen mixed woodland forests in Alberta. Alberta Environmental Centre (AECV95-R1), Vegreville, AB, and Canadian Forest Service (Project No. 0001A), Edmonton, AB, Canada.
- STEPHENS, S. L. 1997. Fire history of a mixed oak-pine forest in the foothills of the Sierra Nevada, El Dorado County, California. Pp. 191-198 in N. H. Pillsbury, J. Verner, and W. D. Tietje, (technical coordinators). Proceedings of a symposium on oak woodlands: ecology, management, and urban interface issues. USDA Forest Service General Technical Report PSW-160. USDA Forest Service, Pacific Southwest Research Station, Albany, CA.
- STERLING, J. C. 1999. Gray Flycatcher (*Empidonax wrightii*). In A. Poole, and F. Gill (editors). The Birds of North America, No. 458. The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists' Union, Washington, DC.
- STEUTER, A. A. 1991. Human impacts on biodiversity in America: thoughts from the grassland biome. Conservation Biology 5:136-137.
- STEWART, G., AND A. C. HULL. 1949. Cheatgrass (*Bromus tectorum* L.) an ecologic intruder in southern Idaho. Ecology 30:58-74.
- STEWART, O. C. 1955. Forest and grass burning in the Mountain West. Southwestern Lore 21:5-9.
- STEWART, O. C. 2002. Forgotten fires: Native Americans and the transient wilderness. University of Oklahoma Press, Norman, OK.
- STODDARD, H. L. 1931. The Bobwhite Quail: its habitats, preservation, and increase. Charles Scribner's Sons, New York, NY.
- STODDARD, H. L. 1963. Fire and birds in the mountains of southern Arizona Pp. 163-175 in Proceedings of the Tall Timbers Fire Ecology Conference #2. Tall Timbers Research Station, Tallahassee, FL.
- STRONG, T. R., AND C. E. BOCK. 1990. Bird species distribution patterns in riparian habitats in southeastern Arizona. Condor 92:866-885.
- STUART, J. D. 1987. Fire history of an old-growth forest of *Sequoia sempervirens* (Taxodiaceae) forest in Humboldt Redwoods State Park, California. Madroño 34:128-141.
- STUART, J. D., AND L. A. SALAZAR. 2000. Fire history of California white fir forests in the coastal mountains of northwestern California. Northwest Science 74: 280-285.
- STUBBS, M. M. 2000. Time-based nitrogen availability and risk of *Taeniatherum caput-medusae* (L.) Nevski invasion in central Oregon. M.S. thesis, Oregon State University, Corvallis, OR.
- SUTHERLAND, E. K. 1997. History of fire in a southern Ohio second-growth mixed-oak forest. Pp. 172-183 in S. G. Pallardy, R. A. Cecich, H. E. Garrett, and P. S. Johnson (editors). Proceedings of the central hardwood forest conference. USDA Forest Service General Technical Report NC-188. USDA Forest Service, North Central Research Station, St. Paul, MN.
- SUTHERLAND, E. K., T. F. HUTCHINSON, AND D. A. YAUSSY. 2003. Introduction, study area description and experimental design. Pp. 1-16 in E. K. Sutherland, and T. F. Hutchinson (editors). Characteristics of mixed-oak forest ecosystems in southern Ohio prior to the reintroduction of fire. USDA Forest Service General Technical Report NE-299. USDA Forest Service, Northeastern Research Station, St. Paul, MN.
- SWANTON, J. R. 1946. The Indians of the southeastern United States. Smithsonian Institution Bureau of American Ethnology Bulletin No. 137. U.S. Government Printing Office, Washington, DC.
- SWENGEL, A. B. 2001. A literature review of insect responses to fire, compared to other conservation managements of open habitat. Biodiversity and Conservation 10: 1141-1169.
- SWENGEL, S. R. 1996. Management responses of three species of declining sparrows in tallgrass prairie. Bird Conservation International 6:241-253.
- SWETNAM, T. W., AND C. H. BAISAN. 1996. Historical fire

- regime patterns in the southwestern United States since AD 1700. Pp. 11–32 in C. D. Allen (editor). Proceedings of the Second La Mesa Fire Symposium. USDA Forest Service General Technical Report RM-GTR-286. USDA Forest Service, Rocky Mountain Research Station, Fort Collins, CO.
- SWETNAM, T. W., C. D. ALLEN, AND J. L. BETANCOURT. 1999. Applied historical ecology: using the past to manage for the future. *Ecological Applications* 9:1189–1206.
- SWIECKI, T. J., AND E. BERNHARDT. 1998. Understanding blue oak regeneration. *Fremontia* 26:19–26.
- SZARO, R. C. 1981. Bird population responses to converting chaparral to grassland and riparian habitats. *Southwestern Naturalist* 26:251–256.
- SZARO, R. C., AND M. D. JAKLE. 1985. Avian use of a desert riparian island and its adjacent scrub habitat. *Condor* 87:511–519.
- TAUSCH, R. J. 1999. Historic woodland development. Pp. 12–19 in S. B. Monsen, R. Stevens, R. J. Tausch, R. Miller, and S. Goodric, (compilers). Proceedings: ecology and management of pinyon-juniper communities within the interior West. USDA Forest Service General Technical Report RMRS-P-9. USDA Forest Service, Rocky Mountain Research Station, Ogden, UT.
- TAUSCH, R. J., AND N. E. WEST. 1995. Plant species composition patterns with differences in tree dominance on a south-western Utah pinon-juniper site. Pp. 16–23 in D. W. Shaw, E. F. Alson, and C. LoSapio (technical coordinators). Proceedings: desired future conditions for pinon-juniper ecosystems. USDA Forest Service General Technical Report RM-258. USDA Forest Service, Rocky Mountain Research Station, Fort Collins, CO.
- TAYLOR, A. H. 1993. Fire history and structure of red fir (*Abies magnifica*) forests, Swain Mountain Experimental Forest, Cascade Range, northeastern California. *Canadian Journal of Forest Research* 23: 1672–1678.
- TAYLOR, A. H., AND C. B. HALPERN. 1991. The structure and dynamics of *Abies magnifica* forests in the southern Cascades Range, U.S.A. *Journal of Vegetation Science* 2:189–200.
- TAYLOR, A. H., AND C. N. SKINNER. 1997. Fire regimes and management of old-growth Douglas-fir forests in the Klamath Mountains of northwestern California. Pp. 203–208 in J. E. Greenlee (editor). Proceedings: fire effects on threatened and endangered species and habitats conference. International Association of Wildland Fire, Fairfield, WA.
- TAYLOR, A. H., AND C. N. SKINNER. 1998. Fire history and landscape dynamics in late-successional reserve, Klamath Mountains, California, U.S.A. *Forest Ecology and Management* 111:285–301.
- TAYLOR, A. H., AND C. N. SKINNER. 2003. Spatial patterns and controls on historical fire regimes and forest structure in the Klamath Mountains. *Ecological Applications* 13:704–719.
- TAYLOR, A. H., AND M. N. SOLEM. 2001. Fire regimes and stand dynamics in an upper montane forest landscape in the southern Cascades, Caribou Wilderness, California. *Journal of the Torrey Botanical Society* 128:350–361.
- TAYLOR, D. L., AND W. J. BARMORE, JR. 1980. Post-fire succession of avifauna in coniferous forests of Yellowstone and Grand Teton National Parks, Wyoming. Pp. 130–145 in Workshop proceedings of the management of western forests and grasslands for nongame birds. USDA Forest Service General Technical Report INT-GTR-86. USDA Forest Service, Intermountain Forest and Range Experiment Station, Ogden, UT.
- TAYLOR, K. L., AND R. W. FONDA. 1990. Woody fuel structure and fire in subalpine fir forests, Olympic National Park, Washington. *Canadian Journal of Forest Research* 20:193–199.
- TEENSMA, P. D. A. 1987. Fire history and fire regimes of the central western Cascades of Oregon. Ph. D. dissertation, University of Oregon, Eugene, OR.
- TEMPLE, S. A. 2002. Dickcissel (*Spiza americana*). In A. Poole, and F. Gill (editors). *The Birds of North America*, No. 703. The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists' Union, Washington, DC.
- THILENIUS, J. F. 1968. The *Quercus garryana* forests of the Willamette Valley, Oregon. *Ecology* 49:1124–1133.
- THOME, D. M., C. J. ZABEL, AND L. V. DILLER. 1999. Forest stand characteristics and reproduction of Northern Spotted Owls in managed north-coastal California forests. *Journal of Wildlife Management* 63:44–59.
- THOMPSON, F. R., III, AND R. M. DEGRAAF. 2001. Conservation approaches for woody, early successional communities in the eastern United States. *Wildlife Society Bulletin* 29:483–494.
- THOMPSON, I. D., H. A. HOGAN, AND W. A. MONTEVECCHI. 1999. Avian communities of mature balsam fir forests in Newfoundland: age-dependence and implications for timber harvesting. *Condor* 101:311–323.
- THORNBURGH, D. A. 1982. Succession in the mixed evergreen forests of northwestern California. Pp. 87–91 in J. Means (editor). *Forest succession and stand development research in the Northwest*. Oregon State University, Corvallis, OR.
- TIEDEMANN, A. R., J. O. KLMMEDSON, AND E. L. BULL. 2000. Solution of forest health problems with prescribed fire: are forest productivity and wildlife at risk? *Forest Ecology and Management* 127:1–18.
- TIETJE, W. D., AND J. K. VREELAND. 1997. Vertebrates diverse and abundant in well-structured oak woodland. *California Agriculture* 51:8–14.
- TIETJE, W. D., J. K. VREELAND, AND W. H. WEITKAMP. 2001. Live oak saplings survive prescribed fire and sprout. *California Agriculture* 55:18–27.
- TIMBROOK, J., J. R. JOHNSON, AND D. D. EARLE. 1982. Vegetation burning by the Chumash. *Journal of California and Great Basin Anthropology* 4:163–186.
- TITTLER, R., S. J. HANNON, AND M. NORTON. 2001. Residual tree retention does not ameliorate negative effects of clearcutting for most boreal songbirds in the short-term. *Ecological Applications* 11:1656–1666.

- TOBALSKE, B. 1997. Lewis' Woodpecker (*Melanerpes lewis*). In A. Poole, and F. Gill (editors). The Birds of North America, No. 284. The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists' Union, Washington, DC.
- TOMOFF, C. S. 1974. Avian species diversity in desert scrub. *Ecology* 55:396–403.
- TOPIK, C. 1989. Plant association and management guide for the grand fir zone, Gifford Pinchot National Forest. USDA Forest Service Technical Report R6-ECOL-TP-006-88. USDA Forest Service, Pacific Northwest Region, Portland, OR.
- TOPIK, C., N. M. HALVERSON, AND D. G. BROCKWAY. 1987. Plant association and management guide for the western hemlock zone, Gifford Pinchot National Forest. USDA Forest Service Technical Report R6-ECOL-230A-1986. USDA Forest Service, Pacific Northwest Region, Portland, OR.
- TOPIK, C., N. M. HALVERSON, AND T. HIGH. 1988. Plant association and management guide for the ponderosa pine, Douglas-fir, and grand fir zones, Mount Hood National Forest. USDA Forest Service Technical Report R6-ECOL-TP004-88. USDA Forest Service, Pacific Northwest Region, Portland, OR.
- TOWNE, G., AND C. OWENSBY. 1984. Long-term effects of annual burning at different dates in ungrazed Kansas tallgrass prairie. *Journal of Range Management* 37: 392–397.
- TURNER, R. M. 1982. Mojave desert scrub. *Desert Plants* 4: 157–168.
- TURNER, R. M., J. E. BOWERS, AND T. L. BURGESS. 1995. Sonoran desert plants. University of Arizona Press, Tucson, AZ.
- TURNER, R. M., R. H. WEBB, J. E. BOWERS, AND J. R. HASTINGS. 2003. The changing mile revisited. University of Arizona Press, Tucson, AZ.
- TVETEN, R. K., AND R. W. FONDA. 1999. Fire effects on prairies and oak woodlands on Fort Lewis, Washington. *Northwest Science* 73:145–158.
- UDVARDY, M. D. F. 1958. Ecological and distributional analysis of North American birds. *Condor* 60:50–66.
- UMBANHOWAR, C. E. 1996. Recent fire history of the northern Great Plains. *American Midland Naturalist* 135: 115–121.
- USDA FOREST SERVICE. 1986. Land and resource management plan: White Mountain National Forest. <http://www.fs.fed.us/r9/white/3_WM_fpr_Web/forest_plan/forest_plan.htm> (12 June 2004).
- USDA FOREST SERVICE. 2000. Managing the impacts of wildfires on communities and the environment: a report to the president in response to the wildfires of 2000. The national fire plan executive summary for the USDA Forest Service. U. S. Department of Agriculture, Washington, DC.
- USDA FOREST SERVICE. 2004. Fire effects information system—glossary. <<http://www.fs.fed.us/database/feis/glossary.html>> (12 September 2004).
- USDA FOREST SERVICE AND USDI BUREAU OF LAND MANAGEMENT. 1994a. Final supplemental environmental impact statement on management of habitat for late-successional and old-growth forest related species within the range of the Northern Spotted Owl. Northwest Forest Plan. 2 Vol. USDA Forest Service and USDI Bureau of Land Management, Portland, OR.
- USDA FOREST SERVICE AND USDI BUREAU OF LAND MANAGEMENT. 1994b. Record of Decision on Management of habitat for late-successional and old-growth related species within the range of the Northern Spotted Owl. Northwest Forest Plan. 2 Vol. USDA Forest Service and USDI Bureau of Land Management, Portland, OR.
- USDA FOREST SERVICE AND USDI BUREAU OF LAND MANAGEMENT. 2000. Final supplemental environmental impact statement for amendment to the survey and manage, protection buffer, and other mitigation measures standards and guidelines. Northwest Forest Plan. 2 Vol. USDA Forest Service and USDI Bureau of Land Management, Portland, OR.
- USDI. 1992. Recovery plan for the Northern Spotted Owl: final draft. U.S. Department of the Interior, Washington, DC.
- USDI. 1995. Recovery plan for the Mexican Spotted Owl (*Strix occidentalis lucida*). Volume 1. USDI Fish and Wildlife Service, Albuquerque, NM.
- USDI. 1996. Effects of military training and fire in the Snake River Birds of Prey National Conservation Area. BLM/IDARNG Research Project Final Report. USGS Biological Resources Division, Snake River Field Station, Boise, ID.
- USDI. 2002. Management considerations for sagebrush (*Artemisia*) in the western United States: a selective summary of current information about the ecology and biology of woody North American sagebrush taxa. USDI Bureau of Land Management, Washington, DC.
- USDI FISH AND WILDLIFE SERVICE 2000. Technical/agency draft revised recovery plan for the Red-cockaded Woodpecker (*Picoides borealis*). USDI Fish and Wildlife Service, Atlanta, GA
- USDI FISH AND WILDLIFE SERVICE. 2002. Birds of conservation concern 2002. Division of Migratory Bird Management, Arlington, Virginia. <<http://migratorybirds.fws.gov/reports/bcc2002.pdf>> (23 July 2004).
- USDI, USDA, DOE, AND OTHERS. 2001. Review and update of the 1995 federal wildland fire management policy. USDI, USDA, DOE, DOD, DOC, USEPA, FEMA, and National Association of State Foresters, National Interagency Fire Center. Boise, Idaho. <http://www.nifc.gov/fire_policy> (23 July 2004).
- VALE, T. R. 1974. Sagebrush conversion projects: an element of contemporary environmental change in the western United States. *Biological Conservation* 6: 274–284.
- VALONE, T. J., AND D. A. KELT. 1999. Fire and grazing in a shrub-invaded arid grassland community: independent or interactive ecological effects? *Journal of Arid Environments* 42:15–28.

- VALONE, T. J., M. MEYER, J. H. BROWN, AND R. M. CHEW. 2002. Timescale of perennial grass recovery in desertified arid grasslands following livestock removal. *Conservation Biology* 16:995–1002.
- VAN HOOSER, W. D., R. A. O'BRIEN, AND D. C. COLLINS. 1993. New Mexico's forest resources. USDA Forest Service Resource Bulletin INT-79. USDA Forest Service, Intermountain Research Station, Ogden, UT.
- VAN HORNE, B. 1983. Density as a misleading indicator of habitat quality. *Journal of Wildlife Management* 47: 893–901.
- VAN LEAR, D. H., AND J. M. WATT. 1993. The role of fire in oak regeneration. Pp. 66–78 in D. L. Loftis, and C. E. McGee (editors). *Oak regeneration: serious problems, practical recommendations*. USDA Forest Service General Technical Report SE-84. USDA Forest Service, Southeastern Forest Experiment Station, Asheville, NC.
- VAN NORMAN, K. 1998. Historical fire regimes in the Little River watershed, southwestern Oregon. M.S. thesis, Oregon State University, Corvallis, OR.
- VANDER HAEGEN, W. M., F. C. DOBLER, AND D. J. PIERCE. 2000. Shrubsteppe bird response to habitat and landscape variables. *Conservation Biology* 14:1145–1160.
- VEBLEN, T. T. 2000. Disturbance patterns in central Rocky Mountain forests. Pp. 33–56 in R. L. Knight, F. W. Smith, S. W. Buskirk, W. H. Romme, and W. L. Baker (editors). *Forest fragmentation in the southern Rocky Mountains*. University Press of Colorado, Niwot, CO.
- VEBLEN, T. T., AND LORENZ, D. C. 1991. *The Colorado Front Range: a century of ecological change*. University of Utah Press, Salt Lake City, UT.
- VEBLEN, T. T., T. KITZBERGER, AND J. DONNEGAN. 2000. Climatic and human influences on fire regimes in ponderosa pine forests in the Colorado Front Range. *Ecological Applications* 10:1178–1195.
- VEIRS, S. D. 1982. Coast redwood forest: Stand dynamics, successional status, and the role of fire. Pp. 119–141 in J. Means (editor). *Forest succession and stand development research in the Northwest*. Forest Research Laboratory, Oregon State University, Corvallis, OR.
- VERMEIRE, L. T., AND T. G. BIDWELL. 1998. Intensive early stocking. Oklahoma Cooperative Extension Service. Extension Facts F-2875. Oklahoma State University, Stillwater, OK.
- VERNER, J. 1988. Blue oak-digger pine. Pp. 80–81 in K. E. Mayer, and W. F. Laudenslayer (editors). *A guide to wildlife habitats of California*. California Department of Forestry, Sacramento, CA.
- VERNER, J., K. L. PURCELL, AND J. G. TURNER. 1997. Bird communities in grazed and ungrazed oak-pine woodlands at the San Joaquin Experimental Range. Pp. 381–390 in N. H. Pillsbury, J. Verner, and W. D. Tietje (technical coordinators). *Proceedings of a symposium on oak woodlands: ecology, management, and urban interface issues*. USDA Forest Service General Technical Report PSW-160. USDA Forest Service, Pacific Southwest Research Station, Albany, CA.
- VERNER, J., E. C. BEEDY, S. L. GRANHOLM, L. V. RITTER, AND E. F. TOTH. 1980. Birds. Pp. 75–319 in J. Verner, and A. S. Boss (technical coordinators). *California wildlife and their habitats: western Sierra Nevada*. USDA Forest Service General Technical Report PSW-37. USDA Forest Service, Pacific Southwest Research Station, Albany, CA.
- VICKERY, P. D. 1992. A regional analysis of endangered, threatened and special concern birds in the northeastern United States. *Transactions of the Northeast Section of the Wildlife Society* 48:1–10.
- VICKERY, P. D. 1996. Grasshopper Sparrow (*Ammodramus savannarum*). In A. Poole, and F. Gill (editors). *The Birds of North America*, No. 239. The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists' Union, Washington, DC.
- VICKERY, P. D. 2002. Effects of the size of prescribed fire on insect predation of northern blazing star, a rare grassland perennial. *Conservation Biology* 16:413–421.
- VICKERY, P. D., AND P. W. DUNWIDDIE. 1997. Introduction. Pages 1–13 in P. D. Vickery, and P. W. Dunwiddie (editors). *Grasslands of northeastern North America*. Massachusetts Audubon Society, Lincoln, MA.
- VICKERY, P. D., AND J. R. HERKERT. 2001. Recent advances in grassland bird research: where do we go from here? *Auk* 118:11–15.
- VICKERY, P. D., M. L. HUNTER, JR., AND S. M. MELVIN. 1994. Effects of habitat area on the distribution of grassland birds in Maine. *Conservation Biology* 8:1087–1097.
- VICKERY, P. D., M. L. HUNTER, JR., AND J. V. WELLS. 1992. Is density an indicator of breeding success? *Auk* 109: 706–710.
- VICKERY, P. D., M. L. HUNTER, JR., AND J. V. WELLS. 1999a. Effects of fire and herbicide treatment on habitat selection of grassland birds in southern Maine. *Studies in Avian Biology* 19:149–159.
- VICKERY, P. D., J. R. HERKERT, F. L. KNOPF, J. RUTH, AND C. E. KELLER. 2000. Grassland birds: An overview of threats and recommended management strategies. Pp. 74–77 in R. Bonney, D. N. Pashley, R. J. Cooper, and L. Niles (editors). *Strategies for bird conservation: the Partners in Flight planning process*. USDA Forest Service Proceedings RMRS-P-16. USDA Forest Service, Rocky Mountain Research Station, Ogden, UT.
- VICKERY, P. D., P. L. TUBARO, J. M. CARDOSO DA SILVA, B. G. PETERJOHN, J. R. HERKERT, AND R. B. CAVALCANTI. 1999b. Conservation of grassland birds in the western hemisphere. *Studies in Avian Biology* 19:2–26.
- VILLARD, M. A., AND J. SCHIECK. 1996. Immediate post-fire nesting by Black-backed Woodpeckers, *Picoides arcticus*, in northern Alberta. *Canadian Field Naturalist* 111:478–479.
- VINTON, M. A., D. C. HARTNETT, E. J. FINCK, AND J. M. BRIGGS. 1993. Interactive effects of fire, bison (*Bison bison*) grazing and plant community composition in tallgrass prairie. *American Midland Naturalist* 129: 10–18.
- VOGL, R. J. 1972. Fire in the southeastern grasslands. *Tall Timbers Fire Ecology Conference* 12:175–198.

- VREELAND, J. K., AND W. D. TIETJE. 1998. Initial response of woodrats to prescribed burning in oak woodland. *Transactions of the Western Section of the Wildlife Society* 34:21–31.
- VREELAND, J. K., AND W. D. TIETJE. 2002. Numerical response of small vertebrates to prescribed fire in a California oak woodland. Pp. 269–280 in R. Standiford, D. McCreary, and K. L. Purcell (technical coordinators). *Proceedings of the fifth symposium on oak woodlands: oaks in California's changing landscape*. USDA Forest Service General Technical Report PSW-184. USDA Forest Service, Pacific Southwest Research Station, Albany, CA.
- WADE D. D. 1993. Societal influences on prescribed burning. *Tall Timbers Fire Ecology Conference* 19: 351–356.
- WADE, D. D., B. L. BROCK, P. H. BROSE, J. B. GRACE, G. A. HOCH, AND W. A. PATTERSON III. 2000. Fire in eastern ecosystems. Pp. 53–96 in J. K. Brown, and J. K. Smith (editors). *Wildland fire in ecosystems: effects of fire on flora*. USDA Forest Service General Technical Report RMRS-GTR-42-vol. 2. USDA Forest Service, Rocky Mountain Research Station, Ogden, UT.
- WADE, D. D., G. CUSTER, J. THORSEN, P. KASKEY, J. KUSH, B. TWOMEY, AND D. VOLTOLINA. 1998. Reintroduction of fire into fire-dependent ecosystems: some southern examples. *Tall Timbers Fire Ecology Conference* 20: 94–98.
- WAHLENBERG, W. G. 1946. *Longleaf pine: its use, ecology, regeneration, protection, growth, and management*. Charles Lathrop Pack Forestry Foundation, Washington, DC.
- WALDROP, T. A., D. L. WHITE, AND S. M. JONES. 1992. Fire regimes for pine-grassland communities in the southeastern United States. *Forest Ecology and Management* 47:195–210.
- WALES, B. C. 2001. The management of insects, diseases, fire, and grazing and implications for terrestrial vertebrates using riparian habitats in Eastern Oregon and Washington. *Northwest Science* 75:119–127.
- WALTERS, C. J., AND C. S. HOLLING. 1990. Large-scale management experiments and learning by doing. *Ecology* 71:2060–2068.
- WAMBOLT, C. L., A. J. HARP, B. L. WELCH, N. SHAW, J. W. CONELLY, K. P. REESE, C. E. BRAUN, D. A. KLEBENOW, E. D. MCARTHUR, J. G. THOMPSON, L. A. TORELL, AND J. A. TANAKA. 2002. Conservation of Greater Sage-Grouse on public lands in the western U.S.: implications of recovery and management policies. Policy Analysis Center for Western Public Lands PACWPL Policy Paper SG-02-02. <<http://sagemap.wr.usgs.gov/documents.asp>> (6 July 2004).
- WARING, R. M., AND J. MAJOR. 1964. Some vegetation of the California coastal redwood region in relation to gradients of moisture, nutrients, light, and temperature. *Ecological Monographs* 34:167–215.
- WARNER, R. E. 1994. Agricultural land use and grassland habitat in Illinois: future shock for grassland birds? *Conservation Biology* 8:147–156.
- WATERHOUSE, F. L., M. H. MATHER, AND D. SEIP. 2002. Distribution and abundance of birds relative to elevation and biogeoclimatic zones in coastal old-growth forests in southern British Columbia. *British Columbia Journal of Ecosystems and Management*, 2:1–13.
- WEBB, R. G. 1970. *Reptiles of Oklahoma*. University of Oklahoma Press, Norman, OK.
- WEBER, M. G., AND B. J. STOCKS. 1998. Forest fires and sustainability in the boreal forests of Canada. *Ambio* 27:545–550.
- WEIK, A. 1998. Conservation of grassland birds in Maine: field survey of breeding birds, 1997–1998. Maine Department of Inland Fisheries and Wildlife Miscellaneous Report, Bangor, ME.
- WEIR, J. M. H., AND E. A. JOHNSON. 1998. Effects of escaped settlement fires and logging on forest composition in the mixed woodland boreal forest. *Canadian Journal of Forest Research* 28:459–467.
- WEIR, J. M. H., E. A. JOHNSON, AND K. MIYANISHI. 2000. Fire frequency and the spatial age mosaic of the mixed-wood boreal forest in western Canada. *Ecological Applications* 10:1162–1177.
- WEISBERG, P. J. 1998. Fire history, fire regimes, and development of forest structure in the central western Oregon Cascades. Ph.D. dissertation, Oregon State University, Corvallis, OR.
- WEISBERG, P. J. AND F. J. SWANSON. 2003. Regional synchrony in fire regimes of western Oregon and Washington, USA. *Forest Ecology and Management* 172:17–28.
- WELCH, B. L. 2002. Bird counts of burned versus unburned big sagebrush sites. USDA Forest Service Research Note RMRS-RN-16. USDA Forest Service, Rocky Mountain Research Station, Ogden, UT.
- WELLS, J. V., AND K. V. ROSENBERG. 1999. Grassland bird conservation in northeastern North America. *Studies in Avian Biology* 19:72–80.
- WELLS, P. 1962. Vegetation in relation to geological substratum and fire in the San Luis Obispo Quadrangle. *Ecological Monographs* 32:79–103.
- WELLS, P. V. 1970. Historical factors controlling vegetation patterns and floristic distributions in the central plains region of North America. Pp. 211–221 in W. Dort, Jr., and J. K. Jones (editors). *Pleistocene and recent environments of the central Great Plains*. University of Kansas Department of Geology Special Publication No. 3. University of Kansas Press, Lawrence, KS.
- WELSTEAD, K. 2002. Factors affecting nest predation of artificial and real Brewer's Sparrow nests in the South Okanagan and Similkameen Valleys, British Columbia. M.S. thesis, University of British Columbia, Vancouver, Canada.
- WELTZIN, J. F., S. ARCHER, AND R. K. HEITSCHMIDT. 1997. Small mammal regulation of vegetation structure in a temperate savanna. *Ecology* 78:751–763.
- WENDEL, G. W., AND H. C. SMITH. 1986. Effects of a prescribed fire in a central Appalachian oak-hickory stand. USDA Forest Service Research Paper NE-594. USDA Forest Service, Northeastern Forest Experiment Station, St. Paul, MN.

- WEST, N. E. 1979. Basic synecological relationships of sagebrush-dominated lands in the Great Basin and the Colorado Plateau. Pp. 33-41 in *The sagebrush ecosystem: a symposium*. Utah State University, College of Natural Resources, Logan, UT.
- WEST, N. E. 1983. Western intermountain sagebrush steppe. Pp. 351-397 in N. E. West (editor). *Ecosystems of the world 5: temperate deserts and semi-deserts*. Elsevier Scientific Publishing Co., New York, NY.
- WEST, N. E. 2000. Synecology and disturbance regimes of sagebrush steppe ecosystems. Pp. 15-26 in P. G. Entwistle, A. M. DeBolt, J. H. Kaltenecker, and K. Steenhof (compilers). *Proceedings: sagebrush steppe ecosystems symposium*. Bureau of Land Management Publication No. BLM/ID/PT-001001+1150. Boise, ID.
- WEST, N. E., AND J. A. YOUNG. 2000. Intermountain valleys and lower mountain slopes. Pp. 255-284 in M. G. Barbour, and W. D. Billings (editors). *North American terrestrial vegetation*. Second Edition. Cambridge University Press, Cambridge, UK.
- WESTEMEIER, R. L., AND J. E. BUHNERKEMPE. 1983. Responses of nesting wildlife to prairie grass management in prairie chicken sanctuaries in Illinois. Pp. 36-46 in R. Brewer (editor). *Proceedings of the Eighth North American Prairie Conference*. Western Michigan University, Kalamazoo, MI.
- WESTOBY, M. 1981. Elements of a theory of vegetation dynamics in arid rangelands. *Israel Journal of Botany* 28:169-194.
- WHELAN, R. J. 1995. *The ecology of fire*. Cambridge University Press, Cambridge, UK.
- WHISENANT, S. G. 1990. Changing fire frequencies on Idaho's Snake River plains: ecological and management implications. Pp. 4-10 in E. D. McArthur, E. M. Romney, S. D. Smith, and P. T. Tueller (editors). *Proceedings symposium on cheatgrass invasion, shrub die-off, and other aspects of shrub biology and management*. USDA Forest Service General Technical Report INT-276. USDA Forest Service, Intermountain Research Station, Ogden, UT.
- WHITE, A. S. 1983. The effects of thirteen years of annual prescribed burning on a *Quercus ellipsoidalis* community in Minnesota. *Ecology* 64:1081-1085.
- WHITE, A. S. 1985. Presettlement regeneration patterns in a southwestern ponderosa pine stand. *Ecology* 66:589-594.
- WHITE, D. H., B. R. CHAPMAN, J. H. BRUNJES, IV, R. V. RAFTOVICH, JR., AND J. T. SEGNAK. 1999. Abundance and reproduction of songbirds in burned and unburned pine forests of the Georgia Piedmont. *Journal of Field Ornithology* 70:414-424.
- WHITE, P. S. 1979. Pattern, process, and natural disturbance in vegetation. *Botanical Review* 45:229-299.
- WHITE, P. S., AND S. T. A. PICKETT. 1985. Natural disturbance and patch dynamics: an introduction. Pp. 3-13 in S. T. A. Pickett, and P. S. White (editors). *The ecology of natural disturbance and patch dynamics*. Academic Press, New York, NY.
- WHITE HOUSE. 2002. Healthy forests: an initiative for wildfire prevention and stronger communities. <www.whitehouse.gov/infocus/healthyforests/Healthy_Forests_v2.pdf> (10 June 2004)
- WHITFORD, W. G. 1997. Desertification and animal biodiversity in the desert grasslands of North America. *Journal of Arid Environments* 37:709-720.
- WHITFORD, W. G. 2002. *Ecology of desert systems*. Academic Press, San Diego, CA.
- WHITNEY, G. G. 1994. *From coastal wilderness to fruited plain*. Cambridge University Press, Cambridge, UK.
- WHITTAKER, R. H. 1954. The ecology of serpentine soils. IV. The vegetational response to serpentine soils. *Ecology* 35:275-288.
- WHITTAKER, R. H. 1960. Vegetation of the Siskiyou Mountains, Oregon and California. *Ecological Monographs* 30:279-338.
- WHITTAKER, R. H. 1961. Vegetation history of the Pacific Coast states and the "central" significance of the Klamath region. *Madroño* 16:5-23.
- WHITTAKER, R. H., AND W. A. NIERING. 1964. Vegetation of the Santa Catalina Mountains, Arizona. I. Ecological classification and distribution of species. *Journal of the Arizona Academy of Science* 3:9-34.
- WHITTAKER, R. H., AND W. A. NIERING. 1965. Vegetation of the Santa Catalina Mountains, Arizona. II. A gradient analysis of the south slope. *Ecology* 46:429-452.
- WIENS, J. A. 1973. Pattern and process in grassland bird communities. *Ecological Monographs* 43:237-270.
- WIENS, J. A., AND J. T. ROTENBERRY. 1985. Response of breeding passerine birds to rangeland alteration in a North American shrubsteppe locality. *Journal of Applied Ecology* 22:655-668.
- WIENS, J. A., J. T. ROTENBERRY, AND B. VAN HORNE. 1986. A lesson in the limitations of field experiments: shrubsteppe birds and habitat alterations. *Ecology* 67:365-376.
- WIENS, J. A., J. T. ROTENBERRY, AND B. VAN HORNE. 1987. Habitat occupancy patterns of North American shrubsteppe birds: the effects of spatial scale. *Oikos* 48:132-147.
- WIGLEY, T. B., K. V. MILLER, D. S. DE CALESTA, AND M. W. THOMAS. 2002. Herbicides as an alternative to prescribed burning for achieving wildlife management objectives. Pp. 124-138 in W. Ford, K. R. Russell, and C. E. Moorman (editors). *The role of fire in nongame wildlife management and community restoration: traditional uses and new directions*. USDA Forest Service General Technical Report NE-288. USDA Forest Service, Newtown Square, PA.
- WIKARS, L. O. 2002. Dependence on fire in wood-living insects: an experiment with burned and unburned spruce and birch logs. *Journal of Insect Conservation* 6:1-12.
- WILLIAMS, G. W. 2002. Aboriginal use of fire: are there any "natural" plant communities? Pp. 179-214 in C. E. Kay, and R. T. Simmons (editors). *Wilderness and political ecology*. University of Utah Press, Salt Lake City, UT.
- WILLIAMS, M. 1989. *Americans and their forests. A historical geography*. Cambridge University Press, Cambridge, UK.

- WILLS, R. D., AND J. D. STUART. 1994. Fire history and stand development of a Douglas-fir/hardwood forest in northern California. *Northwest Science* 68:205–212.
- WIMBERLY, M. C., T. A. SPIES, C. J. LONG, AND C. WHITLOCK. 2000. Simulating historical variability in the amount of old forest in the Oregon Coast Range. *Conservation Biology* 14:167–180.
- WINNE, J. C. 1997. History of vegetation and fire on the Pineo Ridge pine grassland barrens of Washington, County, Maine. Pp. 25–52 in P. D. Vickery, and P. W. Dunwiddie (editors). *Grasslands of northeastern North America*. Massachusetts Audubon Society, Lincoln, MA.
- WINTER, M. 1998. Effect of habitat fragmentation grassland-nesting birds in southwestern Missouri. Ph.D. dissertation, University of Missouri, Columbia, MO.
- WINWARD, A. H., AND E. W. TISDALE. 1977. Taxonomy of the *Artemisia tridentata* complex in Idaho. University of Idaho Forest, Wildlife, and Range Experiment Station Bulletin Number 19, Moscow, ID.
- WIRTZ, W. O., II. 1982. Postfire community structure of birds and rodents in southern California chaparral. Pp. 241–246 in C. E. Conrad, and W. C. Oechel (technical coordinators). *Dynamics and management of Mediterranean-type ecosystems*. USDA Forest Service General Technical Report PSW-58. USDA Forest Service, Pacific Southwest Forest and Range Experiment Station, Berkeley, CA.
- WITHGOTT, J. H., AND K. G. SMITH. 1998. Brown-headed Nuthatch. (*Sitta pusilla*). In A. Poole, and F. Gill (editors). *The Birds of North America*, No. 349. The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists' Union, Washington, DC.
- WOOLFENDEN, G. E., AND J. W. FITZPATRICK. 1984. The Florida Scrub Jay. *Monographs in Population Biology* 20. Princeton University Press, Princeton, NJ.
- WOOLFOLK, E. J., AND J. N. REPPERT. 1963. Then and now: changes in California annual-type range vegetation. USDA Forest Service Research Note PSW-N24. USDA Forest Service, Pacific Southwest Research Station, Berkeley, CA.
- WRIGHT, H. A., AND A. W. BAILEY. 1982. *Fire ecology*. John Wiley and Sons, New York, NY.
- YOSEF, R. 1996. Loggerhead Shrike (*Lanius ludovicianus*). In A. Poole, and F. Gill (editors). *The Birds of North America*, No. 231. The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists' Union, Washington, DC.
- YOUNG, J. A. 1994. History and use of semiarid plant communities—changes in vegetation. Pp. 5–8 in S. B. Monsen, and S. G. Kitchen (compilers). *Proceedings—ecology and management of annual rangelands*. USDA Forest Service General Technical Report INT-GTR-313. USDA Forest Service, Intermountain Research Station, Ogden, UT.
- YOUNG, J. A., AND R. A. EVANS. 1973. Downy brome—intruder in the plant succession of big sagebrush communities. *Journal of Range Management* 26: 410–415.
- YOUNG, J. A., AND R. A. EVANS. 1978. Population dynamics after wildfires in sagebrush grasslands. *Journal of Range Management* 31:283–289.
- YOUNG, J. A., AND R. A. EVANS. 1981. Demography and fire history of a western juniper stand. *Journal of Range Management* 34:501–501.
- YOUNG, J. A., AND R. A. EVANS. 1989. Dispersal and germination of big sagebrush (*Artemisia tridentata*) seeds. *Weed Science* 37:201–206.
- YOUNG, J. A., R. E. ECKERT, AND R. A. EVANS. 1979. Historical perspectives regarding the sagebrush ecosystem. Pp. 1–13 in *The sagebrush ecosystem: a symposium*. Utah State University, College of Natural Resources, Logan, UT.
- ZABEL, C. J., J. R. DUNK, H. B. STAUFFER, L. M. ROBERTS, B. S. MULDER, AND A. WRIGHT. 2003. Northern Spotted Owl habitat models for research and management application in California. *Ecological Applications* 13: 1027–1040.
- ZIMMERMAN, J. L. 1984. Nest predation and its relationship to habitat and nest density in Dickcissels. *Condor* 86: 68–72.
- ZIMMERMAN, J. L. 1988. Breeding season habitat selection by the Henslow's Sparrow (*Ammodramus henslowii*) in Kansas. *Wilson Bulletin* 100:17–24.
- ZIMMERMAN, J. L. 1992. Density-independent factors affecting the avian diversity of the tallgrass prairie community. *Wilson Bulletin* 104:85–94.
- ZIMMERMAN, J. L. 1993. *The birds of Konza: the avian ecology of the tallgrass prairie*. University of Kansas Press, Lawrence, KS.
- ZIMMERMAN, J. L. 1997. Avian community responses to fire, grazing, and drought in the tallgrass prairie. Pp. 167–180 in F. Knopf, and F. Sampson (editors). *Ecology and conservation of Great Plains vertebrates*. Springer-Verlag, New York, NY.
- ZINKE, P. J. 1977. The redwood forest and associated north coast forests. Pp. 679–698 in M. G. Barbour, and J. Major (editors). *Terrestrial vegetation of California*. Wiley-Interscience, New York, NY.
- ZOBEL, D. B., A. MCKEE, G. M. HAWK, AND C. T. DYRNESS. 1976. Relationships of environment to composition, structure, and diversity of forest communities of the central western Cascades of Oregon. *Ecological Monographs* 46:135–156.
- ZUCKERBERG, B. 2002. *Grassland restoration on Nantucket Island, MA: effects of burning and mowing on shrubland and grassland songbirds*. M.S. thesis, University of Massachusetts, Amherst, MA.