


2006 Aerial Insect and Disease Survey Silverton, Colorado USGS 100K TOPO!: 37107-E1



1:100,000

Legend

Causal Agent(s) **Not Flown in 2006**

Use of the Number System
 Example: 5-20 = The first number before the dash is the causal agent code. The number after the dash is the number of dead "faded" trees in the polygon or point. When recent dead trees are not counted, an intensity code of 1 (light), 4 (moderate), and 14 (high) may be used after the causal agent code. Periodically, trees per acre estimates are used after the causal agent code instead of number of dead "faded" trees (or an intensity code). For example: 5-12A = The first number before the dash is the causal agent code. The number after the dash is an estimate of the number of dead "faded" trees in the polygon per acre. In this case it would be an estimate that, on the average, one tree per every two acres would be a dead "faded" tree. In another example: 5-2A = that on the average, an estimated three trees per acre are dead "faded" trees. A "/" is used as a separator when a point polygon has more than one causal agent code.

| Code | Causal Agent | Primary Host | Code | Causal Agent | Primary Host | Code | Causal Agent | Primary Host |
|------|--|------------------|------|---------------------------------------|-----------------------|------|--------------------------------------|-------------------|
| 1 | Douglas-fir beetle | Douglas-fir | 107 | Ring-necked Puffin | Lodgepole Pine | 108 | For meadow trapping | Cottonwood/Poplar |
| 2 | Engelmann Spruce Beetle | Engelmann Spruce | 50 | White pine blister rust | 5-Needle Pine | 109 | road salt | Softwoods |
| 3 | Mountain pine beetle | Ponderosa Pine | 51 | Oak mast dieback | Softwoods | 110 | airborne nematode | Scotts Pine |
| 4 | Mountain pine beetle | Lodgepole Pine | 52 | Elytrodemia | Ponderosa Pine | 111 | oak wilt | Oak |
| 5 | Mountain pine beetle | 5-Needle Pine | 53 | Inducible 605, 50 & 65 | All Tree Species | 112 | spine tip | All Tree Species |
| 6 | Western pine beetle | Ponderosa Pine | 54 | Air pollutants | All Tree Species | 113 | headed chestnut borer | Oak |
| 7 | Fir Engiwever | White Fir | 55 | Chemical damage | Softwoods | 114 | anthracnose like foliar disease | Bur Oak |
| 8 | Douglas-fir engraver beetle | Douglas-fir | 56 | Lophodermium pinastri | Softwoods | 115 | Dieback | All Tree Species |
| 9 | Western balsam bark beetle | Subalpine Fir | 57 | Rhabdocline pseudotsugae | Douglas-fir | 116 | Mortality | All Tree Species |
| 10 | Unidentified bark beetle | Softwoods | 58 | Lophodermium acicola | Softwoods | 117 | Discoloration | All Tree Species |
| 11 | Pine engraver | Lodgepole Pine | 59 | Lachnospila acicola | Softwoods | 118 | Herbicide | All Tree Species |
| 12 | Pine engraver | Ponderosa Pine | 60 | Lophodermium concolor | Softwoods | 119 | Quaking Aspen | Quaking Aspen |
| 13 | Pine engraver | Lodgepole Pine | 61 | Ostrya sp. | Softwoods | 120 | Marssonina blight | Quaking Aspen |
| 14 | Pine engraver | Ponderosa Pine | 62 | Needle cast (Hypodermataceae) | All Tree Species | 200 | Dieback (ash) | Ash |
| 15 | Ponderosa pine needle miner | Ponderosa Pine | 63 | Root Rot | All Tree Species | 201 | Dieback (hardwood) | Hardwoods |
| 16 | Ponderosa pine needle miner | Jack Pine | 64 | Unidentified disease | Softwoods | 202 | Dieback (oak) | Oak |
| 17 | Jack pine budworm | Douglas-fir | 65 | Winter damage light | All Tree Species | 203 | Mortality (old cottonwood) | Cottonwood/Poplar |
| 18 | Spruce budworm, light defol. | Douglas-fir | 66 | Winter damage medium | All Tree Species | 210 | Mortality (eastern cedar) | Eastern Red Cedar |
| 19 | Spruce budworm, medium defol. | Douglas-fir | 67 | Winter damage heavy | All Tree Species | 211 | Mortality (oak) | Oak |
| 20 | Spruce budworm, heavy defol. | Douglas-fir | 68 | Opilidia | Softwoods | 212 | Mortality (spruce) | Spruce |
| 21 | Douglas-fir tussock moth | Douglas-fir | 69 | Prionyn black stain | Common Prinyon | 220 | Discoloration (ash) | Ash |
| 22 | Pine butterfly | Ponderosa Pine | 70 | Fire | All Tree Species | 221 | Discoloration (conifer) | Softwoods |
| 23 | Pine sawfly | Ponderosa Pine | 71 | Fire | Softwoods | 222 | Discoloration (cottonwood) | Cottonwood/Poplar |
| 24 | Tree caterpillars | Hardwoods | 72 | Windthrow | All Tree Species | 223 | Discoloration (eastern cedar) | Eastern Red Cedar |
| 25 | Leaf beetles | Hardwoods | 73 | Windthrow | All Tree Species | 224 | Discoloration (hardwood) | Hardwoods |
| 26 | Oak leaf roller | Hardwoods | 74 | High water damage | All Tree Species | 225 | Discoloration (oak) | Oak |
| 27 | Pine needle-sheath miner | Ponderosa Pine | 75 | Avalanche | All Tree Species | 226 | Discoloration (spruce) | Spruce |
| 28 | Pine sawfly | Ponderosa Pine | 76 | Avian dieback-multiple agents | Quaking Aspen | 230 | Herbicide (eastern cedar) | Eastern Red Cedar |
| 29 | Pine tussock moth | Ponderosa Pine | 77 | Prionyn mortality | Common Prinyon | 231 | Flagging (hardwood) | Hardwoods |
| 30 | Variable oak leaf defoliation | Hardwoods | 78 | Juniper mortality-unknown agents | Juniper | 232 | Discoloration (eastern cedar) | Eastern Red Cedar |
| 31 | Carlinworms | Hardwoods | 79 | Juniper mortality-unknown agents | Gambel Oak | 233 | Discoloration (spruce) | Spruce |
| 32 | Unidentified defoliator | All Tree Species | 80 | Limber pine dieback-multiple agents | Limber Pine | 234 | Unidentified defoliator (cottonwood) | Cottonwood/Poplar |
| 33 | Variable oak leaf defoliation | Hardwoods | 81 | Limber pine dieback-multiple agents | Limber Pine | 235 | Unidentified defoliator (oak) | Oak |
| 34 | Unidentified defoliator | All Tree Species | 82 | Hail damage | All Tree Species | 236 | Unidentified defoliator (hardwood) | Hardwoods |
| 35 | Heliconiastrum annosum (Fomes annosus) | Softwoods | 83 | Hail damage | All Tree Species | 237 | Mortality (pine) | Pine |
| 36 | Armillaria ostoyae (Armillaria mellea) | Softwoods | 84 | Unknown polygon | Unknown | | | |
| 37 | Porygonus schweinitzi | Softwoods | 100 | old prison mortality | Common Prinyon | | | |
| 38 | Phonopis | Softwoods | 101 | dead tall top | Lodgepole Pine | | | |
| 39 | Cytospora | All Tree Species | 102 | slush elm disease | Elm | | | |
| 40 | Western gall rust | Unknown | 103 | dieback blight | Ponderosa Pine | | | |
| 41 | Comandra rust | Unknown | 104 | lps. nutlets | Spruce, White Spruce | | | |
| 42 | Stactiform rust | Lodgepole Pine | 105 | straght killed narrow leaf cottonwood | Narrowleaf Cottonwood | | | |

USGS 100K Quad - Location Map



How Aerial Surveys Are Conducted

Data represented on this map are based on aerial observations manually recorded onto a map. This procedure is considered both an art form and a form of scientific data collection, and is highly subjective. An observer only has a few seconds to recognize the color difference between healthy and damaged trees of different species; diagnose causal agents correctly; estimate intensity; delineate the extent of damage; and precisely record this information on a georeferenced map. Air turbulence, cloud shadows, distance from aircraft, haze, smoke, and observer experience can all affect the quality of the survey. These data summaries provide an estimate of conditions on the ground and may differ from estimates derived by other methods.

Aerial surveys provide information on the current status for many causal agents, and are important when examining insect activity trends by comparing historical and current survey data over large areas.

Overview surveys are a snap shot in time and therefore may not be timed to accurately capture the true extent or severity of a particular disturbance activity. Aerial surveys can be thought of as the first stage in a multi-stage sampling design. Other remote sensing approaches, including aerial photography, electro-optical sensors, and specially designed aerial surveys with modified flight patterns, can be used to more accurately delineate the extent and severity of a particular disturbance agent. The preceding methods are often more costly than overview surveys, and are generally reserved to address situations of sufficient environmental, economic, or political importance.

Area surveyed by Erik Johnson & Pat Ahern 09/13 - 09/14 2006;
 Erik Johnson, Patrick Ahern, Kelly Rogers 09/11 - 09/13 2006;
 Al Dymerski 09/12-09/21 2006 Map Created: 01/04/2006
 Projection: UTM NAD83 Zone 13
 Author: J. Ross, USDA Forest Service

DIRECT ALL INQUIRIES TO:



Colorado State Forest Service
 Colorado State University
 Fort Collins, Colorado 80523

USDA Forest Service, Region 2
 Renewable Resources
 Forest Health Management
 PO Box 25127
 Lakewood, Colorado 80225

****DISCLAIMER****
 Due to the nature of aerial surveys, the data on this map will only provide rough estimates of location, intensity and the resulting trend information for agents detectable from the air. Many of the most destructive diseases are not represented on this map because these agents are not detectable from aerial surveys. The data presented on this map should only be used as a partial indicator of insect and disease activity, and should be validated on the ground for actual location and causal agent. Shaded areas show locations where tree mortality or defoliation were apparent from the air. Intensity of damage is variable and not all trees in shaded areas are dead or defoliated.

The insect and disease data represented on this map are available digitally from the USDA Forest Service, Region Two Forest Health Management group. The cooperators reserve the right to correct, update, modify or replace GIS products. Using this map for purposes other than those for which it was intended may yield inaccurate or misleading results.

A data dictionary and digital copies of this map and the insect and disease data are available at: <http://www.fs.fed.us/r2/resources/fhm/aerialsurvey/>