



# 2006 Aerial Insect and Disease Survey Fort Collins, Colorado USGS 100K TOPO!: 40105-E1



1:100,000

Legend		Causal Agent(s)	Not Flown in 2006		
<p>Use of the Number System</p> <p>Example: 5-25 = The first number before the dash is the causal agent code. The number after the dash is the number of dead "fader" trees in the polygon or point. When recent dead trees are not counted, an intensity code of L-light, M-moderate, and H-high may be used after the causal agent code. Periodically, trees per acreage estimates are used after the causal agent code instead of number of dead "fader" trees (or an intensity code). For example: 5-1/2A = The first number before the dash is the causal agent code. The number after the dash is an estimation of the number of dead "fader" trees in the polygon per acre. In this case it would be an estimation that, on the average, one tree per every two acres would be a dead "fader" tree. In another example: 5-3A = that on the average, an estimated three trees per acre are dead "fader" trees. A "-" is used as a separator when a point/polygon has more than one causal agent code.</p>					
Code	Causal Agent	Primary Host	Code	Causal Agent	Primary Host
1	Douglas-fir beetle	Douglas-fir	42	Aspen	Lodgepole Pine
2	Engelmann Spruce Beetle	Engelmann Spruce	50	White pine blister rust	Cottontail Poplar
3	Mountain pine beetle	Ponderosa Pine	51	Deaf mistletoe	Cottontail Poplar
4	Mountain pine beetle	Lodgepole Pine	52	Elytrodema	Softwoods
5	Mountain pine beetle	Sitka Spruce	53	Includes #50, 51 & 52	All Tree Species
6	Western pine beetle	Ponderosa Pine	54	Aspen bark beetle	All Tree Species
7	Fire Engulfer	White Fir	55	Chemical damage	All Tree Species
8	Fire Engulfer	Subalpine Fir	56	Lophodermium pinis	Softwoods
9	Douglas-fir engraver beetle	Douglas-fir	57	Rhabdocline pseudotsugae	Douglas-fir
10	Western balsam bark beetle	Subalpine Fir	58	Lophodermium pinis	Softwoods
11	Unidentified bark beetle	Lodgepole Pine	59	Lecanosticella aculeata	Softwoods
12	Pine engraver	Ponderosa Pine	60	Lophodermium concolor	Softwoods
13	Pine engraver	Lodgepole Pine	61	Dobsonia sp.	Softwoods
14	Pine engraver	Ponderosa Pine	62	Needle cast (Hypodermataceae)	Softwoods
15	Ponderosa pine needle miner	Lodgepole Pine	64	Unidentified disease	Softwoods
16	Lodgepole pine needle miner	Ponderosa Pine	65	Winter damage light	All Tree Species
17	Jack pine budworm	Jack Pine	66	Winter damage medium	All Tree Species
18	Spruce budworm, light defol.	Douglas-fir	67	Winter damage heavy	All Tree Species
19	Spruce budworm, heavy defol.	Douglas-fir	68	Diptera	Softwoods
20	Douglas-fir tussock moth	Ponderosa Pine	69	Pinus black stain	Common Pinus
21	Pine looper	Ponderosa Pine	70	Mortality (old cottonwood)	Cottontail Poplar
22	Pine looper	Ponderosa Pine	71	Mortality (western cedar)	Eastern Red Cedar
23	Text caterpillars	Hardwoods	72	Mortality (oak)	Oak
24	Leaf beetles	Hardwoods	73	Mortality (spruce)	Spruce
25	Oak leaf roller	Hardwoods	74	Unidentified (oak)	Ash
26	Pine needle-needle miner	Ponderosa Pine	75	Aspen decline-multiple agent(s)	Softwoods
27	Pine sawflies	Ponderosa Pine	76	Pinus pine mortality	Common Pinus
28	Pine tussock moth	Ponderosa Pine	77	Unidentified (oak)	Cottontail Poplar
29	Carleworms	Hardwoods	78	Unidentified (eastern cedar)	Eastern Red Cedar
30	Variable oak leaf caterpillar	Hardwoods	79	Unidentified (hardwood)	Hardwoods
31	Unidentified defoliator	All Tree Species	80	Unidentified (oak)	Oak
32	Unidentified defoliator	All Tree Species	81	Unidentified (spruce)	Spruce
33	Unidentified defoliator (Fomes annosus)	Softwoods	82	Herbicide (cottonwood)	Cottontail Poplar
34	Armillaria ostoyae (Armillaria mellea)	Softwoods	83	Herbicide (eastern cedar)	Eastern Red Cedar
35	Polygonia schweinitzi	Softwoods	84	Unidentified (hardwood)	Hardwoods
36	Phragmites	Softwoods	85	Unidentified (oak)	Oak
37	Cytospora	All Tree Species	86	Unidentified (oak)	Oak
38	Western gall rust	Unknown	87	Unidentified (oak)	Oak
39	Comandra gall rust	Unknown	88	Unidentified (oak)	Oak
40	Stemflow rust	Lodgepole Pine	89	Unidentified (oak)	Oak
41	Stemflow rust	Lodgepole Pine	90	Unidentified (oak)	Oak
42	Stemflow rust	Lodgepole Pine	91	Unidentified (oak)	Oak
43	Stemflow rust	Lodgepole Pine	92	Unidentified (oak)	Oak
44	Stemflow rust	Lodgepole Pine	93	Unidentified (oak)	Oak
45	Stemflow rust	Lodgepole Pine	94	Unidentified (oak)	Oak
46	Stemflow rust	Lodgepole Pine	95	Unidentified (oak)	Oak
47	Stemflow rust	Lodgepole Pine	96	Unidentified (oak)	Oak
48	Stemflow rust	Lodgepole Pine	97	Unidentified (oak)	Oak
49	Stemflow rust	Lodgepole Pine	98	Unidentified (oak)	Oak
50	Stemflow rust	Lodgepole Pine	99	Unidentified (oak)	Oak
51	Stemflow rust	Lodgepole Pine	100	Unidentified (oak)	Oak
52	Stemflow rust	Lodgepole Pine	101	Unidentified (oak)	Oak
53	Stemflow rust	Lodgepole Pine	102	Unidentified (oak)	Oak
54	Stemflow rust	Lodgepole Pine	103	Unidentified (oak)	Oak
55	Stemflow rust	Lodgepole Pine	104	Unidentified (oak)	Oak
56	Stemflow rust	Lodgepole Pine	105	Unidentified (oak)	Oak



**How Aerial Surveys Are Conducted**

Data represented on this map are based on aerial observations manually recorded onto a map. This procedure is considered both an art form and a form of scientific data collection, and is highly subjective. An observer only has a few seconds to recognize the color difference between healthy and damaged trees of different species; diagnose causal agents correctly; estimate intensity; delineate the extent of damage; and precisely record this information on a georeferenced map. Air turbulence, cloud shadows, distance from aircraft, haze, smoke, and observer experience can all affect the quality of the survey. These data summaries provide an estimate of conditions on the ground and may differ from estimates derived by other methods.

Aerial surveys provide information on the current status for many causal agents, and are important when examining insect activity trends by comparing historical and current survey data over large areas.

Overview surveys are a snap shot in time and therefore may not be timed to accurately capture the true extent or severity of a particular disturbance activity. Aerial surveys can be thought of as the first stage in a multi-stage sampling design. Other remote sensing approaches, including aerial photography, electro-optical sensors, and specially designed aerial surveys with modified flight patterns, can be used to more accurately delineate the extent and severity of a particular disturbance agent. The preceding methods are often more costly than overview surveys, and are generally reserved to address situations of sufficient environmental, economic, or political importance.

**Area surveyed by William Ciesla, Meg Halford, Ingrid Aguayo**  
07/10 - 07/14 2006  
Map Created: 01/03/2007  
Projection: UTM NAD83 Zone 13  
Author: J. Ross, USDA Forest Service

**DIRECT ALL INQUIRIES TO:**

**Colorado State Forest Service**  
Colorado State University  
Fort Collins, Colorado 80523

**USDA Forest Service, Region 2**  
Renewable Resources  
Forest Health Management  
PO Box 25127  
Lakewood, Colorado 80225

Due to the nature of aerial surveys, the data on this map will only provide rough estimates of location, intensity and the resulting trend information for agents detectable from the air. Many of the most destructive diseases are not represented on this map because these agents are not detectable from aerial surveys. The data presented on this map should only be used as a partial indicator of insect and disease activity, and should be validated on the ground for actual location and causal agent. Shaded areas show locations where tree mortality or defoliation were apparent from the air. Intensity of damage is variable and not all trees in shaded areas are dead or defoliated.

\*\*\*\*\*DISCLAIMER\*\*\*\*\*

The insect and disease data represented on this map are available digitally from the USDA Forest Service, Region Two Forest Health Management group. The cooperators reserve the right to correct, update, modify or replace GIS products. Using this map for purposes other than those for which it was intended may yield inaccurate or misleading results.

A data dictionary and digital copies of this map and the insect and disease data are available at: <http://www.fs.fed.us/r2/resources/fhm/aerialsurvey/>