


# 2007 Aerial Insect and Disease Survey Farson, Wyoming USGS 100K TOPO!: 42109-A1




1:100,000

### Legend

**Causal Agent(s)** (Red box) **Not Flown** (Grey box)

Use of the Number System  
 Example: 5-25 = The first number before the dash is the causal agent code. The number after the dash is the number of dead "fader" trees in the polygon or point. When recent dead trees are not counted, an intensity code of L-light, M-moderate, and H-high may be used after the causal agent code. Periodically, trees per acreage estimates are used after the causal agent code instead of number of dead "fader" trees (or an intensity code). For example: 5-12(A) = The first number before the dash is the causal agent code. The number after the dash is an estimation of the number of dead "fader" trees in the polygon per acre. In this case it would be an estimation that, on the average, one tree per every two acres would be a dead "fader" tree. In another example: 5-3A = that on the average, an estimated three trees per acre are dead "fader" trees. A // is used as a separator when a point polygon has more than one causal agent code.

Code	Causal Agent	Primary Host	Code	Causal Agent	Primary Host
1	Douglas-fir beetle	Douglas-fir	49	Atropella	Lodgepole Pine
2	Engelmann Spruce Beetle	Engelmann Spruce	50	White pine blister rust	5-Needle Pine
5	Mountain pine beetle	Ponderosa Pine	61	Dwarf mistletoe	Softwoods
6	Mountain pine beetle	Lodgepole Pine	62	Rhinodactylus	Ponderosa Pine
7	Mountain pine beetle	5-Needle Pine	63	Incluses #65, 66 & 68	All Tree Species
8	Western pine beetle	Ponderosa Pine	64	Air pollution	All Tree Species
9	Fire Engraver	White Fir	65	Chemical damage	All Tree Species
10	Douglas-fir engraver beetle	Douglas-fir	66	Lophodermium pinasti	Softwoods
11	Western balsam bark beetle	Subalpine Fir	67	Rhabdocline pseudotsugae	Douglas-fir
12	Unidentified bark beetle	Softwoods	68	Lophodermella arcuata	Softwoods
13	Pine engraver	Lodgepole Pine	69	Larobotrya acicola	Softwoods
14	Pine engraver	Ponderosa Pine	70	Lophodermella concolor	Softwoods
15	Ponderosa pine needle miner	Lodgepole Pine	71	Dobsonia par	Softwoods
16	Lodgepole pine needle miner	Ponderosa Pine	72	Needle cast (Hypodermataceae)	Softwoods
17	Jack pine budworm	Jack Pine	73	Root Rot	All Tree Species
18	Spruce budworm, light defol.	Douglas-fir	74	Unidentified disease	Softwoods
19	Spruce budworm, medium defol.	Douglas-fir	75	Winter damage light	All Tree Species
20	Spruce budworm, heavy defol.	Douglas-fir	76	Winter damage medium	All Tree Species
21	Douglas-fir spruce moth	Douglas-fir	77	Winter damage heavy	All Tree Species
22	Pine butterfly	Ponderosa Pine	78	Diploids	Softwoods
23	Pine looper	Ponderosa Pine	79	Pinus bark stain	Common Pinion
24	Pine tortrix	Ponderosa Pine	80	Hail damage	All Tree Species
25	Teak caterpillars	Hardwoods	81	Unknown polygon	Common Pinion
26	Leaf beetles	Hardwoods	82	Unknown polygon	Common Pinion
27	Oak leaf roller	Hardwoods	83	High water damage	All Tree Species
28	Pine needle-shaft miner	Ponderosa Pine	84	Aniselmis	All Tree Species
29	Pine hawkmoth	Ponderosa Pine	75	Aspen decline-multiple agents	Quaking Aspen
30	Pine hawkmoth	Ponderosa Pine	76	Pinus pine mortality	Common Pinion
31	Cankworms	Hardwoods	77	Juniper mortality-unknown agents	Juniper
32	Variegated oak leaf caterpillar	Hardwoods	78	Quaking Oak	Quaking Oak
33	Armillaria ostroyae (Armillaria mellea)	Softwoods	79	Lumber Pine	Lumber Pine
34	Unidentified defoliator	All Tree Species	80	Hail damage	All Tree Species
35	Heterobasidion annosum (Fomes annosus)	Softwoods	81	Unknown polygon	Common Pinion
36	Phomopsis	Softwoods	82	Unknown polygon	Common Pinion
37	Polyonyx schweinitzi	Softwoods	100	old pinion mortality	Lodgepole Pine
38	Phomopsis	Softwoods	101	dead salt top	Lodgepole Pine
39	Cytospora	All Tree Species	102	slutch elm disease	Elm
40	Western gall rust	Unknown	103	alibaba blight	Ponderosa Pine
41	Comandra rust	Unknown	104	low numbers	Spruce, White Spruce
42	Stactoflora rust	Lodgepole Pine	105	drought killed narrow leaf cottonwood	Narrowleaf Cottonwood
43			106	fox squirrel ragging	Cottonwood/Poplar
44			107	fall webworm	Cottonwood/Poplar
45			108	road salt	Softwoods
46			109	pinewood nematode	Scotch Pine
47			110	oak wilt	Oak
48			111	storage disease	White Spruce
			112	spring ice	Oak
			113	twined chestnut borer	Oak
			114	anthracose like foliar disease	Bur Oak
			115	Dieback	All Tree Species
			116	Flagging	All Tree Species
			117	Discoloration	All Tree Species
			118	Heterobasidion	All Tree Species
			119	Quaking Aspen	Quaking Aspen
			120	aspen tortrix	Quaking Aspen
			121	Manisoma blight	Quaking Aspen
			200	Dieback (ash)	Ash
			201	Dieback (cottonwood)	Cottonwood/Poplar
			202	Dieback (hardwood)	Hardwoods
			203	Dieback (oak)	Oak
			204	Mortality (oak cottonwood)	Oak
			211	Mortality (eastern cedar)	Eastern Red Cedar
			212	Mortality (hardwood)	Hardwoods
			213	Mortality (oak)	Oak
			214	Mortality (spruce)	Spruce
			220	Discoloration (ash)	Ash
			221	Discoloration (hardwood)	Hardwoods
			222	Discoloration (cottonwood)	Cottonwood/Poplar
			223	Discoloration (eastern cedar)	Eastern Red Cedar
			224	Discoloration (hardwood)	Hardwoods
			225	Discoloration (oak)	Oak
			226	Discoloration (spruce)	Spruce
			230	Heterobasidion (cottonwood)	Cottonwood/Poplar
			231	Heterobasidion (eastern cedar)	Eastern Red Cedar
			240	Flagging (hardwood)	Hardwoods
			250	Unidentified defoliator (cottonwood)	Cottonwood/Poplar
			251	Unidentified defoliator (elm)	Elm
			252	Unidentified defoliator (hardwood)	Hardwoods
			300	Mortality (pine)	Pine



### How Aerial Surveys Are Conducted

Data represented on this map are based on aerial observations manually recorded onto a map. This procedure is considered both an art form and a form of scientific data collection, and is highly subjective. An observer only has a few seconds to recognize the color difference between healthy and damaged trees of different species; diagnose causal agents correctly; estimate intensity; delineate the extent of damage; and precisely record this information on a georeferenced map. Air turbulence, cloud shadows, distance from aircraft, haze, smoke, and observer experience can all affect the quality of the survey. These data summaries provide an estimate of conditions on the ground and may differ from estimates derived by other methods.

Aerial surveys provide information on the current status for many causal agents, and are important when examining insect activity trends by comparing historical and current survey data over large areas.

Overview surveys are a snap shot in time and therefore may not be timed to accurately capture the true extent or severity of a particular disturbance activity. Aerial surveys can be thought of as the first stage in a multi-stage sampling design. Other remote sensing approaches, including aerial photography, electro-optical sensors, and specially designed aerial surveys with modified flight patterns, can be used to more accurately delineate the extent and severity of a particular disturbance agent. The preceding methods are often more costly than overview surveys, and are generally reserved to address situations of sufficient environmental, economic, or political importance.

**Area surveyed by AI Dymerski  
Map Created: 12/30/2007  
Projection: UTM NAD83 Zone 13  
Author: J. Ross, USDA Forest Service**

### DIRECT ALL INQUIRIES TO:

Wyoming State Forestry Division  
 1100 West 22nd Street  
 Cheyenne, Wyoming 82002

USDA Forest Service, Region 2  
 Renewable Resources  
 Forest Health Management  
 P.O. Box 25127  
 Lakewood, Colorado 80225

\*\*\*\*DISCLAIMER\*\*\*\*  
 Due to the nature of aerial surveys, the data on this map will only provide rough estimates of location, intensity and the resulting trend information for agents detectable from the air. Many of the most destructive diseases are not represented on this map because these agents are not detectable from aerial surveys. The data presented on this map should only be used as a partial indicator of insect and disease activity, and should be validated on the ground for actual location and causal agent. Shaded areas show locations where tree mortality or defoliation were apparent from the air. Intensity of damage is variable and not all trees in shaded areas are dead or defoliated.

The insect and disease data represented on this map are available digitally from the USDA Forest Service, Region Two Forest Health Management group. The cooperators reserve the right to correct, update, modify or replace GIS products. Using this map for purposes other than those for which it was intended may yield inaccurate or misleading results.

A data dictionary and digital copies of this map and the insect and disease data are available at: <http://www.fs.fed.us/r2/resources/fhm/aerialsurvey/>