


2006 Aerial Insect and Disease Survey

USGS 100K Quad: Shelton - A147123; 2D



Defoliators		Mortality Agents	
Code	Damaging Agent	Code	Damaging Agent
AS	Spruce aphid	1	Douglas-fir beetle
BB	Western blackheaded budworm	2	Douglas-fir engraver
BM	Modic budworm	3	Spruce beetle
BP	Sugar pine tortrix	4	Fir engraver
BS	Western spruce budworm	5	Western balsam bark beetle
BY	Burns' light/colobormella	6B	Mountain pine beetle
CH	Larch	6C	Mountain pine beetle
CL	Western hemlock looper	6K	Mountain pine beetle
LG	Green striped forest looper	6P	Mountain pine beetle
LL	Larch looper	6W	Mountain pine beetle
LZ	Black pine needle scale	6W	Mountain pine beetle
MD	Douglas-fir budmoth	8	Western pine beetle
ML	Larch budmoth	9	Western pine beetle
MN	Douglas-fir needle midge	9	Western pine beetle
MS	Spruce budmoth	9	Western pine beetle
ND	Needle miner	BEAR	Bear damage
NJ	Needle miner	LW	Flatheaded wood borer
NK	Needle miner	PL	Black stain root disease
NL	Needle miner	RD	Pink Cedars root disease
NP	Needle miner	WATR	Water damage
NT	Needle miner		
NW	Needle miner		
OB	Western oak looper		
PH	Pine butterfly		
PI	Pine needle cast		
PB	Phantom hemlock looper		
PM	Panorama moth		
PN	Pine needlehead miner		
PS	Pine needle scale		
RC	Needle cast		
S	Spider mite		
SA	Sawfly		
SD	Sawfly		
SH	Sawfly		
SL	Sawfly		
SM	Satin moth		
SNC	Swiss needle cast		
SP	Sawfly		
SW	Sawfly		
TA	Tent caterpillar, alder		
TC	Tent caterpillar, other		
TM	Douglas-fir tussock moth		
TS	Ashpen		

USGS 100K Quad: Shelton - A147123; 2D
2006 Aerial Insect and Disease Detection Survey
Mapscale: 1:100,000
Date: November 27, 2006

Legend

- Defoliating Agents
- Mortality Agents
- Other Damage
- WaDNR Managed Lands

2006 Large Fires
 Source: Northwest Coordination Center

Other Damaging Agents

Code	Damaging Agent	Primary Host
AB	Balsam woolly adelgid	True fir
AC	Cooley spruce gall adelgid	Spruce, Douglas-fir
AM	Leaf discoloration	Maple
BR	Blester nut	Five-needle pine
CC	Cytospora canker	True fir
DH	Dying hemlock	Hemlock
FIRE	Fir	All species
GP	Gouty pitch midge	Ponderosa pine
HA	Hail	All species
HD	Heartwood decline	Hardwoods
NF	Areas not flown	
CLT	No damage detected	
PMD	Pacific madrone decline	Pacific madrone
PR	Leaf rust in poplars	Poplars
RD	Red bait	All species
SLD	Slide	All species
UNKD	Unknown defoliation	All species
UNKM	Unknown mortality	All species
WATR	Water damage	All species
WIND	Windthrow	All species
WNTR	Winter Damage	All species

The cause of damage is described by a symbol listed above and is followed by: number of trees affected; number of tree/care (example: 5A) or intensity of damage (L-Light, M-Moderate, H-Heavy).

The map base was created with TOPOI (Copyright 2001, National Geographic); available online at: www.ngmapstore.com

A data dictionary, digital copies of this map and ArcGIS insect and disease data are available at: www.fs.fed.us/r6/nr/fid/data.shtml

How the Aerial Surveys Are Conducted

Data represented on this map are based on trees visibly affected by forest insects and diseases detected and recorded during aerial survey flights conducted by the USDA Forest Service and the Washington Department of Natural Resources. Observers have just a few seconds to recognize the color difference between healthy and damaged trees of different species; diagnose causal agents correctly; estimate intensity; delineate the extent of damage; and precisely record this information on a georeferenced, digital map. Air turbulence, cloud shadows, distance from aircraft, haze, smoke and observer experience can all affect the quality of the survey. These data summaries provide an estimate of conditions on the ground and may differ from estimates derived by other methods.

The aerial survey provides information on the current status for many causal agents, and is important when examining insect activity trends by comparing historical and current survey data over large areas.

Overview surveys are a 'snap shot' in time and therefore may not be timed to accurately capture the true extent or severity of a particular disturbance activity. Specially designed surveys with modified flight patterns and timing may be conducted to more accurately delineate the extent and severity of a particular disturbance agent. Special surveys, such as Swiss needle cast surveys, are conducted when resources are available to address situations of sufficient economic, political or environmental importance.

DIRECT ALL INQUIRIES TO:

Washington State Department of Natural Resources
 Resource Protection
 Forest Health
 1111 Washington St. SE
 Olympia, WA 98504

-- OR --

USDA Forest Service, Region 6
 Natural Resources
 Forest Health Protection
 PO Box 3623
 Portland, Oregon 97208

DISCLAIMER
 The insect and disease data presented should only be used as an indicator of insect and disease activity, and should be ground-checked for precise location, extent, severity and causal agent. Color coded polygons show locations where trees were recently killed or defoliated. Intensity of damage is variable and not all trees within coded polygons are dead or defoliated. The cooperators reserve the right to correct, update, modify or replace GIS products without notice. Using this map for purposes other than those for which it was intended may yield inaccurate or misleading results.