

# Nitrogen Mineralization in Aspen/Conifer Soils After a Natural Fire

Michael C. Amacher, Dale L. Bartos, Tracy Christopherson, Amber D. Johnson, and Debra E. Kutterer<sup>1</sup>

**Abstract**—We measured the effects of the 1996 Pole Creek fire, Fishlake National Forest, Utah, on available soil N and net N mineralization for three summers after the fire using an ion exchange membrane (IEM) soil core incubation method. Fire in mixed aspen/conifer increased the amount of available  $\text{NH}_4$ , and a subsequent net increase in soil nitrification was observed. Release of  $\text{NH}_4$  from the ash and a net increase in soil nitrification were still observed 3 years after the fire. The fire-caused increase in available soil N provides a source of N for aspen regrowth.

## Introduction

Aspen (*Populus tremuloides*) provide many ecological benefits to resource users, including protection of watersheds from erosion, some protection against rapid wildfire advance, increased biological diversity in the species-rich grass-forb understory, wood fiber, wildlife habitat, forage for domestic livestock and native ungulates, recreational sites, aesthetic considerations (e.g., fall leaf colors), and more water yield than conifers (Bartos and Campbell 1998a). Some authors have suggested that aspen are in decline throughout the Interior Western United States (Bartos and Campbell 1998a), although the extent of decline has been the subject of much debate. National Forest inventory and analysis data indicate that about 60% of aspen stands have disappeared since the mid-19<sup>th</sup> century (Bartos and Campbell 1998b). Decline of aspen stands is attributed to natural succession (e.g., invasion of conifers) (Harniss 1981), fire suppression (Jones and Debye 1985), and overbrowsing by domestic livestock and native ungulates (Kay 1990).

Western aspen, which reproduce primarily by suckering from lateral roots, often need disturbances to stimulate the suckering response (Schier 1981). In the past, these disturbances included lightning-caused fires that killed mature aspen, and insects and diseases that weakened old decadent stands before toppling by wind. Suppression of human-caused and lightning-caused fires throughout the twentieth century has greatly reduced one of the main forces of aspen regeneration from the landscape. Even when a disturbance such as fire or wind does stimulate aspen regeneration, overbrowsing of new aspen suckers by domestic livestock and native ungulates quickly suppresses regeneration. Evidence indicates that overbrowsing and conifer invasion of declining aspen stands have led to the precarious state of aspen in the Interior Western United States, although this assessment is not universally accepted.

In addition to stimulating aspen suckering, fire can also exert a large influence on soil properties in burned stands, depending on the intensity of the burn. Increases in plant available nutrients usually occur following fire because of the transfer of nutrients from the ash to the soil (Schlesinger 1997). These

---

<sup>1</sup>Rocky Mountain Research Station, USDA Forest Service, Logan, UT.

increases are usually considered to be short lived because of nutrient uptake by new vegetative growth and losses from leaching (primarily  $\text{NO}_3$ ) and erosion.

During June and July of 1996, an 8,000-acre natural fire (the Pole Creek fire) occurred in the area of Betenson and Grindstone Flats on the Beaver Ranger District of the Fishlake National Forest, Utah. This moderate-intensity fire burned in mixed stands of older aspen and invading conifers, primarily sub-alpine fir (*Abies lasiocarpa*) and Engelmann spruce (*Picea engelmannii*). After the fire, some cutting of fire-killed aspen and conifers was done to thin the stands, and four exclosures were constructed by Utah Division of Wildlife Resources and Fishlake National Forest personnel to protect aspen suckers from browsing by domestic livestock and native ungulates. Two exclosures were built in the Grindstone Flat area and two in the nearby Betenson Flat area.

We present results of the effects of a natural fire on available soil N and net soil N mineralization during the first, second, and third summers after the fire year. In particular, we studied the effects of fire on the amounts of extractable soil  $\text{NH}_4$  and  $\text{NO}_3$ , amounts and rates of leaching of  $\text{NH}_4$  and  $\text{NO}_3$  from plant litter or ash layers, and net changes in some reactions of the internal soil N cycle (e.g., nitrification, mineralization).


## Methods

We measured net soil N mineralization/immobilization during the summer months in 1997, 1998, and 1999, following the 1996 Pole Creek fire using an ion exchange membrane (IEM) soil core incubation technique. The method was similar to that developed by DiStefano and Gholz (1986) except that we used ion exchange membranes (Ionics Inc., Watertown, MA) in place of ion exchange resin bags. The IEMs have identical ion exchange properties to the more familiar resin beads and have shown considerable promise for assessing soil nutrient availability (Cooperband and Logan 1994).

### ***Ion Exchange Membrane Soil Core Method***

The assembly of the soil cores with IEMs is shown in figure 1. The soil core is enclosed in a 2-inch diameter x 4-inch long plastic liner (Forestry Suppliers, Jackson, MS). End caps were used to hold the IEMs against each end of the soil core. Circular holes (1.5-inch diameter) were cut in the center of the 2-inch diameter end caps. Cation (CEM) and anion (AEM) exchange membranes were prepared by cutting 2-inch diameter circles from the original membrane sheets and drilling nine evenly spaced  $\frac{1}{8}$ -inch holes in each circle to allow movement of air and slow percolation of water through the soil cores. The CEMs and AEMs were soaked five times in 1 M HCl and 1 M  $\text{NaHCO}_3$ , respectively, rinsed with deionized water, and sealed in zip-lock plastic bags for transport to the field. To protect the IEMs from abrasion and excessive desiccation and to direct infiltrating water to flow across the surfaces of the IEMs to increase the efficiency of ion exchange, porous screens were prepared by cutting 2-inch diameter circles from 114-um Spectra/Mesh polyethylene filter screens.

We installed the IEM soil cores in the following manner. Leaf litter (unburned areas) or ash layer (burned areas) was first removed from the soil surface and set aside. The soil coring head with a plastic liner was driven into the soil using a slide hammer attachment (Forestry Suppliers, Jackson, MS) until flush with the top of the soil. The intact soil core was removed from the coring



**Figure 1**—Ion exchange membrane soil core assembly.

head, the ends of the core were trimmed, and the excess soil was returned to the soil hole. The end caps, plastic screens, and IEMs were assembled as shown in figure 1, the IEM soil core was placed in the hole with the top end up, and the core was covered with leaf litter or ash. Core locations were marked with flagged rebar stakes. An additional soil core was taken adjacent to the installed IEM soil core to determine initial soil exchangeable  $\text{NH}_4$  and  $\text{NO}_3$  levels. The initial cores were removed from the coring tool and placed in labeled zip-lock plastic bags. The cores were transported to the laboratory in a cooler and stored at  $<4^\circ\text{C}$  until extraction (usually within 2 weeks of sampling).

The IEM soil cores were installed in burned and unburned areas on Grindstone and Betenson Flats: four cores were installed in each of the four exclosures in burned aspen/conifer stands, four cores were installed in each of three unburned aspen/conifer stands near the burned areas, and four cores were installed in each of two unburned aspen stands that regenerated as a result of a fire in 1958. Soil temperature in each area was monitored at a 4-inch depth using Optic StowAway temperature loggers (Onset Computer Corp., Bourne, MA).

Two incubation periods were used each summer: 6/25 to 7/22/97 (27 d), 7/22 to 9/23/97 (63 d), 6/23–24 to 7/21–22/98 (27–28 d), 7/21–22 to 9/19/98 (59–60 d), 7/1 to 7/29/99 (28 d), and 7/29 to 9/19/99 (52 d). At the end of each incubation period, the leaf litter or ash layer was removed from the tops of the cores and the cores were removed from the ground and placed in labeled zip-lock plastic bags. The cores were transported to the laboratory in a cooler and stored at  $<4^\circ\text{C}$  until extraction. New initial cores for the second incubation period were collected when the IEM soil cores from the first period were removed and the new IEM soil cores were installed.

The IEM soil cores were disassembled, and top and bottom CEM-AEM pairs were rinsed with deionized water to remove soil particles and placed in

separate labeled petri dishes. The soil cores were removed from the plastic liners, returned to the zip-lock bags, and mixed by kneading the bags. Each CEM-AEM pair was extracted with 100 mL of 2 M KCl in 250-mL wide-mouth plastic jars by shaking for 1 h on an orbital shaker at 150 rpm. The extracts were frozen until analysis.

Water content of a representative subsample of each soil core was determined gravimetrically at 105 °C. A representative field-moist subsample of each soil core equivalent to 10 g of oven-dry soil was extracted with 100 mL of 2 M KCl in a 250-mL wide-mouth plastic jar by shaking for 1 h on an orbital shaker at 150 rpm. The extracts were filtered through Whatman 40 filter paper and frozen until analysis. The 2 M KCl extracts of IEMs and soil cores were analyzed for  $\text{NH}_4\text{-N}$  and  $\text{NO}_3\text{-N}$  by the automated phenate and Cd reduction methods, respectively, using a LACHAT flow injection analysis system (Zellweger Analytics, Milwaukee, WI).

The remaining soil from each soil core was air dried at room temperature and weighed. The total weight of each soil core was calculated by summing the oven-dry weight of the subsample used for determining water content, the oven-dry weight of the subsample used for determining extractable  $\text{NH}_4$  and  $\text{NO}_3$ , and the air-dry weight of the remaining soil in each core.

Amounts of  $\text{NH}_4\text{-N}$  and  $\text{NO}_3\text{-N}$  accumulated by the IEMs ( $\text{ug/cm}^2$ ) were calculated by multiplying the  $\text{NH}_4$  and  $\text{NO}_3$  concentrations in the KCl extracts by the extract volume and dividing by the surface area of the IEM ( $20.27 \text{ cm}^2$ ). Rates of  $\text{NH}_4$  or  $\text{NO}_3$  accumulation by the cores ( $\text{ug/cm}^2/\text{mo}$ ) were calculated by dividing the accumulated amounts by the incubation time in days and multiplying by 30 days per month. The IEMs at the top of the soil core remove  $\text{NH}_4$  and  $\text{NO}_3$  in precipitation or leachate passing through the leaf litter or ash layers. The IEMs at the bottom of the soil core remove  $\text{NH}_4$  and  $\text{NO}_3$  leached from the soil core. Concentrations of  $\text{NH}_4$  and  $\text{NO}_3$  in the top and bottom IEMs on a soil weight basis for the entire soil core ( $\text{mg/kg}$ ) were calculated by multiplying the amounts on the IEMs by the IEM surface area and dividing by the dry weight of the soil core.

Total  $\text{NH}_4$  and  $\text{NO}_3$  concentrations in the IEM soil cores were obtained by summing the concentrations of  $\text{NH}_4$  or  $\text{NO}_3$  in the soil core plus the amounts extracted from the bottom IEMs. Following terminology and definitions given by Stevenson (1986) and DiStefano and Gholz (1986), net ammonification is equal to total  $\text{NH}_4$  in the IEM soil core minus  $\text{NH}_4$  in the initial soil core. Net nitrification is equal to total  $\text{NO}_3$  in the IEM soil core minus  $\text{NO}_3$  in the initial soil core. Net mineralization (or immobilization) is equal to net ammonification plus net nitrification. Net ammonification, nitrification, and mineralization rates were calculated by dividing the net amounts by the incubation time. An example of these calculations for one of the soil cores from the burn area is shown in table 1.

### **Data Analysis**

Because the data were nonnormally distributed, a Kruskal-Wallis (K-W) analysis of variance on ranks was run on each measured or derived N variable for each incubation period in each year with stand type as the factor and unburned aspen, unburned mixed aspen/conifer, and burned mixed aspen/conifer as levels within stand type (SigmaStat, ver. 2.0, SPSS Science, Chicago, IL). If a statistically significant difference among stand types was found ( $p < 0.05$ ), a pairwise multiple comparisons test was conducted using Dunn's method to isolate which stand type was different from the others ( $p < 0.05$ ). Also,

**Table 1**—Net ammonification, nitrification, and mineralization/immobilization for an IEM soil core from the Betenson 1 enclosure (7/22–9/23/97).

Initial soil core NH <sub>4</sub> -N, mg/kg	39.0
Final IEM soil core NH <sub>4</sub> -N, mg/kg	107.7
Bottom CEM NH <sub>4</sub> -N, mg/kg	0.3
Final total (bottom CEM + IEM soil core) NH <sub>4</sub> -N, mg/kg	108.0
Net (final total – initial) NH <sub>4</sub> -N, mg/kg	69.1
Net ammonification rate (net NH <sub>4</sub> -N / incubation time), mg/kg/mo	32.9
Initial soil core NO <sub>3</sub> -N, mg/kg	11.6
Final IEM soil core NO <sub>3</sub> -N, mg/kg	2.0
Bottom AEM NO <sub>3</sub> -N, mg/kg	10.0
Final total (bottom AEM + IEM soil core) NO <sub>3</sub> -N, mg/kg	12.0
Net (final total – initial) NO <sub>3</sub> -N, mg/kg	0.4
Net nitrification rate (net NO <sub>3</sub> -N / incubation time), mg/kg/mo	0.2
Net mineralization/immobilization rate (net ammonification + net nitrification), mg/kg/mo	33.1

Kolmogorod-Smirnov (K-S) tests were run to determine if the distribution of each variable for each incubation in each year differed by stand type (SYSTAT ver. 8.0, SPSS Science, Chicago, IL).

## Results


The Pole Creek fire affected the post-fire rates of accumulation of NH<sub>4</sub> and NO<sub>3</sub> by the top and bottom IEMs in the incubated soil cores (table 2, figures 2 and 3). Rates of NH<sub>4</sub> accumulation by top and bottom CEMs in burned area soil cores were significantly greater than those in unburned areas during the first 1997 incubation, but were not significantly different during the second.

Higher rates of NH<sub>4</sub> accumulation by the top CEMs in unburned areas were observed in 1998 than in 1997. This resulted in no significant differences


**Table 2**—Median rates of NH<sub>4</sub> and NO<sub>3</sub> accumulation by IEMs at the top and bottom of IEM soil cores for two incubations in each of 3 years in unburned aspen, unburned aspen/conifer, and burned aspen/conifer. Different letters indicate that the median values for each stand type are significantly different within each incubation and year.

Stand	Incubation number and year					
	1997		1998		1999	
	1	2	1	2	1	2
	----- ug/cm <sup>2</sup> /mo -----					
<b>Top CEM NH<sub>4</sub>-N</b>						
Unburned aspen	1.5 a	3.2 a	4.5 a	3.6 ab	1.0 a	1.2 a
Unburned aspen/conifer	1.6 a	3.1 a	7.6 a	5.6 a	3.0 ab	1.7 a
Burned aspen/conifer	10.3 b	1.0 a	6.4 a	0.2 b	9.8 b	1.6 a
<b>Bottom CEM NH<sub>4</sub>-N</b>						
Unburned aspen	1.5 a	0.8 a	1.3 a	0.8 a	0.6 a	0.5 ab
Unburned aspen/conifer	1.7 a	0.4 a	1.3 ab	1.1 a	1.2 a	0.2 a
Burned aspen/conifer	4.5 b	0.8 a	2.6 b	1.0 a	1.2 a	0.8 b
<b>Top AEM NO<sub>3</sub>-N</b>						
Unburned aspen	0.8 a	0.2 a	0.9 a	0.4 a	1.0 a	0.4 a
Unburned aspen/conifer	0.6 a	0.4 a	0.8 a	1.0 a	1.4 a	0.2 a
Burned aspen/conifer	0.6 a	44.4 b	78.6 b	91.0 b	29.1 b	26.6 b
<b>Bottom AEM NO<sub>3</sub>-N</b>						
Unburned aspen	0.5 a	0.2 a	0.8 a	0.5 a	0.8 a	0.0 a
Unburned aspen/conifer	0.6 a	0.3 a	0.9 a	0.3 a	0.8 a	0.1 a
Burned aspen/conifer	4.8 b	94.6 b	26.6 b	85.6 b	4.6 b	7.2 b

**Figure 2**—Box plots of rates of  $\text{NH}_4$  accumulation by cation exchange membranes (CEM) at the top and bottom of IEM soil cores for two incubation periods in each of 3 years in unburned aspen, unburned aspen/conifer, and burned aspen/conifer. The 25<sup>th</sup> and 75<sup>th</sup> percentiles are shown as a box centered about the median (50<sup>th</sup> percentile), the 10<sup>th</sup> and 90<sup>th</sup> percentiles are shown as error bars, and the 5<sup>th</sup> and 95<sup>th</sup> percentiles and outliers are shown as points.



**Figure 3**—Box plots of rates of  $\text{NO}_3$  accumulation by anion exchange membranes (AEM) at the top and bottom of IEM soil cores for two incubation periods in each of 3 years in unburned aspen, unburned aspen/conifer, and burned aspen/conifer. The 25<sup>th</sup> and 75<sup>th</sup> percentiles are shown as a box centered about the median (50<sup>th</sup> percentile), the 10<sup>th</sup> and 90<sup>th</sup> percentiles are shown as error bars, and the 5<sup>th</sup> and 95<sup>th</sup> percentiles and outliers are shown as points.



between burned and unburned areas for rates of  $\text{NH}_4$  accumulation by the top CEMs during the first 1998 incubation. During the second 1998 incubation, higher rates of  $\text{NH}_4$  accumulation by the top CEMs were found in unburned than burned areas. In contrast, higher rates of  $\text{NH}_4$  accumulation by the bottom CEMs were found in burned areas than unburned areas during the first 1998 incubation. No significant differences between burned and unburned areas were found in  $\text{NH}_4$  accumulation rates by the bottom CEMs during the second 1998 incubation.

Higher rates of  $\text{NH}_4$  accumulation were again found in the top CEMs in burned areas compared to unburned areas during the first 1999 incubation, but not during the second. Stand type had no effect on  $\text{NH}_4$  accumulation rates by the bottom CEMs during the first incubation in 1999, and although significant differences due to stand type were observed during the second 1999 incubation, these differences were minor because accumulation rates were low during this period.

During the first 1997 incubation,  $\text{NO}_3$  accumulation rates by the top AEMs in burned areas were not significantly different from those in unburned areas. However,  $\text{NO}_3$  accumulation rates by bottom AEMs were significantly greater in burned areas than unburned areas during the first 1997 incubation. During the second 1997 incubation,  $\text{NO}_3$  accumulation rates by both top and bottom AEMs were significantly greater in burned areas than in unburned areas. The higher rates of  $\text{NO}_3$  accumulation by top and bottom AEMs in burned areas compared to unburned areas continued in 1998 and 1999.

Higher rates of  $\text{NH}_4$  accumulation by the CEMs were found during the first incubation than during the second, whereas higher rates of  $\text{NO}_3$  accumulation by the AEMs were found during the second incubation than during the first. The longer second incubation allowed more time for greater quantities of  $\text{NH}_4$  and  $\text{NO}_3$  to accumulate on the IEMs, but the quantities accumulated on the IEMs are divided by incubation time to obtain rates of accumulation to adjust for the time difference between the first and second incubation. This shift in the rates of accumulation by the IEMs and relative distribution of inorganic N species from the first to the second incubation period indicates an increase in net soil nitrification rates in burned areas as summer progressed. This pattern continued for  $\text{NH}_4$  in top and bottom CEMs and  $\text{NO}_3$  in bottom AEMs in 1998 and 1999, but not for  $\text{NO}_3$  in top AEMs. Rates of  $\text{NO}_3$  accumulation by top AEMs during the first incubation were similar to those during the second incubation in 1998 and 1999.


The distribution of  $\text{NH}_4$  and  $\text{NO}_3$  among the top and bottom IEMs tended to follow the relative mobility of the ions. Nitrate is more mobile than  $\text{NH}_4$ , so similar rates of  $\text{NO}_3$  accumulation were found in top and bottom AEMs in burned areas. In contrast, higher rates of  $\text{NH}_4$  accumulation were found in top than bottom CEMs.

During the first 1997 incubation, significantly more  $\text{NH}_4$  and  $\text{NO}_3$  were extracted from IEM soil cores (including bottom IEMs) in burned areas than from those in unburned areas (table 3, figure 4). Extractable  $\text{NH}_4$  in IEM soil

**Table 3**—Median soil + bottom CEM  $\text{NH}_4$  and soil + bottom AEM  $\text{NO}_3$  in IEM soil cores for two incubation periods in each of 3 years in unburned aspen, unburned aspen/conifer, and burned aspen/conifer. Different letters indicate that the median values for each stand type are significantly different within each incubation and year.

Stand	Incubation number and year					
	1997		1998		1999	
	1	2	1	2	1	2
----- mg/kg -----						
<b>Soil + CEM <math>\text{NH}_4</math>-N</b>						
Unburned aspen	3.8 a	2.0 a	2.2 a	3.2 a	1.8 ab	1.4 ab
Unburned aspen/conifer	5.1 a	5.2 a	10.8 a	13.0 a	3.5 a	4.0 a
Burned aspen/conifer	22.3 b	7.2 a	1.0 b	1.0 b	1.0 b	0.6 b
<b>Soil + AEM <math>\text{NO}_3</math>-N</b>						
Unburned aspen	0.6 a	0.6 a	1.5 a	4.6 a	0.1 a	2.1 a
Unburned aspen/conifer	0.5 a	0.8 a	1.4 a	1.7 a	0.2 a	2.0 a
Burned aspen/conifer	10.2 b	35.4 b	13.9 b	32.7 b	3.8 b	8.8 b

**Figure 4**—Box plots of soil + CEM  $\text{NH}_4$  and soil + AEM  $\text{NO}_3$  in IEM soil cores for two incubation periods in each of 3 years in unburned aspen, unburned aspen/conifer, and burned aspen/conifer. The 25<sup>th</sup> and 75<sup>th</sup> percentiles are shown as a box centered about the median (50<sup>th</sup> percentile), the 10<sup>th</sup> and 90<sup>th</sup> percentiles are shown as error bars, and the 5<sup>th</sup> and 95<sup>th</sup> percentiles and outliers are shown as points.




cores was not significantly different between burned and unburned areas during the second 1997 incubation. Significantly more  $\text{NO}_3$  was extracted from IEM soil cores in burned areas than those from unburned areas during both 1997 incubations (table 3, figure 4). No significant differences were found between unburned aspen and unburned aspen/conifer for IEM soil core  $\text{NH}_4$  or  $\text{NO}_3$  during the first or second 1997 incubations.

In contrast with the 1997 results, more  $\text{NH}_4$  tended to be extracted from IEM soil cores in unburned areas than in burned areas in 1998. Soil core  $\text{NO}_3$  levels in burned areas in 1998 were similar to those observed in the same areas in 1997. Soil core  $\text{NO}_3$  levels in unburned areas in 1998 tended to be higher than those observed in 1997, but were still significantly less than those in burned areas. The pattern of higher soil core  $\text{NH}_4$  levels in unburned areas than burned areas and higher soil core  $\text{NO}_3$  levels in burned than unburned areas continued in 1999.

To obtain an estimate of the total supply of inorganic N available for plant uptake or microbial activity during the summer months, the total amounts of  $\text{NH}_4$  and  $\text{NO}_3$  accumulated by the top and bottom IEMs from the first and second incubations were added to the total amounts of these ions remaining in the soil cores at the end of the second incubation period (table 4, figure 5).

**Table 4**—Median total IEM (top and bottom CEMs and AEMs from first and second incubations) + soil (IEM soil core from second incubation)  $\text{NH}_4 + \text{NO}_3$ -N in unburned aspen, unburned aspen/conifer, and burned aspen/conifer in 1997, 1998, and 1999. Different letters indicate that the median values for each stand type are significantly different within each year.

Stand	Year		
	1997	1998	1999
	----- mg/kg -----		
Unburned aspen	4.6 a	10.1 a	4.4 a
Unburned aspen/conifer	9.0 a	21.1 a	8.2 ab
Burned aspen/conifer	69.4 b	67.8 b	20.2 b



**Figure 5**—Box plots of total IEM (top and bottom CEMs and AEMs from first and second incubations) + soil (IEM soil core from second incubation)  $\text{NH}_4 + \text{NO}_3\text{-N}$  in unburned aspen, unburned aspen/conifer, and burned aspen/conifer in 1997, 1998, and 1999. The 25<sup>th</sup> and 75<sup>th</sup> percentiles are shown as a box centered about the median (50<sup>th</sup> percentile), the 10<sup>th</sup> and 90<sup>th</sup> percentiles are shown as error bars, and the 5<sup>th</sup> and 95<sup>th</sup> percentiles and outliers are shown as points.


Similar levels of  $\text{NH}_4 + \text{NO}_3\text{-N}$  were available for plant uptake or microbial activity in burned areas in 1997 and 1998, and these levels were significantly greater than those in unburned areas. In unburned areas, more  $\text{NH}_4 + \text{NO}_3\text{-N}$  was observed in 1998 than in 1997. In 1999,  $\text{NH}_4 + \text{NO}_3\text{-N}$  in unburned areas had declined to levels observed in 1997. Amounts of  $\text{NH}_4 + \text{NO}_3\text{-N}$  in burned areas also decreased in 1999, but were still significantly higher than those in unburned aspen but not in unburned aspen/conifer.

In 1997, net ammonification, nitrification, and mineralization/immobilization rates were not significantly different between burned and unburned areas or between unburned aspen and unburned aspen/conifer stands (table 5, figure 6). Even though median rates were not significantly different, there tended to be a greater net loss of  $\text{NH}_4$  in burned areas during the early part of summer (first incubation) than later in the summer (second incubation). Net gains in soil  $\text{NO}_3$  in burned areas tended to be greater during the latter part

**Table 5**—Median net ammonification, nitrification, and mineralization/immobilization rates in IEM soil cores for two incubation periods in each of 3 years in unburned aspen, unburned aspen/conifer, and burned aspen/conifer. Different letters indicate that the median values for each stand type are significantly different within each incubation and year.

Stand	Incubation number and year					
	1997		1998		1999	
	1	2	1	2	1	2
----- mg/kg -----						
<b>Net ammonification rate</b>						
Unburned aspen	0.7 a	-0.2 a	0.9 ab	1.2 ab	1.5 a	0.6 ab
Unburned aspen/conifer	1.8 a	1.4 a	4.2 a	5.4 a	0.8 a	1.6 a
Burned aspen/conifer	-6.3 a	-7.3 a	-0.2 b	0.1 b	-0.5 a	-0.1 b
<b>Net nitrification rate</b>						
Unburned aspen	-0.2 a	0.2 a	0.5 a	1.8 ab	-0.3 a	0.5 a
Unburned aspen/conifer	-0.3 a	0.2 a	0.2 a	0.2 a	-0.9 a	0.8 a
Burned aspen/conifer	4.2 a	12.2 a	2.6 a	7.5 b	-0.6 a	2.7 b
<b>Net N mineralization rate</b>						
Unburned aspen	0.5 a	0.5 a	1.8 a	3.0 a	1.6 a	1.4 a
Unburned aspen/conifer	1.6 a	1.8 a	4.2 a	6.9 a	0.2 a	3.4 a
Burned aspen/conifer	8.4 a	3.9 a	2.4 a	7.4 a	-0.5 a	2.5 a

**Figure 6**—Box plots of net ammonification, nitrification, and mineralization/immobilization rates in IEM soil cores for two incubation periods in each of 3 years in unburned aspen, unburned aspen/conifer, and burned aspen/conifer. The 25<sup>th</sup> and 75<sup>th</sup> percentiles are shown as a box centered about the median (50<sup>th</sup> percentile), the 10<sup>th</sup> and 90<sup>th</sup> percentiles are shown as error bars, and the 5<sup>th</sup> and 95<sup>th</sup> percentiles and outliers are shown as points.



of summer than in the early part. The loss in  $\text{NH}_4$  was largely balanced by the gain in  $\text{NO}_3$  (nitrification), although some loss of  $\text{NH}_4$  may have been from direct microbial utilization of this form of N. In 1998 and 1999, net ammonification rates tended to be higher in unburned areas than burned areas, whereas, the opposite was observed for net nitrification rates. Because of the relative balance of net ammonification and net nitrification rates, net mineralization/immobilization rates in burned and unburned areas tended to be similar in 1998 and 1999.

For cases where the K-W tests revealed that median values of the various N variables for the different stand types differed significantly, K-S tests showed that the data distributions also differed as the Box plots indicate (figures 2, 3, 4, 5, and 6). The range of values tended to be much greater in burned than in unburned areas.

## Discussion

In undisturbed (unburned) aspen and mixed aspen/conifer soils, the various microbial mineralization and immobilization reactions of the internal soil N cycle tend to be balanced. There is usually little free  $\text{NH}_4$  and  $\text{NO}_3$  in the mineral soil (bottom of figures 2 and 3, figures 4 and 5) because it is quickly utilized by microorganisms or taken up by plant roots. Thus, net mineralization/immobilization rates tend to be low (figure 6). Low or unchanged levels of  $\text{NH}_4$  and  $\text{NO}_3$  and low or unchanged rates of net mineralization do not mean that active mineralization is not taking place or that the internal soil N cycle is not operating (Stevenson 1986). Gross rates of the microbially mediated reactions may be high, but since the reactions tend to be balanced, net rates are low (Stevenson 1986).

Depending on its intensity, fire alters many soil properties including exerting a strong influence on soil microbial mineralization/immobilization

processes (Schlesinger 1997). Losses of gaseous and particulate N occur during combustion of leaf litter and soil organic matter. Microbial populations and enzyme activity may decrease, but available nutrients are also returned to the soil by the residual ash (Schlesinger 1997). Even though total soil N may be lower because of combustion, residual N in the ash is subjected to rapid mineralization and nitrification provided the fire is not too intense and microbial and enzyme populations can recover (Schlesinger 1997).

The source of the high amounts of  $\text{NH}_4$  accumulated by the CEMs and in the soil cores in the burn area the first summer after the fire may include  $\text{NH}_4$  as a combustion product in the ash layer, release from mineral sources of  $\text{NH}_4$ , and rapid mineralization of combustion products. Much of this available  $\text{NH}_4$  was probably carried over in the soil from the summer in which the fire occurred (6 to 7/96). Nitrification (microbial oxidation of  $\text{NH}_4$  to  $\text{NO}_3$ ) occurred mostly during the first summer after the fire (6 to 9/97) because of the observed decrease in  $\text{NH}_4$  and increase in  $\text{NO}_3$  accumulated by the IEMs and soil cores from the first to the second incubation period. The IEM soil core incubation method provides a snapshot of soil N mineralization/immobilization for the incubation periods studied, but since we did not have soil cores in place immediately after the fire, we could not quantify soil N transformations in late summer and fall of the fire year (7 to 9/96).

Nitrification is influenced by many factors, including temperature, moisture, and the substrates  $\text{NH}_4$ ,  $\text{O}_2$ , and  $\text{CO}_2$  (Stevenson 1986). The high amounts of available  $\text{NH}_4$  found in burn area soils provide a ready substrate for surviving or post-fire invading populations of nitrifying microorganisms. Summertime soil temperature ranges and means in the burn area were higher than in unburned areas because the relatively open canopy and dark colored ash served to trap heat (data not shown), thus helping to accelerate nitrification. Adequate soil moisture was available in burned areas (mean soil core water content of 24.1% in 1997) for nitrification to occur throughout the summer.

Although denitrification and leaching can potentially remove much of the  $\text{NO}_3$  produced by post-fire nitrification, the large amounts of available soil  $\text{NH}_4$  and  $\text{NO}_3$  present during the growing season provide a ready source of mineral N for uptake by aspen suckers. A strong aspen suckering response was observed on Betenson and Grindstone Flats and in other burned areas on the Fishlake National Forest during the first summer after the fires (Amacher and Bartos 1998). Leaching of  $\text{NH}_4$  from the ash layer and subsequent nitrification of the leached  $\text{NH}_4$  continued through the second and third summers after the fire. A longer residence time of post-fire soil  $\text{NO}_3$  provides more opportunity for regenerating aspen clones to exploit this source of N. Thus, available N from ash may be an important source of N to young aspen until N mineralization in leaf litter and soil organic matter can resupply the nutritional needs of aspen.

## Acknowledgments

This work was supported by a USDA-CSREES Rangelands Special Grant. We thank Scott Walker of the Utah Division of Wildlife Resources and Bob Campbell, Dandy Pollack, and Doug Sorenson of the Fishlake National Forest for their assistance throughout the project. We also thank Utah State University Analytical Labs for assistance with the sample analyses.

## References

- Amacher, M. C.; Bartos, D. L. 1998. Soil properties associated with aspen to conifer succession and aspen restoration treatments. In: Abstracts: 51<sup>st</sup> annual meeting; Society for Range Management; 1998 Feb. 8-12; Guadalajara, Jalisco, Mexico. Denver, CO: Society for Range Management: 55.
- Bartos, D. L.; Campbell, R. B., Jr. 1998a. Decline of quaking aspen in the Interior West—examples from Utah. *Rangelands* 20(1): 17-24.
- Bartos, D. L.; Campbell, R. B., Jr. 1998b. Water depletion and other ecosystem values forfeited when conifer forests displace aspen communities. In: Potts, D. F., ed. *Rangeland management and water resources*; 1998 May 27-29; Reno, NV. Herndon, VA: American Water Resources Assoc.: 427-434.
- Cooperband, L. R.; Logan, T. J. 1994. Measuring in situ changes in labile soil phosphorus with anion-exchange membranes. *Soil Sci. Soc. Am. J.* 58: 105-114.
- DiStefano, J. F.; Gholz, H. L. 1986. A proposed use of ion exchange resins to measure nitrogen mineralization and nitrification in intact soil cores. *Commun. Soil Sci. Plant Anal.* 17: 989-998.
- Harniss, R. O. 1981. Ecological succession in aspen and its consequences. In: Debyle, N. V., ed. *Situation management of two intermountain species: aspen and coyotes*. Vol. 1. Aspen; 1981 April 23-24; Logan, UT. Logan, UT: Utah State Univ.: 31-39.
- Jones, J. R.; Debyle, N. V. 1985. Fire. In: Debyle, N. V.; Winokur, R. P., eds. *Aspen: ecology and management in the Western United States*. Gen. Tech. Rep. RM-119. Ft. Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station: 77-81.
- Kay, C. E. 1990. *Yellowstone's northern elk herd: a critical evaluation of the "natural regulation" paradigm*. Logan, UT: Utah State Univ. Dissertation. 490 p.
- Schier, G. A. 1981. Aspen regeneration. In: Debyle, N. V., ed. *Situation management of two intermountain species: aspen and coyotes*. Vol. 1. Aspen; 1981 April 23-24; Logan, UT. Logan, UT: Utah State Univ.: 15-21.
- Schlesinger, W. H. 1997. *Biogeochemistry. An analysis of global change*. 2<sup>nd</sup> ed. San Diego: Academic Press. 588 p.
- Stevenson, F. J. 1986. *Cycles of soil*. New York: John Wiley & Sons. 380 p.

# Aspen Forest Products

