

LANDFIRE Biophysical Setting Model

Biophysical Setting 1311250

Inter-Mountain Basins Big Sagebrush Steppe

- This BPS is lumped with:
 This BPS is split into multiple models:

General Information

Contributors (also see the Comments field) **Date** 2/23/2005

Modeler 1 Mike Zielinski mike_zielinski@nv.blm.gov **Reviewer**
ov

Modeler 2 Louis lprovencher@tnc.org **Reviewer**
Provencher

Modeler 3 **Reviewer**

Vegetation Type

Upland Savanna and
Shrub-Steppe

Dominant Species

ARTRW8
AGSP

Map Zone

13

Model Zone

- | | |
|---|--|
| <input type="checkbox"/> Alaska | <input type="checkbox"/> Northern Plains |
| <input type="checkbox"/> California | <input type="checkbox"/> N-Cent. Rockies |
| <input type="checkbox"/> | <input type="checkbox"/> |
| <input checked="" type="checkbox"/> Great Basin | <input type="checkbox"/> Pacific Northwest |
| <input type="checkbox"/> Great Lakes | <input type="checkbox"/> South Central |
| <input type="checkbox"/> Hawaii | <input type="checkbox"/> Southeast |
| <input type="checkbox"/> Northeast | <input type="checkbox"/> S. Appalachians |
| | <input type="checkbox"/> Southwest |

General Model Sources

- Literature
 Local Data
 Expert Estimate

STTH2
POSA12

Geographic Range

This widespread matrix-forming ecological system occurs throughout much of the Columbia Plateau and northern Great Basin and WYg and is found at slightly higher elevations farther south.

Biophysical Site Description

Sagebrush steppe is found in continental, semi-arid climate with highly variable annual precipitation greater than 7-12in (~180-300mm) (McArthur 2000) that may also include 14in precipitation zone. Common on foothills, undulating terraces, slopes and plateaus, but also in basins and valley bottoms. Soil depths range from shallow to moderately deep, well-drained with an effective rooting depth of <40in (~1m). NRCS Range Sites: Loamy 8-10in and 10-12in precipitation zones, and shallow loam 10-14in precipitation zones.

Vegetation Description

This shrub-steppe is dominated by perennial grasses and forbs (>25% cover) with *Artemisia tridentata* ssp. *tridentata*, *Artemisia tridentata* ssp. *wyomingensis*, and/or *Purshia tridentata* dominating or codominating the open to moderately dense (10-40% cover) shrub layer. In southern ID and northern UT, *Artemisia tridentata* ssp. *wyomingensis* dominates large landscapes. *Atriplex confertifolia*, *Chrysothamnus viscidiflorus*, *Ericameria nauseosa* or *Tetradymia* spp may be common especially in disturbed stands. Associated graminoids include *Achnatherum hymenoides*, *Elymus lanceolatus* ssp. *lanceolatus*, *Festuca idahoensis*, *Festuca campestris*, *Koeleria macrantha*, *Poa secunda* and *Pseudoroegneria spicata*. Common forbs are *Phlox hoodii*, *Arenaria* spp, and *Astragalus* spp. Areas with deeper soils more commonly support *Artemisia tridentata* ssp. *tridentata* but have largely been converted

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

for other land uses.

The sagebrush steppe landscape is a mosaic of shrub-dominated and herbaceous-dominated phases (West 2000). Forbs have low diversity but are important for wildlife, including the greater sage grouse. Species diversity is lower in Wyoming big sagebrush communities than in other big sagebrush types (FEIS). Wyoming big sagebrush communities are critical habitat for greater sage grouse and other sagebrush obligate species.

Disturbance Description

Historically, fire was the principal disturbance within this vegetation type; other disturbances included insects (eg, moths and grasshoppers that eat leaves, moth larval grubs that eat roots; return interval of 75yrs), periods of drought and wet cycles and shifts in climate (return interval of 100yrs). Intervals between natural wildfires varied between 25yrs (northern Yellowstone National Park [Houston 1973], cited in West 2000) and 100yrs+ (West 2000). West (1983) and Miller and Eddelman (2000) cite mean FRI <100yrs for replacement fire. FEIS cites fire return interval ranges between 10-70yrs with mean of 40yrs for Wyoming sagebrush steppe. Studies cited in FEIS may underestimate FRIs or not hold up to scrutiny (Welch and Criddle 2003). It was assumed that dominant fires were stand replacement (mean FRIs of 75-94yrs) due to the continuity of fine fuel typical of steppe ecosystems. Mixed severity (25-75% of area inside burn perimeter topkilled) played a minor role during mid-development. Assuming a MFI of 75yrs (from the total fire probability), the mean FRI of mixed severity fire was 20% of fires, thus a mean FRI of 375yrs, during mid-development. Re-establishment following fire is from seed germination and establishment. Establishment is dependent upon soil seedbank and/or proximity of seed sources, fire size and continuity and climatic conditions.

Adjacency or Identification Concerns

The NatureServe description of BpS 1125 includes different species of sagebrush and steppe ecosystems that are structurally and ecologically different such as *Artemisia tridentata* ssp. *tridentata* and *Artemisia tridentata* ssp. *wyomingensis*. We highly recommend that, at least, *Artemisia tridentata* ssp. *tridentata*, which is a taller shrub found in drainages and deeper soils, be separated from the other shrubs. Ultimately, the two sagebrush species should be modeled separately. *Artemisia tripartita* ssp. *tripartita* is not part of this system in NV because it is generally associated with frigid soils (thus more typically mountain big sagebrush) under snow pockets. Bitterbrush is not found in a large area of northcentral NV on the more alkaline soils of Pleistocene Lake Lahontan.

Wyoming big sagebrush is known to hybridize with other subspecies of the big sagebrush complex; ie, basin big sagebrush (*A. tridentata* ssp. *tridentata*) and mountain big sagebrush (*A. tridentata* ssp. *vaseyana*) (Freeman et al 1991, McArthur et al 1998). Across ecotones, populations of Wyoming big sagebrush probably intergrade with basin big sagebrush and mountain big sagebrush. Soils and elevation may help determine which species is present.

Invasion of cheatgrass has transformed this ecological system into large areas of uncharacteristic annual grasslands and shrublands with understories where annual grasses replaced perennial grasses.

Native Uncharacteristic Conditions

Scale Description

Sagebrush steppe covers vast landscapes >10000ac with inclusions of low sagebrush and basin big sagebrush. Historic disturbance (fire) likely ranged from small (<10ac) to large (>10000ac) depending on conditions, time since last ignition and fuel loading. Assumed the average patch size is 250ac.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

Issues/Problems

West (2000) cites species wide range in FRI (25-100yrs+). West (1983) and Miller and Eddelman (2000) recommend a FRI of <100yrs for replacement fire. FEIS gives 10-70yr range (40yr average) (but see Welch and Criddle 2003). Current scientific opinion (Mike Pellant, BLM Range Ecologist on the Great Basin Restoration Initiative) puts the natural fire return interval at about 100yrs (confirmed by Stephen Bunting and Dave Pyke). Given uncertainties and opinions of reviewers, a MFI of 75yrs was chosen. Without this shorter MFI and differences in fire behavior, there would be no difference between Wyoming sagebrush steppe from the Snake River plains and Wyoming big sagebrush semi-desert from central NV, UT and eastern CA. Because replacement fire is by far dominant over mixed severity fire, a FRG IV was selected to the recommendation of reviewers.

Comments

BpS 1125 for MZ12 and MZ17 was obtained by slightly modifying the description of BpS 1125 for MZ16 developed by Don Major (dmajor@tnc.org). BpS 1125 for MZ16 is completely based on R2SBWYse developed by Eric Limbach (eric_limbach@blm.gov) for Wyoming big sagebrush steppe and reviewed by Krista Waid-Gollnick/Sarah Heidi (krista_waid@blm.gov), Stanley Kitchen (skitchen@fs.fed.edu), Michael Zielinski (mike_zielinski@nv.blm.gov), Jolie Pollet (jpollet@blm.gov) and Gary Back (gback@srk.com).

As a result of final QC for LANDFIRE National by Kori Blankenship the user-defined min and max fire return intervals for mixed severity fire were deleted because they were not consistent with the modeled fire return interval for this fire severity type.

Vegetation Classes

Class A 20 %

Early Development 1 Open

Upper Layer Lifeform

Herbaceous
 Shrub
 Tree

Fuel Model
1

Indicator Species and Canopy Position
 AGSP Upper
 STTH2 Upper
 POSA12 Upper
 ARTRW8 Upper
 Upper

Structure Data (for upper layer lifeform)

	Min	Max
Cover	0 %	60 %
Height	Herb 0m	Herb 1.0m
Tree Size Class	None	

Upper layer lifeform differs from dominant lifeform.

Vegetation is primarily herbaceous with a few scattered shrubs accounting for less than five percent cover and less than one meter.

Description
 Perennial grasses and forbs dominate where woody shrub canopy has been top killed/removed by wildfire. Shrub cover <6%. (~ 0-19yrs). Replacement fire every 120yrs on average resets succession back to zero. Succession to class B after 20yrs.

Class B 50 %

Mid Development 1 Open

Upper Layer Lifeform

Herbaceous
 Shrub
 Tree

Fuel Model
1

Indicator Species and Canopy Position
 AGSP Lower
 STTH2 Lower
 ARTRW8 Upper
 Upper

Structure Data (for upper layer lifeform)

	Min	Max
Cover	0 %	20 %
Height	Shrub 0m	Shrub 3.0m
Tree Size Class	None	

Upper layer lifeform differs from dominant lifeform.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

POSA12
Lower

Description

Shrubs dominate (5-25% cover) with diverse perennial grass and forb understory (20-60yrs). MFI is 75yrs with 80% replacement fire (mean FRI of 94yrs) and 20% mixed severity fire (mean FRI of 375yrs). Mixed severity fire, insect/disease (return interval of 75yrs) and weather related stress (return interval of 100yrs) maintains vegetation in class B. Succession to class C after 40yrs.

Class C 30 %

Late Development 1 Closed

Upper Layer Lifeform

- Herbaceous
 Shrub
 Tree

Fuel Model
2

Indicator Species and Canopy Position

ARTRW8
Upper
AGSP
Lower
STTH2
Lower
POSA12
Lower

Structure Data (for upper layer lifeform)

	Min	Max
Cover	21 %	40 %
Height	Shrub 0m	Shrub 3.0m
Tree Size Class	None	

Upper layer lifeform differs from dominant lifeform.

Description

Mature shrub canopy >25% cover with proportional reduction in understory productivity as canopy cover increases. The mean FRI for replacement fire is 75yrs. Insect/diseases (return interval of 75yrs), and weather related stress (return interval of 100yrs) thin the shrub canopy, causing a transition to class B. Succession from class C to C.

Class D 0 %

[Not Used] [Not Used]

Upper Layer Lifeform

- Herbaceous
 Shrub
 Tree

Fuel Model

Indicator Species and Canopy Position

Structure Data (for upper layer lifeform)

	Min	Max
Cover	%	%
Height		
Tree Size Class		

Upper layer lifeform differs from dominant lifeform.

Description

Class E 0 %

[Not Used] [Not Used]

Upper Layer Lifeform

- Herbaceous
 Shrub
 Tree

Fuel Model

Indicator Species and Canopy Position

Structure Data (for upper layer lifeform)

	Min	Max
Cover	%	%
Height		
Tree Size Class		

Upper layer lifeform differs from dominant lifeform.

Description

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

Disturbances

Fire Regime Group:** IV

Historical Fire Size (acres)

Avg 250

Min 10

Max 10000

Sources of Fire Regime Data

- Literature
- Local Data
- Expert Estimate

Additional Disturbances Modeled

- Insects/Disease
- Native Grazing
- Other (optional 1)
- Wind/Weather/Stress
- Competition
- Other (optional 2)

Fire Intervals

	Avg FI	Min FI	Max FI	Probability	Percent of All Fires
Replacement	92	30	120	0.01087	89
Mixed	714			0.00140	11
Surface					
All Fires	81			0.01228	

Fire Intervals (FI):

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class.

References

Brown, J.K. and J. Kapler-Smith, eds. 2000. Wildland fire in ecosystems: effects of fire on flora. Gen. Tech. Rep. RMRS-GTR-42-vol. 2. Ogden, UT: USDA Forest Service, Rocky Mountain Research Station. 257 pp.

Freeman, D.C., W.A. Turner, E.D. McArthur and J.H. Graham. 1991. Characterization of a narrow hybrid zone between two subspecies of big sagebrush (*Artemisia tridentata*: Asteraceae). *American Journal of Botany*. 78(6): 805-815.

Houston, D.B. 1973. Wildfires in northern Yellowstone National Park. *Ecology* 54: 1111-1117.

Howard, Janet L. 1999. *Artemisia tridentata* subsp. *wyomingensis*. In: Fire Effects Information System, [Online]. USDA Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). Available: <http://www.fs.fed.us/database/feis/> [2007, June 21].

McArthur, E.D., D.C. Freeman and J.H. Graham. 1998. Narrow hybrid zone between two subspecies of big sagebrush (*Artemisia tridentata*: Asteraceae). VI. Respiration and water potential. *Canadian Journal of Botany*. 76(4): 567-574.

McArthur, E.D. 2000. Sagebrush systematics and distribution. Pages 9-14 in: P.G. Entwistle, A.M. DeBolt, J.H. Kaltenecker and K. Steenhof, compilers. Proceedings: Sagebrush Steppe Ecosystems Symposium. Bureau of Land Management Publication No. BLM/ID/PT-001001+1150, Boise, Idaho, USA.

Miller, R.F. and L.L. Eddleman. 2000. Spatial and temporal changes of sage grouse habitat in the sagebrush biome. Oregon State University Agricultural Experiment Station Technical Bulletin 151, Corvallis, Oregon. 35 pp.

NatureServe. 2007. International Ecological Classification Standard: Terrestrial Ecological Classifications. NatureServe Central Databases. Arlington, VA. Data current as of 10 February 2007.

Peters, E.F. and S.C. Bunting. 1994. Fire conditions pre- and post-occurrence of annual grasses on the

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

Snake River plain. Pages 31-36 in: Proceedings - Ecology, management, and restoration of Intermountain rangelands symposium. USDA Forest Service INT-GTR-313, Ogden, Utah.

Welch, B.L. and C. Criddle. 2003. Countering Misinformation Concerning Big Sagebrush. Research Paper RMRS-RP-40. Ogden, UT: USDA Forest Service, Rocky Mountain Research Station. 28 pp.

West, N.E. 1983. Western Intermountain sagebrush steppe. Pages 351-395 in: N.E. West, ed. Ecosystems of the World 5: Temperate deserts and semi-deserts. Elsevier Scientific Publishing Co., New York, NY.

West, N.E. 2000. Synecology and disturbance regimes of sagebrush steppe ecosystems. Pages 15-26 in: P.G. Entwhistle, A.M. DeBolt, J.H. Kaltenecker and K. Steenhof, compilers. Proceedings: Sagebrush Steppe Ecosystems Symposium. Bureau of Land Management Publication No. BLM/ID/PT-001001+1150, Boise, Idaho, USA.

Whisnant, S.G. 1990. Changing fire frequencies on Idaho's Snake River plains: Ecological and management implications. Pages 4-10 in: E.D. McArthur, E.M. Romme, S.D. Smith and P.T. Tueller, eds. Proceedings of a symposium on cheatgrass invasion, shrub die-off, and other aspects of shrub biology and management. USDA Forest Service Gen. Tech. Rep. INT-276. Intermountain Forest and Range Experiment Station, Ogden, Utah.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.