

LANDFIRE Biophysical Setting Model

Biophysical Setting 1511250

Inter-Mountain Basins Big Sagebrush Steppe

- This BPS is lumped with:
 This BPS is split into multiple models:

General Information

Contributors (also see the Comments field)

Date 2/23/2005

Modeler 1 Don Major dmajor@tnc.org

Reviewer Tim Christiansen tchristiansen@tnc.org

Modeler 2

Reviewer

Modeler 3

Reviewer

Vegetation Type

Upland
Savannah/Shrub
Steppe

Dominant Species

ARTRW8
PSSP6

Map Zone

15

Model Zone

Alaska
California

Northern Plains
N-Cent. Rockies

General Model Sources

- Literature
 Local Data
 Expert Estimate

STTH2
POSA12

Great Basin
 Great Lakes
 Hawaii
 Northeast

Pacific Northwest
 South Central
 Southeast
 S. Appalachians
 Southwest

Geographic Range

This widespread matrix-forming ecological system occurs throughout much of the Columbia Plateau, northern Great Basin and WY. It is found at slightly higher elevations farther south.

Biophysical Site Description

Sagebrush steppe is found in continental, semi-arid climate where annual precipitation generally ranges from 7-12in (~180-300mm) (McArthur 2000), but occasionally ranging into the 14in precipitation zone. This system is common on foothills, undulating terraces, slopes and plateaus, but also in basins and valley bottoms. Soil depths range from shallow to moderately deep, well-drained with an effective rooting depth of less than 40in (~ 1m). NRCS Range Site is (Droughty) Loamy 8-10in precipitation zone.

Vegetation Description

This shrub-steppe is dominated by perennial grasses and forbs (>25% cover) with *Artemisia tridentata* ssp. *tridentata*, *Artemisia tridentata* ssp. *xericensis*, *Artemisia tridentata* ssp. *wyomingensis*, *Artemisia tripartita* ssp. *tripartita* and/or *Purshia tridentata* dominating or codominating the open to moderately dense (10-40% cover) shrub layer. In southern ID and northern UT, *Artemisia tridentata* ssp. *wyomingensis* dominates large landscape. *Atriplex confertifolia*, *Chrysothamnus viscidiflorus*, *Ericameria nauseosa*, *Tetradymia* spp or *Artemisia frigida* may be common, especially in disturbed stands. Associated graminoids include *Achnatherum hymenoides*, *Calamagrostis montanensis*, *Elymus lanceolatus* ssp. *lanceolatus*, *Festuca idahoensis*, *Festuca campestris*, *Koeleria macrantha*, *Poa secunda* and *Pseudoroegneria spicata*. Common forbs are *Phlox hoodii*, *Arenaria* spp and *Astragalus* spp. Areas with deeper soils more commonly support *Artemisia tridentata* ssp. *tridentata* but have largely been converted for other land uses.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

The sagebrush steppe landscape is a mosaic of shrub-dominated and herbaceous-dominated phases (West 2000). Forbs have low diversity but are important for wildlife, including the greater sage grouse. Species diversity is lower in Wyoming big sagebrush communities than in other big sagebrush types (Howard 1999). Wyoming big sagebrush communities are critical habitat for greater sage grouse and other sagebrush obligate species.

Disturbance Description

Historically, fire was the principal disturbance within this vegetation type. Intervals between natural wildfires varied between 25yrs (eg, northern Yellowstone National Park, West 2000) and 100yrs+ (West 2000). West (1983) and Miller and Eddelman (2000) cite mean FRIs of <100yrs for replacement fire. Howard (1999) cites mean FRIs of 10 to 70yrs with a mean of 40yrs for Wyoming sagebrush steppe. Studies cited in Howard (1999) may underestimate FRIs or not hold up to scrutiny (Welch and Criddle 2003). This BpS was modeled with a MFI of 75yrs for the system as a whole. Eighty percent of fires modeled were replacement severity. Re-establishment following fire is from seed germination and establishment. Establishment is dependent upon soil seedbank and/or proximity of seed sources, fire size and continuity, and climatic conditions.

Other disturbances included insects (eg, moths and grasshoppers that eat leaves, moth larval grubs that eat roots) with an average return interval of 75yrs. Periods of drought, wet cycles and shifts in climate would have also affected this system with an average return interval of 100yrs.

Adjacency or Identification Concerns

The NatureServe description of Inter-Mountain Basin Big Sagebrush Steppe (1125) includes different species of sagebrush and steppe ecosystems that are structurally and ecologically different such as *Artemisia tridentata* ssp. *tridentata*, *Artemisia tridentata* ssp. *wyomingensis* and *Artemisia tripartita* ssp. *tripartita*. We highly recommend that, at least, *Artemisia tridentata* ssp. *tridentata*, which is a taller shrub found in drainages and deeper soils, be separated from the other shrubs. Ultimately, all three sagebrush species should be modeled separately.

Wyoming big sagebrush is known to hybridize with other subspecies of the big sagebrush complex; ie, basin big sagebrush *A. tridentata* ssp. *tridentata* and mountain big sagebrush *A. tridentata* ssp. *vaseyana* (Freeman et al 1991, McArthur et al 1998). Across ecotones, populations of Wyoming big sagebrush probably intergrade with basin big sagebrush and mountain big sagebrush. Soils and elevation may help determine which species is present.

Invasion of cheatgrass has transformed this ecological system into large areas of uncharacteristic annual grasslands and shrublands with understories where annual grasses replaced perennial grasses.

Native Uncharacteristic Conditions

Scale Description

Sagebrush steppe covers vast landscapes >10000ac with inclusions of low sagebrush and basin big sagebrush. Historic disturbance (fire) likely ranged from small (<10ac) to large (>10000ac) depending on conditions, time since last ignition and fuel loading. The average patch size was assumed to be approximately 250ac.

Issues/Problems

West (2000) cites wide range in FRI (25-100yrs+). West (1983) and Miller and Eddelman (2000)

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

recommend a FRI of <100yrs for replacement fire. Howard (1999) gives 10-70yr range (40yr average) (but see Welch and Criddle 2003). Current scientific opinion (Mike Pellant, BLM Range Ecologist on the Great Basin Restoration Initiative, Stephen Bunting, and David Pyke, personal communication) puts the natural fire return interval at about 100yrs. Given uncertainties and opinions of reviewers, a MFI of 75yrs was modeled. Without this shorter MFI and differences in fire behavior, there would be no difference between Wyoming sagebrush steppe from the Snake River plains and Wyoming big sagebrush semi-desert from central NV, UT, and eastern CA. Because replacement fire is by far dominant over mixed severity fire, a FRG IV was selected by recommendation of reviewers.

Comments

This model is identical to the model for the same BpS in MZ16 (Utah High Plateaus) with minor descriptive changes based on peer review for MZ23 and MZ24. BpS 1125 is completely based on the Rapid Assessment PNVG R2SBWYse developed by Eric Limbach (eric_limbach@blm.gov) for Wyoming big sagebrush steppe and reviewed by Krista Waid-Gollnick, Sarah Heidi (krista_waid@blm.gov), Stanley Kitchen (skitchen@fs.fed.edu), Michael Zielinski (mike_zielinski@nv.blm.gov), Jolie Pollet (jpollet@blm.gov) and Gary Back (gback@srk.com).

As a result of final QC for LANDFIRE National by Kori Blankenship the user-defined min and max fire return intervals for mixed severity fire were deleted because they were not consistent with the modeled fire return interval for this fire severity type.

Vegetation Classes															
<p>Class A 20 %</p> <p>Early Development 1 Open</p> <p><u>Upper Layer Lifeform</u></p> <p><input type="checkbox"/> Herbaceous</p> <p><input checked="" type="checkbox"/> Shrub</p> <p><input type="checkbox"/> Tree</p> <p><u>Fuel Model</u></p> <p>1</p> <p><u>Description</u></p> <p>Perennial grasses and forbs dominate where woody shrub canopy has been top killed / removed by wildfire. Shrub cover less than five percent. Replacement fire every 120yrs on average resets succession back to zero. Succession to class B after 20yrs.</p>	<p><u>Indicator Species and Canopy Position</u></p> <p>PSSP6 Upper</p> <p>STTH2 Upper</p> <p>POSA12 Upper</p> <p>ARTRW8 Upper</p>	<p><u>Structure Data (for upper layer lifeform)</u></p> <table border="1"> <thead> <tr> <th></th> <th>Min</th> <th>Max</th> </tr> </thead> <tbody> <tr> <td>Cover</td> <td>0 %</td> <td>4 %</td> </tr> <tr> <td>Height</td> <td>Shrub 0m</td> <td>Shrub 1.0m</td> </tr> <tr> <td>Tree Size Class</td> <td colspan="2">None</td> </tr> </tbody> </table> <p><input checked="" type="checkbox"/> Upper layer lifeform differs from dominant lifeform.</p> <p>Vegetation is primarily herbaceous with a few scattered shrubs.</p>		Min	Max	Cover	0 %	4 %	Height	Shrub 0m	Shrub 1.0m	Tree Size Class	None		
	Min	Max													
Cover	0 %	4 %													
Height	Shrub 0m	Shrub 1.0m													
Tree Size Class	None														
<p>Class B 50 %</p> <p>Mid Development 1 Open</p> <p><u>Upper Layer Lifeform</u></p> <p><input type="checkbox"/> Herbaceous</p> <p><input checked="" type="checkbox"/> Shrub</p> <p><input type="checkbox"/> Tree</p> <p><u>Fuel Model</u></p> <p>1</p> <p><u>Description</u></p>	<p><u>Indicator Species and Canopy Position</u></p> <p>PSSP6 Lower</p> <p>STTH2 Lower</p> <p>ARTRW8 Upper</p> <p>POSA12 Lower</p>	<p><u>Structure Data (for upper layer lifeform)</u></p> <table border="1"> <thead> <tr> <th></th> <th>Min</th> <th>Max</th> </tr> </thead> <tbody> <tr> <td>Cover</td> <td>5 %</td> <td>25 %</td> </tr> <tr> <td>Height</td> <td>Shrub 0m</td> <td>Shrub 3.0m</td> </tr> <tr> <td>Tree Size Class</td> <td colspan="2">None</td> </tr> </tbody> </table> <p><input type="checkbox"/> Upper layer lifeform differs from dominant lifeform.</p>		Min	Max	Cover	5 %	25 %	Height	Shrub 0m	Shrub 3.0m	Tree Size Class	None		
	Min	Max													
Cover	5 %	25 %													
Height	Shrub 0m	Shrub 3.0m													
Tree Size Class	None														

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

Shrubs dominate (5-25% cover) with diverse perennial grass and forb understory. MFI is 75yrs with 80% replacement fire (mean FRI of 94yrs) and 20% mixed severity fire (mean FRI of 375yrs). Mixed severity fire, insect/disease (return interval of 75yrs) and weather related stress (return interval of 100yrs) maintains vegetation in class B. Succession to class C after 40yrs.

<p>Class C 30 %</p> <p>Late Development 1 Closed</p> <p>Upper Layer Lifeform</p> <p><input type="checkbox"/> Herbaceous <input checked="" type="checkbox"/> Shrub <input type="checkbox"/> Tree</p> <p style="margin-left: 100px;">Fuel Model 2</p>	<p>Indicator Species and Canopy Position</p> <p>ARTRW8 Upper PSSP6 Lower STTH2 Lower POSA12 Lower</p>	<p>Structure Data (for upper layer lifeform)</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th style="text-align: center;"><i>Min</i></th> <th style="text-align: center;"><i>Max</i></th> </tr> </thead> <tbody> <tr> <td>Cover</td> <td style="text-align: center;">26 %</td> <td style="text-align: center;">35 %</td> </tr> <tr> <td>Height</td> <td style="text-align: center;">Shrub 0m</td> <td style="text-align: center;">Shrub 3.0m</td> </tr> <tr> <td>Tree Size Class</td> <td colspan="2" style="text-align: center;">None</td> </tr> </tbody> </table> <p><input type="checkbox"/> Upper layer lifeform differs from dominant lifeform.</p>		<i>Min</i>	<i>Max</i>	Cover	26 %	35 %	Height	Shrub 0m	Shrub 3.0m	Tree Size Class	None	
	<i>Min</i>	<i>Max</i>												
Cover	26 %	35 %												
Height	Shrub 0m	Shrub 3.0m												
Tree Size Class	None													

Description

Mature shrub canopy >25% cover with proportional reduction in understory productivity as canopy cover increases. The mean FRI for replacement fire is 75yrs. Insect/diseases (return interval of 75yrs) and weather related stress (return interval of 100yrs) thin the shrub canopy, causing a transition to class B. Otherwise, this class persists indefinitely.

<p>Class D 0 %</p> <p>[Not Used] [Not Used]</p> <p>Upper Layer Lifeform</p> <p><input type="checkbox"/> Herbaceous <input type="checkbox"/> Shrub <input type="checkbox"/> Tree</p> <p style="margin-left: 100px;">Fuel Model</p>	<p>Indicator Species and Canopy Position</p>	<p>Structure Data (for upper layer lifeform)</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th style="text-align: center;"><i>Min</i></th> <th style="text-align: center;"><i>Max</i></th> </tr> </thead> <tbody> <tr> <td>Cover</td> <td style="text-align: center;">%</td> <td style="text-align: center;">%</td> </tr> <tr> <td>Height</td> <td></td> <td></td> </tr> <tr> <td>Tree Size Class</td> <td colspan="2"></td> </tr> </tbody> </table> <p><input type="checkbox"/> Upper layer lifeform differs from dominant lifeform.</p>		<i>Min</i>	<i>Max</i>	Cover	%	%	Height			Tree Size Class		
	<i>Min</i>	<i>Max</i>												
Cover	%	%												
Height														
Tree Size Class														

Description

<p>Class E 0 %</p> <p>[Not Used] [Not Used]</p> <p>Upper Layer Lifeform</p> <p><input type="checkbox"/> Herbaceous <input type="checkbox"/> Shrub <input type="checkbox"/> Tree</p> <p style="margin-left: 100px;">Fuel Model</p>	<p>Indicator Species and Canopy Position</p>	<p>Structure Data (for upper layer lifeform)</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th style="text-align: center;"><i>Min</i></th> <th style="text-align: center;"><i>Max</i></th> </tr> </thead> <tbody> <tr> <td>Cover</td> <td style="text-align: center;">%</td> <td style="text-align: center;">%</td> </tr> <tr> <td>Height</td> <td></td> <td></td> </tr> <tr> <td>Tree Size Class</td> <td colspan="2"></td> </tr> </tbody> </table> <p><input type="checkbox"/> Upper layer lifeform differs from dominant lifeform.</p>		<i>Min</i>	<i>Max</i>	Cover	%	%	Height			Tree Size Class		
	<i>Min</i>	<i>Max</i>												
Cover	%	%												
Height														
Tree Size Class														

Description

Disturbances

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

Fire Regime Group:** IV

Historical Fire Size (acres)

Avg 250

Min 10

Max 10000

Sources of Fire Regime Data

- Literature
- Local Data
- Expert Estimate

Additional Disturbances Modeled

- Insects/Disease
- Native Grazing
- Other (optional 1)
- Wind/Weather/Stress
- Competition
- Other (optional 2)

Fire Intervals	<i>Avg FI</i>	<i>Min FI</i>	<i>Max FI</i>	<i>Probability</i>	<i>Percent of All Fires</i>
<i>Replacement</i>	92	30	120	0.01087	89
<i>Mixed</i>	714			0.00140	11
<i>Surface</i>					
<i>All Fires</i>	81			0.01228	

Fire Intervals (FI):

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class.

References

Brown, J.K. and J. Kapler-Smith, eds. 2000. Wildland fire in ecosystems: effects of fire on flora. Gen. Tech. Rep. RMRS-GTR-42-vol. 2. Ogden, UT: USDA Forest Service, Rocky Mountain Research Station. 257 pp.

Freeman, D.C, W.A. Turner, E.D. McArthur and J.H. Graham. 1991. Characterization of a narrow hybrid zone between two subspecies of big sagebrush (*Artemisia tridentata*: Asteraceae). American Journal of Botany. 78(6): 805-815.

Houston, D.B. 1973. Wildfires in northern Yellowstone National Park. Ecology 54: 1111-1117.

Howard, J.L. 1999. *Artemisia tridentata* spp. *wyomingensis*. In: Fire Effects Information System [Online]. USDA Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). Available: <http://www.fs.fed.us/database/feis>. Accessed 23 February 2005.

McArthur, E.D., D.C. Freeman and J.H. Graham. 1998. Narrow hybrid zone between two subspecies of big sagebrush (*Artemisia tridentata*: Asteraceae). VI. Respiration and water potential. Canadian Journal of Botany. 76(4): 567-574.

McArthur, E.D. 2000. Sagebrush systematics and distribution. Pages 9-14. in: P.G. Entwistle, A.M. DeBolt, J.H. Kaltenecker and K. Steenhof, compilers. Proceedings: Sagebrush Steppe Ecosystems Symposium. Bureau of Land Management Publication No. BLM/ID/PT-001001+1150, Boise, Idaho, USA.

Miller, R.F. and L.L. Eddleman. 2000. Spatial and temporal changes of sage grouse habitat in the sagebrush biome. Oregon State University Agricultural Experiment Station Technical Bulletin 151, Corvallis, Oregon. 35 pp.

NatureServe. 2007. International Ecological Classification Standard: Terrestrial Ecological Classifications. NatureServe Central Databases. Arlington, VA. Data current as of 10 February 2007.

Peters, E.F. and S.C. Bunting. 1994. Fire conditions pre- and post-occurrence of annual grasses on the Snake River plain. Pages 31-36. in Proceedings - Ecology, management, and restoration of Intermountain

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

rangelands symposium. USDA Forest Service INT-GTR-313, Ogden, Utah.

Welch, B.L. and C. Criddle. 2003. Countering Misinformation Concerning Big Sagebrush. Research Paper RMRS-RP-40. Ogden, UT: USDA Forest Service, Rocky Mountain Research Station. 28 pp.

West, N.E. 1983. Western Intermountain sagebrush steppe. Pages 351-395. in: N.E. West, ed. Ecosystems of the World 5: Temperate deserts and semi-deserts. Elsevier Scientific Publishing Co., New York, NY.

West, N.E. 2000. Synecology and disturbance regimes of sagebrush steppe ecosystems. Pages 15-26. in: P.G. Entwhistle, A.M. DeBolt, J.H. Kaltenecker and K. Steenhof, compilers. Proceedings: Sagebrush Steppe Ecosystems Symposium. Bureau of Land Management Publication No. BLM/ID/PT-001001+1150, Boise, Idaho, USA.

Whisnant, S.G. 1990. Changing fire frequencies on Idaho's Snake River plains: Ecological and management implications. Pages 4-10 in E.D. McArthur, E.M. Romme, S.D. Smith and P.T. Tueller, eds. Proceedings of a symposium on cheatgrass invasion, shrub die-off, and other aspects of shrub biology and management. U.S. Forest Service Gen. Tech. Rep. INT-276. Intermountain Forest and Range Experiment Station, Ogden, Utah.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.