

LANDFIRE Biophysical Setting Model

Biophysical Setting 3310800

Inter-Mountain Basins Big Sagebrush Shrubland

- This BPS is lumped with: 11250
 This BPS is split into multiple models:

General Information

Contributors (also see the Comments field) **Date** 10/27/2004

Modeler 1 Joe Vinyard joe_vinyard@co.blm.gov **Reviewer** Vic Ecklund vecklund@csu.org
Modeler 2 Ken Holsinger ken_holsinger@co.blm.gov **Reviewer** Chuck Kostecka kostecka@webaccess.net
v
Modeler 3 anonymous **Reviewer**

<u>Vegetation Type</u>	<u>Dominant Species</u>	<u>Map Zone</u>	<u>Model Zone</u>	
Upland Shrubland	ARTR2 JUNIP	33	<input type="checkbox"/> Alaska	<input checked="" type="checkbox"/> Northern Plains
<u>General Model Sources</u>	PURSH		<input type="checkbox"/> California	<input type="checkbox"/> N-Cent. Rockies
<input checked="" type="checkbox"/> Literature	SAVE4		<input type="checkbox"/> Great Basin	<input type="checkbox"/> Pacific Northwest
<input type="checkbox"/> Local Data	ATRIP		<input type="checkbox"/> Great Lakes	<input type="checkbox"/> South Central
<input checked="" type="checkbox"/> Expert Estimate	ERNA10		<input type="checkbox"/> Hawaii	<input type="checkbox"/> Southeast
	CHVI8		<input type="checkbox"/> Northeast	<input type="checkbox"/> S. Appalachians
	SYOR2			<input type="checkbox"/> Southwest

Geographic Range

Mid elevation of the Central Rockies through MT, Mountain areas of UT and northwest NM/northeast AZ.

Biophysical Site Description

This vegetation type is found on all aspects. Pure stands are found in areas with deeper soils and less topographic relief, but it is also common on slopes with a gradual shift to a mixed mountain shrub community on steeper slopes and in drainages. Elevation ranges from 1500-2300m (5000-7600ft), and precipitation from 11-20in. Soils are deep, well drained with a pH +/- 7.0. Soil moistures are udic (not dry for as long as 90 cumulative days) and soil temperatures cryic (very cold soils of the Rocky Mountain Region).

Vegetation Description

Dominant shrubs include *Artemisia tridentata* ssp. *vaseyana*, *Purshia tridentata* and *Symphoricarpos oreophilus*. Other common shrubs include *Amelanchier alnifolia*, *Chrysothamnus viscidiflorus*, *Cercocarpus montanus*, *Tetradymia canescens* and *Artemisia novae*. Other shrubs may be locally common. Herbaceous cover is moderate to abundant ranging from 40-85%. Common grasses include: *Festuca idahoensis*, *Elymus elymoides*, *Pascopyrum smithii*, *Elymus trachycaulus*, *Hesperostipa comata*, *Nassella viridula*, *Poa fenderiana* and *Poa juncifolia* var *ampla*. Indicative forbs include *Eriogonum umbellatum*, *Antennaria rosae*, *Balsamorhiza sagittata*, *Lupinus argenteus*, *Delphinium nuttallianum*, *Phlox multiflora* and *Viola nuttallii*.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

Disturbance Description

Mountain sagebrush steppe dominated by mountain sage, western snowberry and bitterbrush with a grass and forb understory is believed to be the major presettlement vegetation type for the area, although the exact composition of the community before settlement is unknown. Fire is a major disturbance factor for mountain sage (Blaisdell et al 1984, FEIS Database). Mountain big sagebrush has the fastest recovery rate of the three subspecies, may be as short as 15yrs (see FEIS, local data from various monitoring groups - NPS, BLM, TNC, etc). Fire size for this type is larger than other big sagebrush species because of greater fine fuel load but some unburned pockets remain. The fire return intervals reported in the literature for this sage type is 50yrs or more (Welch and Criddle 2003). Assuming that recovery rates are correlated with composite fire return intervals, one could posit with some certainty that the fire return interval lies somewhere between 40-75yrs. Ranges lie between 30yrs near Ponderosa Pine communities and other productive sites (maintaining more early seral types) up to 100yrs on north aspects and on rocky slopes (maintaining more late seral types).

Adjacency or Identification Concerns

Differentiation of mountain big sagebrush steppe from Wyoming big sagebrush may be difficult at the ecotone due to physical similarities and hybridization zones (i.e., species concepts become blurred).

Native Uncharacteristic Conditions

Scale Description

Size of disturbance extent will be limited by the variation of topographical features, age classes of the sage over the landscape, and vegetation types, all typical of mountain terrain. Average patch size 100-500ac with larger sizes during drought.

Issues/Problems

Reviewer questions existence of mixed severity component in this model.

Comments

NOTE 2/12/08: As a result of final QC for LANDFIRE National by Jennifer Long it was noticed that the time since disturbance function in the VDDT model for BpS 1080 was not attributed appropriately. According to LANDFIRE rules, time since disturbance should be set to less than or equal to one less than the class duration. This ensures that the time since disturbance function occurs before a deterministic transition shifts pixels to another class. In the case of BpS 1080 in MZ33, time since disturbance in class C was set to 49, although according to LANDFIRE rules it should have been set less than or equal to 33 because the duration of class C was 34yrs. When time since disturbance is set to 49 in class C as originally modeled the percent of the landscape in classes A-D are 15, 30, 25 and 30 respectively, as shown in this description document. When time since disturbance is set to 33 in class C, as it should have been attributed according to LANDFIRE rules, the percent of the landscape in classes A-D changes to 15, 45, 20 and 25 respectively. Based on the assumption that the modelers achieved the results they wanted with the time since disturbance function set to 49 in class C, it was decided to leave it as is. For future use, this model should be reassessed.

This model was copied from Z28 by Brendan C. Ward, July 18, 2007.

Based on the Rapid Assessment model R3MASB, which was reviewed by Bill Baker (bakerwl@wyo.edu) and Tim Christiansen (christta@wsmr.army.mil).

Mike Babler (mbabler@tnc.org) made species and other edits to match NatureServe description. R3MASB

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

was adapted from ROSBMT, reflects drier climate and longer fire return intervals observed in Southwest Region.

Final quality control (Pohl 8/17/05) resulted in the removal of a VDDT rule violation and slight adjustments (5%) to resulting proportions in classes C and D.

Based on the Rapid Assessment model R3MASB, which was reviewed by Bill Baker (bakerwl@wyo.edu) and Tim Christiansen (christta@wsmr.army.mil).

Vegetation Classes

Class A 15 %		<u>Indicator Species and Canopy Position</u>	<u>Structure Data (for upper layer lifeform)</u>	
			<i>Min</i>	<i>Max</i>
Early Development 1 All Structure		ARTR2	<i>Cover</i>	0 %
<u>Upper Layer Lifeform</u>		Upper	<i>Height</i>	Shrub 0m
<input type="checkbox"/> Herbaceous		PURSH	<i>Tree Size Class</i>	None
<input checked="" type="checkbox"/> Shrub		Upper	<input type="checkbox"/> Upper layer lifeform differs from dominant lifeform.	
<input type="checkbox"/> Tree	<u>Fuel Model</u>	ERNA10		
	2	Upper		
		CHVI8		
		Upper		
<u>Description</u>				
Sagebrush cover ranges from 0-5%. Herbaceous cover is variable, but is typically at least 30%.				
Class B 30 %		<u>Indicator Species and Canopy Position</u>	<u>Structure Data (for upper layer lifeform)</u>	
			<i>Min</i>	<i>Max</i>
Late Development 1 Closed		ARTR2	<i>Cover</i>	31 %
<u>Upper Layer Lifeform</u>		Upper	<i>Height</i>	Shrub 0.6m
<input type="checkbox"/> Herbaceous		PURSH	<i>Tree Size Class</i>	None
<input checked="" type="checkbox"/> Shrub		Upper	<input type="checkbox"/> Upper layer lifeform differs from dominant lifeform.	
<input type="checkbox"/> Tree	<u>Fuel Model</u>	ERNA10		
	6	Upper		
		CHVI8		
		Upper		
<u>Description</u>				
Sagebrush cover is >30%. Predominant grass/forb species will vary across geographic area.				
Class C 25 %		<u>Indicator Species and Canopy Position</u>	<u>Structure Data (for upper layer lifeform)</u>	
			<i>Min</i>	<i>Max</i>
Mid Development 1 Open		ARTR2	<i>Cover</i>	11 %
<u>Upper Layer Lifeform</u>		Upper	<i>Height</i>	Shrub 0.6m
<input type="checkbox"/> Herbaceous		PURSH	<i>Tree Size Class</i>	None
<input checked="" type="checkbox"/> Shrub		Upper	<input type="checkbox"/> Upper layer lifeform differs from dominant lifeform.	
<input type="checkbox"/> Tree	<u>Fuel Model</u>	ERNA10		
	6	Upper		
		CHVI8		
		Upper		
<u>Description</u>				

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

Sagebrush cover ranges from 5-15%. Predominant grass/forb species will vary across geographic area.

<p>Class D 30 %</p> <p>Late Development 1 Open</p> <p>Upper Layer Lifeform</p> <p><input type="checkbox"/> Herbaceous <input checked="" type="checkbox"/> Shrub <input type="checkbox"/> Tree</p> <p style="text-align: right;">Fuel Model 6</p>	<p>Indicator Species and Canopy Position</p> <p>ARTR2 Upper PURSH Upper ERNA10 Upper CHVI8 Upper</p>	<p>Structure Data (for upper layer lifeform)</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th style="text-align: center;">Min</th> <th style="text-align: center;">Max</th> </tr> </thead> <tbody> <tr> <td>Cover</td> <td style="text-align: center;">11 %</td> <td style="text-align: center;">30 %</td> </tr> <tr> <td>Height</td> <td style="text-align: center;">Shrub 1.1m</td> <td style="text-align: center;">Shrub 3.0m</td> </tr> <tr> <td>Tree Size Class</td> <td colspan="2" style="text-align: center;">None</td> </tr> </tbody> </table> <p><input type="checkbox"/> Upper layer lifeform differs from dominant lifeform.</p>		Min	Max	Cover	11 %	30 %	Height	Shrub 1.1m	Shrub 3.0m	Tree Size Class	None	
	Min	Max												
Cover	11 %	30 %												
Height	Shrub 1.1m	Shrub 3.0m												
Tree Size Class	None													
<p>Description</p> <p>Sagebrush cover ranges from 15-30%. Predominant grass/forb species will vary across geographic area.</p>														

<p>Class E 0 %</p> <p>[Not Used] [Not Used]</p> <p>Upper Layer Lifeform</p> <p><input type="checkbox"/> Herbaceous <input type="checkbox"/> Shrub <input type="checkbox"/> Tree</p> <p style="text-align: right;">Fuel Model</p>	<p>Indicator Species and Canopy Position</p>	<p>Structure Data (for upper layer lifeform)</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th style="text-align: center;">Min</th> <th style="text-align: center;">Max</th> </tr> </thead> <tbody> <tr> <td>Cover</td> <td style="text-align: center;">%</td> <td style="text-align: center;">%</td> </tr> <tr> <td>Height</td> <td></td> <td></td> </tr> <tr> <td>Tree Size Class</td> <td colspan="2"></td> </tr> </tbody> </table> <p><input type="checkbox"/> Upper layer lifeform differs from dominant lifeform.</p>		Min	Max	Cover	%	%	Height			Tree Size Class		
	Min	Max												
Cover	%	%												
Height														
Tree Size Class														

Description

Disturbances

<p>Fire Regime Group**: III</p> <p>Historical Fire Size (acres)</p> <p>Avg 0 Min 0 Max 0</p> <p>Sources of Fire Regime Data</p> <p><input type="checkbox"/> Literature <input type="checkbox"/> Local Data <input checked="" type="checkbox"/> Expert Estimate</p> <p>Additional Disturbances Modeled</p> <p><input checked="" type="checkbox"/> Insects/Disease <input checked="" type="checkbox"/> Native Grazing <input type="checkbox"/> Other (optional 1) <input checked="" type="checkbox"/> Wind/Weather/Stress <input type="checkbox"/> Competition <input type="checkbox"/> Other (optional 2)</p>	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">Fire Intervals</th> <th style="text-align: center;">Avg FI</th> <th style="text-align: center;">Min FI</th> <th style="text-align: center;">Max FI</th> <th style="text-align: center;">Probability</th> <th style="text-align: center;">Percent of All Fires</th> </tr> </thead> <tbody> <tr> <td>Replacement</td> <td style="text-align: center;">95</td> <td></td> <td></td> <td style="text-align: center;">0.01053</td> <td style="text-align: center;">76</td> </tr> <tr> <td>Mixed</td> <td style="text-align: center;">300</td> <td></td> <td></td> <td style="text-align: center;">0.00333</td> <td style="text-align: center;">24</td> </tr> <tr> <td>Surface</td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td>All Fires</td> <td style="text-align: center;">72</td> <td></td> <td></td> <td style="text-align: center;">0.01387</td> <td></td> </tr> </tbody> </table> <div style="border: 1px solid black; padding: 5px; margin-top: 10px;"> <p>Fire Intervals (FI): Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class.</p> </div>	Fire Intervals	Avg FI	Min FI	Max FI	Probability	Percent of All Fires	Replacement	95			0.01053	76	Mixed	300			0.00333	24	Surface						All Fires	72			0.01387	
Fire Intervals	Avg FI	Min FI	Max FI	Probability	Percent of All Fires																										
Replacement	95			0.01053	76																										
Mixed	300			0.00333	24																										
Surface																															
All Fires	72			0.01387																											

References

Blaisdell, J.P., R.B. Murray and E.D. McArthur. 1982. Managing Intermountain rangelands--sagebrush-grass ranges. Gen. Tech. Rep. INT-134. Ogden, UT: USDA Forest Service, Intermountain Forest and Range Experiment Station. 41 pp.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.

Blaisdell, J.P. and R.C. Holmgren. 1984. Managing intermountain rangelands-salt-desert shrub ranges. General Technical Report INT-163. Ogden, UT: USDA Forest Service, Intermountain Forest and Range Experiment Station. 52 pp.

Bunting, S.C. 2002. Natural and prescribed fires in big sagebrush steppe -- response of individual species and implication to burning practices. In: Restoration and management of sagebrush/grass communities workshop. Elko, NV. 4-8 Nov. 2002. Available: http://www.rangenet.org/trader/2002_Elko_Sagebrush_Conf.pdf.

Bunting, S.C., B.M Kilgore and C.L. Bushey. 1987. Guidelines for prescribed burning sagebrush-grass rangelands in the northern Great Basin. Gen. Tech. Rep. INT-231. Ogden, UT: USDA Forest Service. 33 pp.

Harniss, Roy O.; Murray, Robert B. 1973. 30 years of vegetal change following burning of sagebrush-grass range. *Journal of Range Management*. 26(5): 322-325.

Hironaka, M.; Fosberg, M.A. and Winward, A.H. 1983. Sagebrush-grass habitat types of southern Idaho. Bulletin Number 35. Moscow, ID: University of Idaho, Forest, Wildlife and Range Experiment Station. 44 pp.

Johnson, Kathleen A. 2000. *Artemisia tridentata* spp. *Vaseyana*. In: Fire Effects Information System, [Online]. USDA Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). Available: <http://www.fs.fed.us/database/feis/> [Accessed 2004, September 28].

Welch, B.L. and C. Criddle. 2003. Countering Misinformation Concerning Big Sagebrush. Research Paper RMRS-RP-40. Ogden, UT: USDA Forest Service, Rocky Mountain. Research Station. 28 pp.

Wright, H.A., L.F. Neuenschwander and C.M. Britton. 1979. The role and use of fire in sagebrush-grass and pinyon-juniper plant communities. Gen. Tech. Rep. INT-GTR-58. Ogden, UT: USDA Forest Service, Intermountain Research Station. 48 pp.

**Fire Regime Groups are: I: 0-35 year frequency, surface severity; II: 0-35 year frequency, replacement severity; III: 35-100+ year frequency, mixed severity; IV: 35-100+ year frequency, replacement severity; V: 200+ year frequency, replacement severity.