

How Can We Use VCIS?

- Local, state, tribal, or federal managers of smoke or other pollutants can use the ventilation index data.
- Maps and graphs of historical ventilation conditions can be:
 - ❖ included in fire- or smoke-management plans.
 - ❖ used to illustrate concerns in public meetings.
 - ❖ overlain with other values at risk.
- The data can be used to help develop plans for avoiding smoke impacts and optimizing the use of prescribed fire at specific places or certain times of the year.
- Spatial patterns of risks to air quality and visibility are illustrated in maps of monthly averaged values.
- Frequency statistics at selected points show temporal dimension of risk.
- Values at risk can be assessed for specific airsheds and compared over time or with other airsheds.

VCIS offers the first quantitative planning tool for smoke management that is nationally consistent.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, or marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program GET Center, 202-720-2600, voice and TDD.


To file a complaint of discrimination, write: USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue SW, Washington, DC 20250-9410, or call 202-720-5964, voice and TDD. USDA is an equal opportunity provider and employer.

What is VCIS?

The ventilation climate information system (VCIS) allows users to assess risks to values of air quality and visibility from historical patterns of ventilation conditions.

It is available through an interactive, Internet map server. The Internet server allows maps of ventilation potential to be overlain with sensitive receptors, terrain features, or political boundaries. The data apply to local, regional, or national scales.

Historical patterns of ventilation conditions are available through an interactive, Internet map server.



Average ventilation index for February afternoon with sensitive receptors (hospitals, airports, and roads).



Fire and Environmental
Research Applications Team

Contact information:

Sue A. Ferguson (206) 732-7800
sferguson@fs.fed.us

USDA Forest Service
Pacific Northwest Research Station
Fire and Environmental Research Applications Team
4043 Roosevelt Way NE
Seattle, WA 98105

VENTILATION CLIMATE INFORMATION SYSTEM

***A Management Tool
for Smoke and other
Pollutants***



The VCIS is available at
<http://www.fs.fed.us/pnw/fera/vent>




Joint Fire
Science Program

March 2002

What's Inside VCIS?


The ventilation climate information system (VCIS) is based on a 40-year database that includes twice-daily values of wind, mixing height, and a ventilation index that is the product of wind speed and mixing height. Data are spatially interpolated to a grid of about 5 kilometers on a side.

The VCIS offers the first nationally consistent maps of surface wind and ventilation index ...



Average ventilation index for June morning.

...and includes the longest climate record of mixing height in the country.




Frequency of mixing height over a point near Honolulu.


The Science Behind VCIS

Wind: Surface winds were generated with a reliable, well-documented meteorological model. The best available upper-air data were used as inputs to the model. Surface observations were used to verify the model. At every observation point, users can evaluate model performance and estimate uncertainty for themselves.

Mixing Height: A standard parcel method was used to calculate mixing height from upper air and surface observations. Mapped values that intersect high terrain were adjusted to well above ground level.



Mixing height.




Mixing height adjusted for local inversions.

A unique set of algorithms were used to derive the likely timing and location of local nighttime inversions. Mapped values of mixing height were adjusted to account for local inversions wherever and whenever they occur.

Ventilation Index: The surface wind speed times the mixing height gives an estimate of ventilation potential. The resulting index in VCIS is lower than is typical because it is influenced by local inversions and derived from winds at the surface instead of higher in the mixed layer.

Table 1—Range of ventilation indices corresponding to each ventilation class

Ventilation index Meters per second	Class
0-1175	Poor
1176-2350	Marginal
2351-3525	Fair
>3526	Good



Frequency of ventilation index over a point in southeast Nebraska.