

REPORT: *ARCANGELIELLA CLARIDGEI* COMB. NOV. AND *A. CORKII* COMB. NOV. TRANSFERRED FROM THE GENUS *ZELLEROMYCES*

James M. Trappe¹ and Andrew W. Claridge²

¹Department of Forest Science, Oregon State University, Corvallis, Oregon 97331-5752, USA. Email: trappej@onid.orst.edu

²Department of Environment and Climate Change, Parks and Wildlife Division, Reserve Conservation Unit, Southern Branch, PO Box 2115, Queanbeyan, NSW 2620, Australia

Abstract

J. M. Vidal's recent reexamination of the holotype of *Arcangeliella borziana* Cavara, the type species of that genus, revealed that it has been misinterpreted by various authors in the past. Vidal synonymised the genus *Zelleromyces* with *Arcangeliella* and transferred all *Zelleromyces* spp. to *Arcangeliella* except for two Australian species, *Z. claridgei* and *Z. corkii*. We complete his transfers here with the new combinations *Arcangeliella claridgei* and *A. corkii*.

J.M. Trappe and A.W. Claridge (2008). Report: *Arcangeliella claridgei* comb. nov. and *A. corkii* comb. nov. transferred from the genus *Zelleromyces*. *Australasian Mycologist* 27 (2): 117.

The genus *Arcangeliella* (Russulaceae) was interpreted by Singer and Smith (1960) to include all *Ladarius*-related, sequestrate species with a columella. They assigned species lacking a columella to their new genus, *Zelleromyces*. However, the type species of *A. borziana* Cavara, as illustrated by Cavara (1900) and recently studied by Vidal (2004), has only a narrow central strand that may be percurrent but can disappear as specimens mature. *Zelleromyces* species abound in Australia, and we have observed that some specimens of a given species may have a rudimentary columella, whereas others do not. We agree with Vidal's conclusion that presence or absence of a columella in astipitate species are not useful taxonomic characters at the generic level and that all *Zelleromyces* spp. should be transferred to *Arcangeliella*. Vidal (2004) did this except for *Z. claridgei* and *Z. corkii*, published by us (Trappe and Claridge 2003) shortly before his paper appeared. Accordingly, we propose these new combinations here:

Arcangeliella claridgei (Trappe in Trappe & Claridge) Trappe & Claridge, comb. nov.

Basionym: *Zelleromyces claridgei* Trappe in Trappe & Claridge, *Australasian Mycologist* 22, 34.2003.

Arcangeliella corkii (Trappe & Claridge) Trappe & Claridge, comb. nov.

Basionym: *Zelleromyces corkii* Trappe & Claridge, *Australasian Mycologist* 23, 35-36. 2003.

Acknowledgements

This study was supported in part by CSIRO Sustainable Ecosystems, Gungahlin, ACT, Australia.

References

- Cavara, F. (1900). *Arcangeliella borziana* nov. gen., nov. sp. Nuova imenogasterea delle abetine di Vallombrosa. *Nuovo Giornale Botanico Italiano, Nuova Serie* 7, 117-128.
- Singer, R. & Smith, A.H. (1960). Studies on secotiaceous fungi IX. The astrogastreaous series. *Memoirs of the Torrey Botanical Club* 21, 1-112.
- Trappe, J.M. & Claridge, A.W. (2003). Australasian sequestrate (truffle-like) fungi. 15. New species from tree line in the Australian Alps. *Australasian Mycologist* 22, 27-38.
- Vidal, J.M. (2004). *Arcangeliella borziana* and *A. stephensii*, two gasteroid fungi often mistaken. A taxonomic revision of *Ladarius*-related sequestrate fungi. *Revista Catalana de Micologia* 26, 59-82.