



FIG. 1. *Aneides ferreus* inside a cavity 75 m above ground in the top of an old-growth Douglas-fir (*Pseudotsuga menziesii*), Coquille River, Oregon, USA.

fir forests than is generally known and also support the hypothesis that nests of arboreal rodents may be important microhabitats for *A. ferreus* (Spickler et al. 2006, *op. cit.*; Forsman and Swingle 2007, *op. cit.*). It remains unknown if use of the forest canopy by *A. ferreus* is restricted to foraging and shelter, or includes occasional breeding as well.

Submitted by WILLIAM W. PRICE, 1110 SW 197th Avenue, Beaverton, Oregon 97006, USA (e-mail: groupb@gmail.com); CLINTON P. LANDON, P.O. Box 2049, Clackamas, Oregon 97015, USA (e-mail: cplandon@gmail.com); and ERIC D. FORSMAN, U.S. Forest Service, Pacific Northwest Research Station, 3200 SW Jefferson Way, Corvallis, Oregon 97331, USA (e-mail: eforzman@fs.fed.us).

CRYPTOBRANCHUS ALLEGANIENSIS ALLEGANIENSIS (Eastern Hellbender). SECRETION PRODUCTION. Of the potential antipredator mechanisms exhibited by amphibians, noxious skin secretions are considered the most effective against potential predators (Brodie et al. 1979. *Copeia* 1979:270-274). Many species produce toxic and irritating skin secretions as adults, but the larvae of many salamanders are palatable to various predators because toxic and distasteful secretions generally do not develop until after metamorphosis (Formanowicz and Brodie 1982. *Copeia*



FIG. 1. Larval Eastern Hellbenders (*Cryptobranchus alleganiensis alleganiensis*) immediately after producing a white secretion composed of sticky and foamy components.

1982:91-97).

Adult *Cryptobranchus alleganiensis* are large amphibians (60+ cm) and probably have few predators. When stressed or being captured, adult hellbenders often produce a milky secretion that is bitter and distasteful when applied to the tongue (Brodie 1971. *Herpetol. Rev.* 3:8), and the secretion may be unpalatable to predators. Larval hellbenders hatch between 23 and 30 mm total length and metamorphose 1.5-2 yrs after hatching, and are probably highly vulnerable to predation due to their small size and slow developmental rate (Nickerson and Mays 1973. *The Hellbenders*. Milwaukee Public Museum. Wisconsin. 106 pp.). It is unknown when the ability to produce secretion develops in hellbenders. In Oct-Nov 2007 several Eastern Hellbender egg clutches were collected in Missouri for captive rearing. On 29 April 2008 at 0900 h, 12 larval hellbenders (25 weeks post hatching, mean TL \pm SE = 91.75 \pm 1.8 mm) were collected in a small net for transport to a separate container for behavioral observations. The larvae immediately produced copious amounts of a secretion that appears to have two components. The first component was water-soluble and had a "foamy soap" appearance (Fig. 1). The second component was very sticky and was not soluble in water (Fig. 1); it remained adhered to the individual hellbenders for up to 48 h. Both components are similar in appearance to those produced by adult hellbenders (Nickerson and Mays 1973, *op. cit.*).

Immediately after secretion production, BGG put a small amount of the foamy secretion on his tongue and experienced a strong bitter sensation lasting for ca. 5 sec. The sensation was not accompanied with burning or numbness. The larval secretion tasted very similar to that produced by adult hellbenders.

Our observations suggest that, unlike most salamanders with aquatic larvae, larval hellbenders are capable of producing noxious skin secretions that might function to deter potential predators.

Submitted by BRIAN G. GALL, 5305 Old Main Hill, Logan, Utah 84322, USA (e-mail: gall@biology.usu.edu); ADAM L. CRANE (e-mail: AdamCrane@MissouriState.edu) and ALICIA MATHIS (e-mail: AliciaMathis@MissouriState.edu), 901 S. National, Springfield, Missouri 65897, USA

- and Conservation of Amphibians and Reptiles. pp. 391-395. SSAR Contrib. Herpetol. II.
- HAGMAN, M., AND R. SHINE. 2008. Deceptive digits: the functional significance of toe waving by cannibalistic cane toads, *Chaunus marinus*. Anim. Behav. 75(1):123-131.
- HOOGMOED, M. S. 1967. Mating and early development of *Gastrotheca marsupiata* (Dumeril and Bibron) in captivity (Hylidae. Anura, Amphibia). Brit. J. Herpetol. 4:1-7.
- IUCN 2009. 2009 IUCN Red List of Threatened Species. <www.iucnredlist.org>. Accessed on 03 July 2009.
- LIPS, K. R., F. BREM, R. BRENES, J. D. REEVE, R. A. ALFORD, J. VOYLES, C. CAREY, L. LIVO, A. P. PESSIER, AND J. P. COLLINS. 2006. Emerging infectious disease and the loss of biodiversity in a Neotropical amphibian community. Proc. Natl. Acad. Sci. 103:3165-3170.
- P. A. BURROWES, J. R. MENDELSON III, AND G. PARRA-OLEA. 2005. Amphibian population declines in Latin America: a synthesis. Biotropica 37:222-226.
- MARTINA, AND G. F. WATSON. 1971. Life history as an aid to generic delimitation in the family Hylidae. Copeia 1971 :78-89.
- MATTHEWS, L. H. 1957. Viviparity in *Gastrotheca* (Amphibia: Anura) and some considerations on the evolution of viviparity. Bull. Soc. Zool. France 82:317-320.
- MEANS, D. E., W. E. DUELLMAN, AND V. CLARK. 2008. Ovipositing behavior in the egg-brooding frog *Stefania ayanganae* (Anura, Hemiphractidae). Phyllomedusa 7(2):143-148
- MENDELSON III, J. R., H. R. DA SILVA, AND A. M. MAGLIA. 2007. Phylogenetic relationships among marsupial frog genera (Anura: Hylidae: Hemiphractinae) based on evidence from morphology and natural history. Zool. J. Linn. Soc. 128:125-148.
- MURPHY, J. B. 1976. Pedal luring in the leptodactylid frog, *Ceratophrys calcarata* Boulenger. Herpetologica 32:339-341
- QUGUANGO-UBILLUS, A., AND L. A. COLOMA. 2008. Notes on behaviour, communication, and reproduction in captive *Hyloxalus toachi* (Anura: Dendrobatidae), an endangered Ecuadorian frog. Int. Zoo. Yb. 42:78-89.
- RADCLIFFE, C. W., D. CHISZAR, K. ESTEP, J. B. MURPHY, AND H. M. SMITH. 1986. Observations on pedal luring and pedal movements in leptodactylid frogs. J. Herpetol. 20(3):300-306.
- SAVAGE, J. M. 2002. The Amphibians and Reptiles of Costa Rica: A Herpetofauna between Two Continents, between Two Seas. University of Chicago Press, Chicago, Illinois. 934 pp.
- SLOGGETT, J. J., AND I. ZEILSTRA. 2008. Waving or tapping? Vibrational stimuli and the general function of toe twitching in frogs and toads (Amphibia: Anura). Anim. Behav. 76(5):e1-e4.
- TODD, B. D. 2007. Parasites lost? An overlooked hypothesis for the evolution of alternative reproductive strategies in amphibians. Am. Nat. 170:793-799.
- WELDON, P. J., B. J. DEMETER, AND R. ROSSCOE. 1993. A survey of sheel skin-eating (dermatophagy) in amphibians and reptiles. J. Herpetol. 27(2):219-228.
- ZIPPEL, K. C. 2002. Conserving the Panamanian golden frog: Proyecto Rana Dorada. Herpetol. Rev. 33:11-12.
- , R. LACY, AND O. BYERS (eds.) 2006. CBSG/WAZA amphibian ex situ conservation planning workshop final report. IUCN/SSC Conservation Breeding Specialist Group. Apple Valley, Minnesota.

NATURAL HISTORY NOTES

The Natural History Notes section is analogous to Geographic Distribution Preferred notes should 1) focus on observations in the field, with little human intrusion; 2) represent more than the isolated documentation of developmental aberrations; and 3) possess a natural history perspective. Individual notes should, with few exceptions, concern only one species, and authors are requested to choose a keyword or short phrase which best describes the nature of their note (e.g., Reproduction, Morphology, Habitat, etc.). Use of figures to illustrate any data is encouraged, but should replace words rather than embellish them. The section's intent is to convey information rather than demonstrate prose. Articles submitted to this section will be reviewed and edited prior to acceptance.

Electronic submission of manuscripts is requested (as Microsoft Word or Rich Text format [rtf] files, as e-mail attachments), Figures can be submitted electronically as JPG files, although higher resolution TIFF or PDF files will be requested for publication. Please DO NOT send graphic files as imbedded figures within a text file. Additional information concerning preparation and submission of graphics files is available on the SSAR web site at: <http://www.ssarherps.org/HRinfo.html>. Manuscripts should be sent to the appropriate section editor: Marc P. Hayes (crocodilians, lizards, and *Sphenodon*; mhayesrana@aol.com); Charles W. Painter (amphibians: charles.painter@state.nm.us); Andrew T. Holycross (snakes; AndrewHolycross@gmail.com); and James Harding (turtles; hardingj@msu.edu).

Standard format for this section is as follows: SCIENTIFIC NAME. COMMON NAME (for the United States and Canada as it appears in Crother [ed.] 2008. *Scientific and Standard English Names of Amphibians and Reptiles of North America North of Mexico*. SSAR Herpetol. Circ. 37:1-84, available from SSAR Publications Secretary. ssar@herplito.com; for Mexico as it appears in Liner and Casas-Andrea 2008. *Standard Spanish, English and Scientific Names of the Amphibians and Reptiles of Mexico*. Herpetol. Circ. 38:1-162), KEYWORD. DATA on the animal. Place of deposition or intended deposition of specimen(s), and catalog number(s). Then skip a line and close with SUBMITTED BY (give name and address in full-s-spell out state names-no abbreviations). (NCN) should be used for common name where none is recognized. References may be briefly cited in text (refer to this issue for citation format).

Recommended citation for notes appearing in this section is: Medina, P., and R. L. Jøglar, 2008. *Eleutherodactylus richmondi*: reproduction. Herpetol. Rev. 39:460.

CAUDATA - SALAMANIDERS

ANEIDES FERREUS (Clouded Salamander). ARBOREAL ACTIVITY. *Aneides ferreus* inhabits the forests of western Oregon and extreme northwestern California. Although thought to be primarily *terrestrial*, *A. ferreus* has occasionally been found as high as 60 m up in trees (Jones et al. 2005. Amphibians of the Pacific Northwest. Seattle Audubon Society. 227 pp.), and two recent reports suggest that it may be more arboreal than previously believed (Spickler et al. 2006. Herpetol. Conserv. Biol. 1:16-26; Forsman and Swingle 2007. Herpetol. Conserv. Biol. 2:113-118). However, it is difficult to evaluate the amount of arboreal activity by this species because almost all sampling efforts have been focused on terrestrial habitats.

On 23 May 2008 near the Coquille River in southwest Oregon we observed two adult *Ai ferreus* in a cavity 75 m above ground in a Douglas-fir tree *tPseudotsuga menziesii* that was 78 m tall and 195 cm diameter at breast height (43.1797°N, 123.8122°W; WGS 84). Exposed when a limb broke off, the small cavity in the trunk contained copious amounts of Red Tree Vole (*Arborimus longicaudus*) fecal pellets. Two days later we photographed what we presumed to be the same two salamanders in the cavity (Fig. 1). Based on the shape of their heads, we suspected they were a male and female. These observations add to the increasing evidence that *A. ferreus* is more active in the canopy of Douglas-