

Partitioning hydrologic contributions to an ‘old-growth’ riparian area in the Huron Mountains of Michigan, USA[†]

Randall K. Kolka^{1*}, Christian P. Giardina², Jason D. McClure³, Alex Mayer³,
and Martin F. Jurgensen³

¹ Forestry Sciences Laboratory, Northern Research Station, USDA Forest Service, Grand Rapids, MN 55744, USA

² Institute of Pacific Islands Forestry, Pacific Southwest Research Station, USDA Forest Service, 60 Nowelo Drive, Hilo, Hawaii 96720, USA

³ Michigan Technological University, Houghton, MI 49931, USA

ABSTRACT

Over the past century, annual snowfall has increased across the ‘snow-belt’ region of the Upper Peninsula of Michigan, yet total annual precipitation has not changed, with potential impacts on hydrological processes and ecosystem composition. Using an integrated hydrochemical approach, we characterized groundwater discharge and quantified the contribution of snow- and rain-derived waters to groundwater for an old-growth riparian area within the Huron Mountains in northern Michigan. We then quantified the relative contribution of lateral, hillslope-derived groundwater and upstream lake water to streamwater, and the extent of hyporheic zone expansion and contraction during one growing season. During a period of above-average snowfall, yet below average growing season precipitation, ~80% of the riparian area’s groundwater reservoir was derived from snowmelt. The relative contribution of groundwater to streamflow ranged from 70% in June to 100% in August. The remainder was derived from upstream lakes and wetlands, which dropped in elevation and relative contribution from June to August. Finally, the extent of the hyporheic zone was small (<50 cm from streambed surface) and contracted towards the stream during the recession limb of the hydrograph. We conclude that if snowfall continues to rise while total annual precipitation declines, in line with climate change scenarios for the region, then water fluxes from snowmelt will increasingly dominate summer baseflow and could potentially increase spring flooding. However, because of overall declines in precipitation, streamflow patterns will likely change towards lower overall flow following the influence of the snowmelt period. Published in 2010 by John Wiley & Sons, Ltd.

KEY WORDS riparian; snowmelt; groundwater; hyporheic; stable isotopes

Received 11 October 2009; Accepted 21 December 2009

INTRODUCTION

The riparian area connects upland and in-stream ecosystems, where the amount and distribution of upland precipitation drive the recruitment, retention, and release of water and its constituents (carbon, nitrogen, etc.) to downstream ecosystems (Naiman and Decamps, 1997). Consequently, efforts to quantify patterns of subsurface and surface water movement through riparian areas are needed to inform simulation models that forecast the effects of chronic (e.g. global temperature increases) or acute (e.g. 100-year flood) climatic events on ecosystem processes (National Research Council, 2002; Naiman *et al.*, 2005).

Across the Upper Peninsula of Michigan, USA, annual precipitation falls nearly equally as snow and rain (Stottlemyer and Toczydowski, 1996). Some climate change scenarios predict an alteration in the amount and seasonality of precipitation inputs for the Lake Superior

region, where predicted increase in temperature leads to drier summers, yet increasing amounts of snow precipitation during winters (Kattenberg *et al.*, 1996; Kunkel *et al.*, 2000). In line with model predictions, there is evidence that annual snowfall in the region has increased over the past century (Norton and Bolsenga, 1993; Leathers and Ellis, 1996; Burnett *et al.*, 2003). Based on Houghton, MI annual records from 1890 to 2007 (<http://www.admin.mtu.edu/alumni/snowfall/>), snowfall has increased at 30 mm yr⁻¹ ($r^2 = 0.45$, $p < 0.01$); increases in snowfall from 1958 to 2007 for Marquette, Michigan is 64.8 mm yr⁻¹ ($r^2 = 0.33$, $p < 0.01$). For Marquette, there does not appear to be a change in total precipitation (Marquette’s $r^2 = 0.008$, $p = 0.58$), in line with climate change models that predict altered patterns of precipitation distribution for the region. The measured increase in ‘lake-effect’ snow within the Upper Peninsula is perceived to be due in part to greater rates of winter evaporation from the surface of Lake Superior, where rising annual temperatures of Lake Superior result in a longer duration of ice-free (evaporative) surface area during winter (Leathers and Ellis, 1996; Burnett *et al.*, 2003). Annual increases in snow precipitation lead

* Correspondence to: Randall K. Kolka, USFS Forestry Sciences Laboratory, 1831 Hwy 169 E, Grand Rapids, MN 55744.
E-mail: rkolka@fs.fed.us

[†] This article is a U.S. Government work and is in the public domain in the USA.

to greater amounts of spring snowmelt, which in turn can lead to elevated pulses of either 'acidic' and/or 'dilution' events that can affect local streams and lakes by altering stream chemistry and productivity (Johannessen and Henriksen, 1978; Rascher *et al.*, 1987; Stottlemyer and Toczydlowski, 1991). Furthermore, where annual precipitation amounts stay constant or even rise, but growing season precipitation decreases (i.e. more snow, less rain), then ecosystem composition, structure and function are likely to change in response to altered hydrological processes. Considering the numerous scenarios and implications for northern ecosystems under a changing climate (Davis *et al.*, 2000; Nijssen *et al.*, 2001), quantifying current connections between climate and forest hydrology in near-pristine north-temperate forests will provide an important baseline for quantifying future changes.

Overland flow and direct precipitation inputs are minor contributors to headwater streams in northern temperate forests, and so field-based hydrology studies in these systems have focused on riparian groundwater fluxes to streamflow (Morrice *et al.*, 1997; Walker and Krabbenhoft, 1998; McGlynn *et al.*, 1999; Vidon and Hill, 2004; Wondzell, 2005). Groundwater contributions of terrestrially derived nutrients and elements contribute to biological and chemical processes within the surface water environment (Hill, 2000), and the functional significance of riparian soils on nutrient cycling and flux regulation is well recognized (Mulholland, 1992; Cirimo and McDonnell, 1997; Schindler and Krabbenhoft, 1998; Baker *et al.*, 2000; Thomas *et al.*, 2001; Valett *et al.*, 2002; McClain *et al.*, 2003). Further, riparian groundwater provides a stable source of water to streams and transpiring vegetation. A change in precipitation inputs to groundwater could potentially alter riparian ecosystem processes, where the distribution (timing, duration, and amount) of inputs are likely to affect solute fluxes and concentrations.

In this study we used three approaches to quantify the water sources to a headwater stream in the Huron Mountain Reserve (HMR) in the Upper Peninsula of Michigan, USA. First, we compared upstream and downstream discharge to estimate groundwater contribution along a stream reach (Harvey and Wagner, 2000). Secondly, we measured the isotopic composition for oxygen (O^{18}/O^{16}) and hydrogen (H^2/H) within stream and source waters to partition ground- and stream-sources (Buttle, 1998; Cey *et al.*, 1998; Winograd *et al.*, 1998; McGlynn *et al.*, 1999; Burns *et al.*, 2001; Katsuyama *et al.*, 2001; Atekwana and Richardson, 2004; Pardo *et al.*, 2004; Wenninger *et al.*, 2004; Monteith *et al.*, 2006; Reddy *et al.*, 2006). Lastly, we used semi-conservative parameters of conductivity, temperature and chloride concentration to create a more complete description of groundwater hydrology (Christopher *et al.*, 1990; Hooper *et al.*, 1990; Stream Solute Workshop, 1990). This diverse hydrochemical approach was used to partition the hillslope's hydrological contribution to the downslope riparian area, stream channel, and hyporheic zone (McGlynn *et al.*, 1999; Burns *et al.*, 2001; Ladouche *et al.*, 2001; Seibert *et al.*, 2003; Wenninger *et al.*, 2004).

The Fisher Creek (FC) riparian area, where the current study was conducted, has no known history of management, including land clearing or harvesting for timber, and so is an important pristine or old-growth site for conducting baseline forest-hydrology research. We build on previous isotope-based hydrology studies of northern hardwood forests (Cey *et al.*, 1998; McGlynn *et al.*, 1999; Buttle *et al.*, 2001; Monteith *et al.*, 2006), by describing the baseflow conditions and relative contributions of an old-growth northern hardwood riparian area. Our objectives were to characterize the riparian subsurface hydrology for two old-growth riparian reaches. Specifically, we quantified groundwater discharge and the relative contributions of: (1) rainfall and snowfall to riparian groundwater; and (2) groundwater and upstream sources to streamwater. We also used the isotopic signature of groundwater and streamwater to examine the extent and dynamics of the hyporheic zone. We hypothesized that: (1) snowmelt would be the dominant source to groundwater during baseflow; (2) lateral groundwater inputs from the hillslope to streamwater would increase in relative contribution along the recession limb of the hydrograph, as longitudinal inputs from upstream sources decrease; and (3) the zone of hyporheic mixing would contract towards the stream during the recession limb of the hydrograph as upstream inputs and surface streamflow decreases.

METHODS

Site description

This study was conducted in the HMR (46°52' north latitude, 87°50' west longitude), a conservation area within the larger, privately owned Huron Mountain Club property in the Upper Peninsula of Michigan, USA near Lake Superior (Figure 1). The HMR contains one of the largest (~2600 ha) pristine old-growth forests within the Great Lakes region (Frelich, 1995; Woods, 2000; Flaspohler and Meine, 2006). The climate of the HMR is characterized by a relatively even distribution of annual precipitation, low potential evapotranspiration, and a strong local influence from the proximity to Lake Superior, which includes moderated temperatures and elevated snowfall (Figure 2; Denton and Barnes, 1988). Three National Oceanic and Atmospheric Administration (NOAA) weather stations located in Marquette, Houghton and Herman are within 55 km of the research site. Based on isohyetal interpolation, we estimate a 30-year average (1971–2000) for mean annual temperature of 4.2 °C, a mean precipitation of 918 mm, and a mean snowfall of 5453 mm for the HMR. Snow-free season rainfall data were collected daily from two locations approximately 3-km north and east of the FC using graduated rain collectors during the study period from April 7 to November 11, 2005 (total = 522 mm). To account for the spring snowmelt, snow water equivalent was found to be 0.72 mm mm⁻¹ (±0.112; *n* = 6) for late March, which for the average measured snowpack depth



Figure 1. Location of Huron Mountains, in the Upper Peninsula of Michigan, USA and close-up of Fisher Creek, Florence Pond Drain, and plot locations within the Huron Mountain Reserve.




Figure 2. Monthly precipitation and snowfall data for nearby Marquette, MI.

of approximately 400 mm was equal to about 290 mm of meltwater precipitation per unit area. Together, snowmelt plus rainfall, the localized 812 mm of precipitation corroborates the below average conditions for the study period.

Our study was conducted along Florence Pond Drain (FPD), a 1st order reach that drains Florence Pond, and FC, a 2nd order reach that drains Trout Lake and other small headwater wetlands (Figure 1). The two reaches were selected based on mature forest condition, and similarity in lithotopography (Montgomery, 1999) and channel characterization (Rosgen, 1994). The FC study area is approximately 600 m in straight length, with an approximate 840 m of stream channel length (watershed area above our monitoring location = 349 ha), while the FPD study area is approximately 300 m in length, with an approximate stream channel length of 540 m (watershed area above our monitoring location = 117 ha).

The riparian areas of FC and FPD have been characterized as a yellow birch (*Betula allegheniensis*), eastern hemlock (*Tsuga canadensis*), northern white cedar (*Thuja occidentalis*), red maple (*Acer rubrum*) forest type (Simpson *et al.*, 1989, 1990), although sugar maple (*Acer saccharum*), white pine (*Pinus strobus*), white spruce (*Picea glauca*), and balsam fir (*Abies balsamea*) also occur. The

canopy is composed of super-dominant individuals of white pine that often exceed 40 m in height and 1.2 m in diameter (Wells and Thompson, 1976), which attest to the absence of timber harvesting activities within the watershed. Analysis of variance (ANOVA) revealed that reaches did not differ with respect to amounts of large woody debris (volume, length, or biomass) or decay class frequency. Because characteristics of large woody debris within a riparian forest often is related to disturbance history (Hedman *et al.*, 1996; Bragg and Kershner, 1999; Naiman *et al.*, 2002; McClure *et al.*, 2004), we expect that the FC and FPD reaches to be similar in disturbance history.

The Huron Mountains consist of Huronian and Archean formations, which are of Precambrian metamorphic origin resulting from uplifting of the Canadian Shield (Dorr and Eschman, 2001). They were subject to glacial processes until ~10 000 ybp. Floodplains of HMR riparian areas consist of deep sandy glacial outwash sediment, and are characterized as sandy stream terraces with deep, moderately well drained loamy sands (Simpson, 1990). Soils are mapped as either Kalkaska Sands or the Evert–Pelkie–Sturgeon Complex, in which all soils are sands with hydraulic conductivity values that range from 0.000423–0.0141 cm s⁻¹ and pH values that range from 3.6–8.4, depending on depth. The higher pH Evert type is composed of 0–10% calcium carbonate, but when soils were sampled to a depth of 50 cm throughout the riparian area, application of 10% HCl did not result in a characteristic bubbling reaction, and so indicates little or no calcium carbonate.

Streamflow

Stream discharge was measured using velocity and cross-sectional area measurements throughout the two reaches. Velocity was measured with a Marsh-McBirney™ electromagnetic flow meter. Discharge was determined for multiple stage heights, while stage was recorded regularly using three-staff gauges positioned within the stream's thalweg at two locations along the FC reach and at one location within the FPD reach.

To predict discharge during periods when no field data were collected, a discharge-to-discharge relationship was developed using regression analysis against a nearby United States Geological Survey (USGS) stage recording data station (<http://waterdata.usgs.gov/nwis/rt>). A discharge-to-discharge power function was determined to be the best fit for FC and FPD discharge. Power functions have been used in similar studies of other regional watersheds (Cey *et al.*, 1998; Goebel, 2001). The Yellow Dog River USGS station located approximately 20 km from FC provided the best continuous discharge data with which to predict FC and FPD flow ($r^2 = 0.76$ for FC, $r^2 = 0.90$ for FPD; $p \leq 0.05$).

Well network

Throughout FC and FPD, 11-m segments of straight reaches were used as the upper and lower bound of each

plot. A total of 15 plots were randomly chosen from a population of 46 selected 11-m segments of similar morphological characteristics within the reaches. Each rectangular plot spans the farthest extent of the historical floodplain on either side of the stream (Figure 1). Within each plot, at least six wells were randomly located at: (1) the approximate bankfull width of the stream, and (2) the estimated floodprone width of the stream, where floodprone is considered the width of the stream at twice the depth of the thalweg. Beyond the floodprone wells on both sides of the stream, a floodplain well was installed at the farthest edge of each plot, which coincided with the farthest extent of the historical floodplain. Terrace wells were positioned above the floodplain ($n = 4$ in FC, $n = 2$ in FPD) irrespective of plot location. There were 11 plots positioned within the FC reach, with a total of 72-bankfull wells, 71-floodprone wells, and 22-floodplain wells. Within the FPD reach, a total of four plots, with 26-bankfull wells, 26-floodprone wells, and 8-floodplain wells were installed.

All wells were installed using a bucket-auger with each bankfull and floodprone well installed to approximately 30 cm below the stream bed elevation; floodprone and terrace wells were installed 2–3 m below ground-surface, to extend below the point of contact with the water table. All wells were constructed of 5.1-cm inside-diameter polyvinyl chloride (PVC) pipe of various lengths with perforations along the underground portion of the pipe. Each perforated section of the pipe was then covered with a nylon filter to keep sediment from entering the wells, while each pipe bottom and top was capped with a fitted PVC cap. All wells were back-filled with native material and capped with 5 cm of Portland cement. Each cap was then sealed with 5 cm of bentonite clay to resist preferential flow down the well exterior.

In addition, streambed wells (or mini-piezometers, as described in Dahm and Valett, 1996) of 1.9-cm diameter PVC were installed into the middle of the stream thalweg; these wells were perforated from 10 to 50 cm below the streambed. At least three mini-piezometers were randomly placed within each plot (see Figure 1; $n = 37$ for FC, $n = 13$ for FPD). All wells were installed in 2004 and not measured or sampled until the spring of 2005 to allow wells to settle.

Water sampling and analyses

Samples of streamwater for the $\delta^{18}\text{O}$ and $\delta^2\text{H}$ analyses were collected by grab-sampling while groundwater samples were collected with a peristaltic pump from wells and mini-piezometers. Groundwater and streamwater samples were taken from 4 of the 15 plots in June and September, 2005. Three of the four plots were located on FC and included the furthest upstream, a mid-stream plot and the furthest downstream plot. The fourth plot selected for water sampling was the furthest downstream plot on FPD. Plots were selected to both understand watershed effects (furthest downstream plots) in both FC and FPD and changes along the stream course in FC. Precipitation

was collected using six rainfall collectors placed throughout the watershed; while snowwater was collected from 10 snowpack cores sampled along two transects spanning a 100-m gradient of elevation. Precipitation samples were collected for six events from June through September, 2005 and snowpack samples were collected on a single occasion in April 2005. All samples were collected into acid-washed high-density polyethylene (HDPE) 125-ml bottles, then kept chilled on ice until filtration within 24-h of collection (0.7- μ glass fibre filters within a polypropylene inline filter holder attached to a peristaltic pump using sterilized tygon tubing). The samples were then kept in a dark refrigerator ($\sim 2^\circ\text{C}$) until sent to the USDA Forestry Sciences Lab in Moscow, Idaho for isotope analysis using Finnigan Delta + Continuous Flow isotope ratio mass spectrometer (IRMS) (Thermo Scientific, Waltham, MA).

Chloride analyses of monthly samples collected from all wells, piezometers and stream grab-stations, were conducted following vacuum filtration with an acid-washed polysulfone filter apparatus (0.7- μ glass fibre filter paper), on a Dionex DX 500TM ion chromatograph analyzer (Dionex, Sunnyvale, CA) at the USDA Forest Service Forestry Sciences Lab in Grand Rapids, MN. Monthly *in situ* measurements of temperature and conductivity were collected for all wells and stream sampling locations using a YSI 556TM multi-parameter meter (YSI Inc., Yellow Springs, OH). Outliers that were more than two times the standard error were omitted before analyses. Precipitation values for conductivity and chloride concentration were assumed to equal those measured at the nearest National Atmospheric Deposition Program site in Chassel, MI approximately 30-km northwest of the study site (<http://nadp.sws.uiuc.edu/sites/>).

Groundwater discharge

Stream gauging method. As a measure of groundwater contribution along each stream reach, the difference in discharge using stream gauging between an upstream and downstream pair of sample points was determined. The difference between upstream and downstream pairs was fitted for a regression (for FPD, $r^2 = 0.93$, $p < 0.05$; for FC, $r^2 = 0.96$, $p < 0.05$) which was used to predict groundwater discharge from the previously described daily discharge estimates. Given local topography constraints, we assume gaining reaches with no losses of streamwater.

Partitioning water sources

Stable isotopes. Isotope data were analysed using a mixing model procedure to partition water sources. This approach assumes no between-population correlation between isotopic signatures. An analysis of model error was performed with the Environmental Protection Agency, Western Ecology Division's free software program IsoError[®] (Phillips and Gregg, 2001, 2003). The simplified model equation used to partition different sources of each chemical and physical parameter, is

expressed as:

$$C_{\text{mixture}} = XC_{\text{sourceA}} + (1 - X)C_{\text{sourceB}}$$

where C_{mixture} is the mean of the water mixture of the two end members, and C_{sourceA} is the mean for one source end member, while C_{sourceB} is the mean for the other end member. Solving for X provides the percent contribution by each end member in the mixture.

To estimate the source contributions of water within the groundwater aquifer mixture (floodplain and terrace wells), the end members were pooled into late-winter snowpack isotopic values and into summer rain precipitation concentration values. For estimating the source contributions within the hyporheic zone (streambed mini-piezometers and bankfull wells), streamwater and groundwater concentration were considered the two end members. The two concentration end members used to partition streamwater were groundwater from streamside wells (mean of bankfull and flood prone locations) and surface water from the upstream Trout Lake and Florence Pond.

Conservative tracers. Because chloride is considered a highly soluble and a somewhat conservative natural tracer (Stream Solute Workshop, 1990; Hart *et al.*, 1999; Hinkle *et al.*, 2001; Thomas *et al.*, 2003), we used chloride concentration to partition hyporheic water sources of groundwater and streamwater. We also used measurements of conductivity and temperature as semi-conservative tracers for water source partitioning (Robson and Neal, 1990; Kleissen *et al.*, 1990; Cey *et al.*, 1998; Battin *et al.*, 2003; Wenninger *et al.*, 2004; Monteith *et al.*, 2006) within the hyporheic zone and streamwater (groundwater vs upstream reservoirs) estimates, respectively.

Statistical analyses

ANOVA was used to test for differences between well positions and sampling locations, while linear and non-linear regressions were performed for isotope and discharge comparisons. The univariate models, descriptive statistics, and Least Significant Differences (LSD) comparisons were performed using SPSS[®] statistical software, while all regressions were performed using SIGMAPLOT[®] software. All differences are considered significant at the $p \leq 0.05$ level, unless otherwise stated.

RESULTS AND DISCUSSION

Groundwater discharge

The stream gauging approach yielded estimates of groundwater discharge for the study period that ranged from 10.8 to 14.4 $\text{m}^3 \text{h}^{-1}$ for FC and 1.8 to 144 $\text{m}^3 \text{h}^{-1}$ for FPD. Groundwater discharge for each reach decreased from onset of snowmelt through the summer growing season, but increased with autumnal precipitation inputs, likely as a result of increased inputs and reduced evapotranspiration demands (Figure 3). The




Figure 3. Groundwater discharge of Florence Pond Drain (FPD) and Fisher Creek (FC), along with the Huron Mountain Reserve's rain precipitation inputs for 2005. Notice the difference in scale (log) for discharge between reaches.

greater range/relative flashiness of FPD, compared with FC, is likely because of a more constrained floodplain along the study reach with a shorter residence time of the relatively steep upslope recharge.

Partitioning water sources

Precipitation inputs. The distinct isotopic signatures of rain and snow precipitation permitted us to quantify the relative contributions of sources of water to groundwater and streamwater. By regressing $\delta^2\text{H}$ (‰) against values of $\delta^{18}\text{O}$ (‰) for precipitation, we developed a local meteoric water line (LMWL) where $\delta^2\text{H} = 7.8 \delta^{18}\text{O} + 14.1$ ($R^2 = 0.99$) (Figure 4). The LMWL very closely approximates the recognized global meteoric water line for precipitation with the formula $\delta^2\text{H} = 8.0 \delta^{18}\text{O} + 10$ (Craig, 1961). Upon superimposing our values for groundwater, streamwater, and upstream reservoirs (lake-waters), a local evaporation line (LEL) with the equation $\delta^2\text{H} = 5.0 \delta^{18}\text{O} - 26.5$ ($R^2 = 0.83$; $p < 0.0001$) is formed. The divergence of the LEL from the LMWL distinguishes the evaporative enrichment that occurs for the site, while the LEL intersection with the LMWL closely approximates the mean annual precipitation signature (Gibson *et al.*, 2005; Reddy *et al.*, 2006), which was approximately -14.5‰ for the FC watershed during 2005. Walker and Krabbenhoft (1998) found a 2-year volume weighted average of approximately -11‰ for a Northern Wisconsin site. Because the FC watershed is closer in proximity to Lake Superior it experiences




Figure 4. Local meteoric water line (LMWL; solid line) of $\delta^2\text{H}$ (‰) and $\delta^{18}\text{O}$ (‰) for all precipitation (snow and rain) collected within the Fisher Creek Watershed, 2005; along with the regressed local evaporation line (LEL; dashed line) for all ground- and surface-waters collected.

more lake-effect snow precipitation; therefore, the relative depletion in signature is expected. The evaporative trend is very similar to Reddy *et al.*'s (2006) LEL for a north-central Minnesota watershed, where most groundwater closely resembled the LMWL. FC's and FPD's groundwater and streamwater do not show much divergence from the LMWL; therefore, evaporative enrichment is relatively low within the stream. Conversely, the values for the upstream lakes (Trout Lake and Florence Pond) indicate strong enrichment, where ^2H preferentially evaporates over ^{18}O during the summer and/or rain dilution continues to enrich the chemical signature of the lakes.

Groundwater sources. The signature of meteoric water has been shown to follow a sine function relationship with the seasons (Dewalle *et al.*, 1997; Gibson *et al.*, 2005; Reddy *et al.*, 2006), but for shorter time steps the LMWL provides an indication of whether ground, stream, or lake-waters are derived from rain or snow. For 2005, FC streamwater was positioned between the range of values found for snow and rain. Isotopic analyses revealed that streamwater closely resembled groundwater (Figure 5), and that both stream and groundwaters were derived primarily from snowmelt, with minor contributions from rain or upstream lakes. Because streamwater during base flow was on average 80% groundwater, and groundwater was 85% snowmelt, we conclude that streamwater during baseflow was up to 70% of snowmelt origin.

The distinctness of our results is sensitive to any isotopic enrichment resulting from the loss of lighter water during sublimation of snow or evaporation of meltwater and/or intercepted rainwater. Several laboratory and *in situ* studies estimated the $\delta^{18}\text{O}$ enrichment of snow to snowmelt water to range from +1.4‰ to +5.6‰ (Herrmann *et al.*, 1981; Cooper *et al.*, 1993; Mast *et al.*, 1995; Suzuki, 1995; Taylor *et al.*, 2002). Mast *et al.* (1995) found an enrichment of +1.4‰ for a Colorado site while Taylor *et al.* (2002) found an enrichment of +4.5‰ for




Figure 5. $\delta^{18}\text{O}$ (‰) for select waters of the Fisher Creek Watershed, 2005. The snowmelt signature was elevated +4‰ from the snowpack signature to account for enrichment processes (estimated from Talyor *et al.*, 2002) used in mixing model analyses. Error bars are present, but hidden under most symbols.

another Colorado site and an enrichment of >5‰ for a Vermont site. To address enrichment in the mixing model, we conservatively assumed that at our site, snowmelt was enriched to a maximum of +4‰ relative to snow. In addition, soil evaporative enrichment can be large in arid climates (Gat, 1998), but an enrichment of +1.5‰ is considered a limit in humid climates (Gat, 1998). Considering this potential enrichment along with other selective enrichment factors (e.g. selective runoff processes and canopy interception), a maximum rain enrichment of 1.5‰ was evaluated within the mixing model.

Based on our analyses of the possible range of enrichment, the 95% confidence limit for the percent of groundwater that was snow derived for May ranged from 73% to 100% (Table I). For August the range was slightly lower at 69–97% snow-derived groundwater, which may be an indication of slight dilution with rain or evaporative enrichment that occurred since May. Given all the potential evaporative enrichment effects, our analysis conservatively suggests that ~85% of groundwater is snowmelt derived. Given evidence of low enrichment from stream values, we believe that this is a low estimate for snowmelt's contribution to groundwater within the FC Watershed. This conservatively high estimate of contribution underscores the importance of snowmelt to this riparian system.

Approximately 50% of annual runoff in the region occurs during and immediately after snowmelt (Stottlemyer and Toczydowski, 1996, 1999, 2006), with snowfall equalling approximately 50% of annual precipitation inputs (Eichenlaub 1970; Stottlemyer and Toczydowski, 2006). Because snowmelt occurs before the growing season commences, during which evapotranspiration losses are minimal, it is likely that a greater percentage of

Table I. Mixing model results for determining proportionality of source waters to groundwater for Fisher Creek, 2005 including standard errors (S.E.) and confidence limits (C.L.).

Groundwater Sample date and $\delta^{18}\text{O}$ corrections ^a	Snowmelt ($N = 8$; March)					Rain ($N = 23$; June–September)			
	$\delta^{18}\text{O}$ (‰)	$\delta^{18}\text{O}$ (‰)	% of groundwater	S.E.	95% C.L.	$\delta^{18}\text{O}$ (‰)	% of groundwater	S.E.	95% C.L.
May ($N = 9$)	-14.3	-19.1				-6.2			
Snow +3‰	-14.3	-16.1	82	4.3	73–91	-6.2	18	4.3	9–27
Snow +4‰; rain +1.5‰	-14.3	-15.1	92	4.3	83–100	-4.7	8	4.3	0–17
August ($N = 5$)	-13.9	-19.1				-6.2			
Snow +3‰	-13.9	-16.1	78	3.9	69–86	-6.2	22	3.9	14–31
Snow +4‰; rain +1.5‰	-13.9	-15.1	88	3.9	80–97	-4.7	12	3.9	3–20

^a Interpolated enrichment corrections based on data from Taylor *et al.*, 2002 & Gat, 1998.

Table II. ANOVA results with LSD for temperature, conductivity, and chloride of waters from the FC watershed by position in riparian landscape, June–October 2005 including standard errors (S.E.) ($n = 7$ samples).

Riparian position	Temperature			Conductivity			Chloride		
	°C	S.E.	LSD	mS cm ⁻¹	S.E.	LSD	ppm	S.E.	LSD
Streamwater	11.5	0.17	A	0.086	0.009	B	0.68	0.01	A
Streambed	10.4	0.12	B	0.149	0.003	A	0.66	0.01	A
Bankfull	9.84	0.07	C	0.148	0.003	A	0.63	0.01	B
Floodprone	9.76	0.07	C	0.141	0.003	A	0.62	0.01	B
Floodplain	8.67	0.10	D	0.085	0.006	B	0.64	0.01	B
Terrace	8.82	0.30	D	0.033	0.002	C	0.53	0.03	C

snowmelt water will reach the riparian groundwater system than growing season precipitation inputs. This may explain why the groundwater and streamwater signatures varied very little between May and August (Figure 5). The uneven distribution of snowmelt to rain precipitation during a moderately droughty 2005 (Figure 2), led to a snow-dominated riparian groundwater reservoir.

Streamwater sources. An isotope mixing model was used to separate groundwater and upstream lake water sources that contributed to streamwater in the FC and FPD reaches of study. Groundwater in May contributed ~75% of the streamwater, while groundwater in August contributed ~90% to streamwater. These results indicate that upstream lakes and ponds in May resemble snowmelt more than in August, and that groundwater is more influential in August because of lower surface water elevations of upstream lakes. Therefore, our hypothesis of an increasing percentage of groundwater derived streamwater during the recession limb of the hydrograph is supported.

Other results from this study support this isotope-based conclusion. Specific conductivity of waters varied by riparian position (Table II), with streambed, bankfull and floodprone positions exhibiting the highest conductivities, while more distant groundwater positions (floodplain + terrace wells) had the lowest conductivities. These results indicate that precipitation with low conductivity (<0.02 mS cm⁻¹) gains ions as it percolates through soils along a course to the stream. McGlynn *et al.* (1999) found similar increases in solute concentrations along

riparian flow paths. Based on mixing model results for conductivity, the June contribution to streamwater by upstream lakes (precipitation values as a proxy) was estimated at 40%, while in August it decreased to 23%. If one separates the analysis by reach, upstream sources contribute less to FPD streamwater than the FC reach, similar to earlier indications derived from isotope data. In addition, both reaches show increases in groundwater contribution from June to August, which corroborate earlier isotope data. The conductivity model results in a greater (2×) contribution from upstream lakes than the isotope model results (25% in May/June to 10% in August); however, the difference between months is similar; with a 12–15% decrease in upstream lake/upstream precipitation inputs to streamwater from June to August. The conductivity and isotope data both suggest that streamwater is dominated by groundwater inputs and upstream inputs to streamwater are reduced in August when surface water elevations in the upstream lakes have dropped. When the late autumnal period is analysed for conductivity, the contribution from upstream lakes rises to nearly 53% in October, which is a 13% increase from June. This pattern in upstream contribution correlates well with the observed hydrograph (Figure 3), where periods of elevated recharge to upstream lakes, such as early season snowmelt and autumn rain, result in reduced contribution of groundwater to streamwater.

Hyporheic zone. Isotope, chloride and water temperature mixing models were performed to describe the extent of the hyporheic zone and the relative contributions of

groundwater and streamwater to within hyporheic waters (Hinkle *et al.*, 2001; Battin *et al.*, 2003). Based on temperature and chloride concentration data, the stream contributed approximately 25% of the water found within waters <50 cm from the streambed (within bankfull and streambed wells) in early June, while the late summer values for these wells were indistinguishable from groundwater. With the onset of autumn rains, the stream contributed nearly 36% to bankfull and streambed wells, as determined from temperature measurements in October. Oxygen isotope signatures resulted in a 38% streamwater contribution to bankfull and streambed wells in late May, and 0% contribution in August because of identical groundwater and streamwater end members. Although there is a difference between the results for determining the percent contributions to the waters within bankfull and streambed wells, all results indicate that the hyporheic boundary was detectable within our sampling design, was consistently close to the streambed's margin (<50 cm), and contracted seasonally in unison with the discharge hydrograph, as hypothesized. For these sandy bottom stream reaches, there was no hyporheic mixing beyond 10-cm during baseflow conditions for the moderately droughty summer.

CONCLUSIONS

The groundwater dynamics during a period of moderate drought provided a snapshot of changing climatic effects on the riparian hydrology within an 'old-growth' Lake Superior watershed. Snowfall is increasing throughout the region, while total precipitation appears to be unchanged changing the annual distribution of precipitation, a predicted outcome from global climate change. Stottlemeyer and Toczydlowski (1996, 1999, 2006) estimated that regional snowpacks contribute ~50% to annual runoff or streamflow. Our estimations indicate that baseflow conditions of an old-growth riparian area are not evenly distributed. The isotope-derived mixing model estimated that snowmelt accounted for approximately 80% of the inputs to the riparian groundwater during 2005 baseflow conditions with the remaining 20% of inputs likely from rain during the previous non-growing season (spring and fall, 2004). Models also allowed for the observation that groundwater's relative contribution to streamflow increases along the recession limb of the hydrograph, as upstream reservoirs decrease in relative influence downstream. Future increases in snowfall, without an increase in rain, may further lead to water fluxes from snowmelt that increasingly dominate summer baseflow and could potentially increase spring flooding. However, because of overall declines in precipitation, streamflow patterns will likely change towards lower overall flow following the influence of the snowmelt period. Changes in flow sources will likely lead to changes in-stream chemistry which could alter the productivity of in-stream ecosystems.

We also observed that in our riparian areas, the hyporheic zone was very small in extent (<50 cm from

the streambed surface) and shrank during the growing season, as streamwater contribution and hyporheic zone expansion and contraction also followed the hydrograph. As the hyporheic zone boundary was indistinguishable during the drought conditions of summer baseflow, an argument could be made that the hyporheic zone processes are minimal and/or contracted during this period, while higher flow conditions expand the boundary and zone of hyporheic mixing, potentially leading to a greater impact on ecosystem processing of nutrients.

Our isotope-based analyses allowed us to track the flux and fate of water of a remote northern old-growth watershed, and highlighted the dominant role that snowmelt plays in the functioning of old-growth riparian ecosystems of the Great Lakes. Future work assessing climate change influences of water sources to streams in the region will be able to build on this research.

ACKNOWLEDGEMENTS

Funding for this research was provided by the Riparian and Aquatic Ecosystems work unit and the Belowground Processes work unit of the Northern Research Station of the USDA Forest Service. We also thank the Huron Mountain Wildlife Foundation for critical logistical and financial support for field sampling. We also would like to thank Joe Ridgeway and numerous field assistants who helped make the project successful.

REFERENCES

- Atekwana EA, Richardson DS. 2004. Geochemical and isotopic evidence of a groundwater source in the Corral Canyon meadow complex, central Nevada, USA. *Hydrological Processes* **18**: 2801–2815.
- Baker MA, Valett HM, Dahm CN. 2000. Organic carbon supply and metabolism in a shallow groundwater ecosystem. *Ecology* **81**: 3133–3148.
- Battin TJ, Kaplan LA, Newbold JD, Hendricks SP. 2003. A mixing model analysis of stream solute dynamics and the contribution of a hyporheic zone to ecosystem function. *Freshwater Biology* **48**: 995–1014.
- Bragg DC, Kershner JL. 1999. Coarse woody debris in riparian zones. *Journal of Forestry* **97**: 30–35.
- Burnett AW, Kirby ME, Mullins HT, Patterson WP. 2003. Increasing Great Lake-Effect snowfall during the twentieth century: a regional response to global warming? *Journal of Climate* **16**: 3535–3542.
- Burns DA, McDonnell JJ, Hooper RP, Peters NE, Freer JE, Kendall C, Beven K. 2001. Quantifying contributions to storm runoff through end-member mixing analysis and hydrologic measurements at the Panola Mountain Research Watershed (Georgia, USA). *Hydrological Processes* **15**: 1903–1924.
- Buttle JM. 1998. Fundamentals of small catchment hydrology. In *Isotope Tracers in Catchment Hydrology*, Kendall C, McDonnell JJ (eds). Elsevier: Amsterdam; 1–49.
- Buttle JM, Hazlett PW, Murray CD, Creed IF, Jeffries DS, Semkin R. 2001. Prediction of groundwater characteristics in forested and harvested basins during spring snowmelt using a topographic index. *Hydrological Processes* **15**: 3389–3407.
- Cey EE, Rudolph DL, Parkin GW, Aravena R. 1998. Quantifying groundwater discharge to a small perennial stream in southern Ontario, Canada. *Journal of Hydrology* **210**: 21–37.
- Christopherson N, Neal C, Hooper RP, Vogt RD, Andersen S. 1990. Modelling streamwater chemistry as a mixture of soilwater end-members—a step towards second-generation acidification models. *Journal of Hydrology* **116**: 307–320.
- Cirino CP, McDonnell JJ. 1997. Linking the hydrologic and biogeochemical controls of nitrogen transport in near-stream zones of temperate-forested catchments: a review. *Journal of Hydrology* **199**: 88–120.

- Cooper LW, Solis C, Kane DL, Hinzman LD. 1993. Application of oxygen-18 tracer techniques to arctic hydrological processes. *Arctic and Alpine Research* **25**: 247–255.
- Craig H. 1961. Isotopic variations in meteoric waters. *Science* **133**: 1702–1703.
- Dahm CN, Valett HM. 1996. Hyporheic zones. In *Methods in Stream Ecology*, Hauer FR, Lamberti GA (eds). Academic Press: San Diego; 107–122.
- Davis M, Douglas C, Calcote R, Cole KL, Winkler MG, Flakne R. 2000. Holocene climate in the western Great Lakes National Parks and lakeshores: implications for future climate change. *Conservation Biology* **14**: 968–983.
- Denton SR, Barnes BV. 1988. An ecological climatic classification of Michigan: a quantitative approach. *Forest Science* **34**: 119–138.
- Dewalle DR, Edwards PJ, Swistock BR, Aravena R, Drimmie RJ. 1997. Seasonal isotope hydrology of three Appalachian forest catchments. *Hydrological Processes* **11**: 1895–1997.
- Dorr JA, Eschman DF. 2001. *Geology of Michigan*. The University of Michigan Press: Ann Arbor.
- Eichenlaub LV. 1970. Lake effect snowfall to the lee of the Great Lakes: its role in Michigan. *Bulletin American Meteorological Society* **51**: 403–412.
- Flaspohler DJ, Meine C. 2006. Planning for wildness: Aldo Leopold's report on Huron Mountain Club. *Journal of Forestry* **104**: 32–42.
- Frelich LE. 1995. Old forests in the Lake States today and before European settlement. *Natural Areas Journal* **15**: 157–167.
- Gat JR. 1998. Stable isotopes, the hydrological cycle and the terrestrial biosphere. In *Stable Isotopes: Integration of Biological, Ecological, and Geochemical Processes*, Griffiths H (ed). BIOS Scientific Publishers Ltd: Oxford; 397–407.
- Gibson JJ, Edwards TWD, Birks SJ, St Amour NA, Buhay WM, McEachern P, Wolfe BB, Peters DL. 2005. Progress in isotope tracer hydrology in Canada. *Hydrological Processes* **19**: 303–327.
- Goebel PC. 2001. *Hydrogeomorphic controls on riparian areas of the Northern Lake States*. PhD Dissertation, Michigan Technological University, Houghton, MI.
- Hart DR, Mulholland PJ, Marzolf ER, DeAngelis DL, Hendricks SP. 1999. Relationships between hydraulic parameters in a small stream under varying flow and seasonal conditions. *Hydrological Processes* **13**: 1497–1510.
- Harvey JW, Wagner BJ. 2000. Quantifying hydrologic interactions between streams and their subsurface hyporheic zones. In *Streams and Ground Waters*. Jones JB, Mulholland PJ (eds). Academic Press: San Diego; 3–44.
- Hedman CW, Van Lear DH, Swank WT. 1996. In-stream large woody debris loading and riparian forest seral stage associations in the southern Appalachian Mountains. *Canadian Journal of Forest Resources* **26**: 1218–1227.
- Herrmann A, Lehrer M, Stichler W. 1981. Isotope input into runoff systems from melting snow covers. *Nordic Hydrology* **12**: 309–318.
- Hill AR. 2000. Stream chemistry and riparian zones. In *Streams and Ground Waters*, Jones JB, Mulholland PJ (eds). Academic Press: San Diego; 83–110.
- Hinkle SR, Duff JH, Triska FJ, Laenen A, Gates EB, Bencala KE, Wentz DA, Silva SR. 2001. Linking hyporheic flow and nitrogen cycling near the Willamette River—a large river in Oregon, USA. *Journal of Hydrology* **244**: 157–180.
- Hooper RP, Christopherson N, Peters NE. 1990. Modelling streamwater chemistry as a mixture of soilwater end-members—an application to the Panola Mountain Catchment, Georgia, U.S.A. *Journal of Hydrology* **116**: 321–343.
- Johannessen M, Henriksen A. 1978. Chemistry of snow meltwater: changes in concentration during melting. *Water Resources Research* **14**: 615–619.
- Katsuyama M, Ohte N, Kobashi S. 2001. A three-component end-member analysis of streamwater hydrochemistry in a small Japanese forested headwater catchment. *Hydrological Processes* **15**: 249–260.
- Kattenberg AF, Giorgi H, Grassi GH, Meehl JFB, Mitchell RJ, Stouffer T, Tokioka AJ, Weaver J, Wigley TML. 1996. Climate models: projections of future climate. In *Climate Change 1995: The Science of Climate Change*, Houghton JT, Fihlo LGM, Callander BA, Harris N, Kattenberg A, Maskell K (eds). Cambridge University Press: Cambridge, UK.
- Kleissen FM, Wheeler HS, Beck MB, Harriman R. 1990. Conservative mixing of water sources: analysis of the behaviour of the Allt A' Mharchaidh Catchment. *Journal of Hydrology* **116**: 365–374.
- Kunkel KE, Wescott NE, Kristovich DAR. 2000. Climate change and lake effect snow. In *Preparing for a Changing Climate: The Potential Consequences of Climate Variability and Change*, Sousounis PJ, Bisanz JM (eds). U.S. EPA, Office of Research and Development Global Change Research Program: Washington D.C.; 25–28.
- Ladouche B, Probst A, Viville D, Idir S, Baque D, Loubet M, Probst J-L, Bariac T. 2001. Hydrograph separation using isotopic, chemical, and hydrological approaches (Strengbach catchment, France). *Journal of Hydrology* **242**: 255–274.
- Leathers DJ, Ellis AW. 1996. Synoptic mechanisms associated with snowfall increases to the lee of Lakes Erie and Ontario. *International Journal of Climatology* **16**: 1117–1135.
- Mast AM, Kendall C, Campbell DH, Clow DW, Black J. 1995. Determination of hydrologic pathways in an alpine-subalpine basin using isotopic and chemical tracers, Loch Vale Watershed, Colorado, USA. In *Biogeochemistry of Seasonally Snow-Covered Catchments*, IAHS Publication 228, Tennesen KA, Williams MW, Tranter M (eds). IAHS Press: Wallingford, UK; 263–270.
- McClain ME, Boyer EW, Dent CL, Gergel SE, Grimm NB, Groffman PM, Hart SC, Harvey JW, Johnston CA, Mayorga E, McDowell WH, Pinay G. 2003. Biogeochemical hot spots and hot moments at the interface of terrestrial and aquatic ecosystems. *Ecosystems* **6**: 301–312.
- McClure JD, Kolka RK, White A. 2004. Effect of forest harvesting Best Management Practices (BMPs) on coarse woody debris in stream and riparian zones in three Appalachian watersheds. *Water, Air, Soil Pollution: Focus* **4**: 245–261.
- McGlynn BL, McDonnell JJ, Shanley JB, Kendall C. 1999. Riparian zone flowpath dynamics during snowmelt in a small headwater catchment. *Journal of Hydrology* **222**: 75–92.
- Monteith SS, Buttle JM, Hazlett PW, Beall FD, Semkin RG, Jeffries DS. 2006. Paired-basin comparison of hydrologic response in harvested and undisturbed hardwood forests during snowmelt in central Ontario: II. Streamflow sources and groundwater residence times. *Hydrological Processes* **20**: 1117–1136.
- Montgomery DR. 1999. Process domains and the river continuum. *Journal of the American Water Resources Association* **35**: 397–410.
- Morrice JA, Valett HM, Dahm CN, Campana ME. 1997. Alluvial characteristics, groundwater-surface water exchange and hydrological retention in headwater streams. *Hydrological Processes* **11**: 253–267.
- Mulholland PJ. 1992. Regulation of nutrient concentrations in a temperate forest stream: Roles of upland, riparian, and instream processes. *Limnology and Oceanography* **37**: 1512–1526.
- Naiman RJ, Decamps H. 1997. The ecology of interfaces: riparian zones. *Annual Review of Ecology, Evolution and Systematics* **28**: 621–658.
- Naiman RJ, Balian EV, Bartz KK, Bilby RE, Latterell JJ. 2002. Dead Wood Dynamics in Stream Ecosystems. USDA Forest Service General Technical Report, PSW-GTR-181. USDA Forest Service, Washington D.C.
- Naiman RJ, Decamps H, McClain ME. 2005. *Riparia: Ecology, Conservation, and Management of Streamside Communities*. Elsevier: London.
- National Research Council. 2002. *Riparian Areas: Functions and Strategies for Management*. National Academy Press: Washington DC.
- Nijssen B, O'Donnell GM, Hamlet AF, Lettenmaier DP. 2001. Hydrologic sensitivity of global rivers to climate change. *Climatic Change* **50**: 143–175.
- Norton DC, Bolsenga SJ. 1993. Spatiotemporal trends in lake effect and continental snowfall in the Laurentian Great Lakes, 1951–1980. *Journal of Climate* **6**: 1943–1956.
- Pardo LH, Kendall C, Pett-Ridge J, Chang CCY. 2004. Evaluating the source of streamwater nitrate using $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ in nitrate in two watersheds in New Hampshire, USA. *Hydrological Processes* **18**: 2699–2712.
- Phillips DL, Gregg JW. 2001. Uncertainty in source partitioning using stable isotopes. *Oecologia* **127**: 171–179.
- Phillips DL, Gregg JW. 2003. Source partitioning using stable isotopes: coping with too many sources. *Oecologia* **136**: 261–269.
- Rascher CM, Driscoll CT, Peters NE. 1987. Concentration and flux of solutes from snow and forest floor during snowmelt in the West-Central Adirondack region of New York. *Biogeochemistry* **3**: 209–224.
- Reddy MM, Schuster P, Kendall C, Reddy MB. 2006. Characterization of surface and ground water $\delta^{18}\text{O}$ seasonal variation and its use for estimating groundwater residence times. *Hydrological Processes* **20**: 1753–1772.
- Robson A, Neal C. 1990. Hydrograph separation using chemical techniques: an application to catchments in Mid-Wales. *Journal of Hydrology* **116**: 345–363.
- Rosgen DL. 1994. A classification of natural rivers. *Catena* **22**: 169–199.

- Schindler JE, Krabbenhoft DP. 1998. The hyporheic zone as a source of dissolved organic carbon and carbon gases to a temperate forested stream. *Biogeochemistry* **43**: 157–174.
- Seibert J, Bishop K, Rodhe A, McDonnell JJ. 2003. Groundwater dynamics along a hillslope: a test of the steady state hypothesis. *Water Resources Research* **39**: 1–9.
- Simpson TB. 1990. *Landscape ecosystem and cover types of the reserve area and adjacent lands of the Huron Mountain Club, Marquette County, Michigan*. PhD Thesis, The University of Michigan, Ann Arbor, USA.
- Simpson TB, Stuart PE, Barnes BV. 1989. *Cover Types at the Huron Mountain Club Marquette County, Michigan (map)*. Huron Mountain Wildlife Foundation: Big Bay, MI.
- Simpson TB, Stuart PE, Barnes BV. 1990. Landscape ecosystems and cover types of the reserve area and adjacent lands of the Huron Mountain Club. *Occasional Papers of the Huron Mountain Wildlife Foundation* **4**: 128 pp.
- Stottlemeyer R, Toczydlowski D. 1991. Stream chemistry and hydrologic pathways during snowmelt in a small watershed adjacent Lake Superior. *Biogeochemistry* **13**: 177–197.
- Stottlemeyer R, Toczydlowski D. 1996. Precipitation, snowpack, stream-water ion chemistry, and flux in a northern Michigan watershed, 1982–1991. *Canadian Journal of Fisheries and Aquatic Sciences* **53**: 2659–2672.
- Stottlemeyer R, Toczydlowski D. 1999. Seasonal change in precipitation, snowpack, snowmelt, soil water, and streamwater chemistry, northern Michigan. *Hydrological Processes* **13**: 2215–2231.
- Stottlemeyer R, Toczydlowski D. 2006. Effect of reduced winter precipitation and increased temperature on watershed solute flux, 1988–2002, Northern Michigan. *Biogeochemistry* **77**: 409–440.
- Stream Solute Workshop. 1990. Concepts and methods for assessing solute dynamics in stream ecosystems. *Journal of the North American Benthological Society* **9**: 95–119.
- Suzuki K. 1995. Hydrochemical study of snow meltwater and snow cover. In *Biogeochemistry of Seasonally Snow-Covered Catchments*, IAHS Publication 228. IAHS Press: Wallingford, UK; 107–114.
- Taylor S, Feng X, Williams M, McNamara J. 2002. How isotopic fractionation of snowmelt affects hydrograph separation. *Hydrologic Processes* **16**: 3683–3690.
- Thomas SA, Valett HM, Mulholland PJ, Fellows CF, Webster JR, Dahm CN, Peterson CG. 2001. Nitrogen retention in headwater streams: the influence of groundwater-surface water exchange. *The Scientific World Journal* **1**: 623–631.
- Thomas SA, Valett HM, Webster JR, Mulholland PJ. 2003. A regression approach to estimating reactive solute uptake in advective and transient storage zones of stream ecosystems. *Advances in Water Resources* **26**: 965–976.
- Valett HM, Crenshaw CL, Wagner PF. 2002. Stream nutrient uptake, forest succession, and biogeochemical theory. *Ecology* **83**: 2888–2901.
- Vidon PGF, Hill AR. 2004. Landscape controls on the hydrology of stream riparian zones. *Journal of Hydrology* **292**: 210–228.
- Walker JF, Krabbenhoft, DF. 1998. Groundwater and surface-water interactions in riparian and lake-dominated systems. In *Isotope Tracers in Catchment Hydrology*, Kendall C, McDonnell JJ (eds). Elsevier: Amsterdam; 467–488.
- Wells JR, Thompson PW. 1976. Vegetation and flora of the Huron Mountains. *Occasional Papers of the Huron Mountain Wildlife Foundation* **3**: 59 pp.
- Wenninger J, Uhlenbrook S, Tilch N, Leibundgut C. 2004. Experimental evidence of fast groundwater responses in a hillslope/floodplain area in the Black Forest Mountains, Germany. *Hydrological Processes* **18**: 3305–3322.
- Winograd IJ, Riggs AC, Coplen TB. 1998. The relative contributions of summer and cool-season precipitation to groundwater recharge, Spring Mountains, Nevada, USA. *Hydrogeology Journal* **6**: 77–93.
- Wondzell SM. 2005. Effect of morphology and discharge on hyporheic exchange flows in two small streams in the Cascade Mountains of Oregon, USA. *Hydrologic Processes* **20**: 267–287.
- Woods KD. 2000. Dynamics in late-successional hemlock-hardwood forests over three decades. *Ecology* **81**: 110–126.