

# Verbenone-releasing flakes protect individual *Pinus contorta* trees from attack by *Dendroctonus ponderosae* and *Dendroctonus valens* (Coleoptera: Curculionidae, Scolytinae)

Nancy E. Gillette, John D. Stein\*, Donald R. Owen†, Jeffrey N. Webster‡, Gary O. Fiddler§, Sylvia R. Mori and David L. Wood¶

USDA Forest Service, PSW Research Station, Berkeley, CA 94701, U.S.A., \*USDA Forest Service, Forest Health Technology, Morgantown, WV 26505, U.S.A., †California Department of Forestry and Fire Protection, Redding, CA 96002, U.S.A., ‡Total Forestry, Anderson, CA 96007, U.S.A., §USDA Forest Service, Redding Silviculture Laboratory, Redding, CA 96002, U.S.A. and ¶Dept. Environmental Science, Policy, and Management, University of California, Berkeley, CA 94720, U.S.A.

- Abstract**
- 1 In a study site in interior northern California, twenty individual lodgepole pines *Pinus contorta* were sprayed with a suspension of DISRUPT Micro-Flake® Verbenone (4,6,6-trimethylbicyclo(3.1)hept-3-en-2-one) Bark Beetle Anti-Aggregant flakes (Hercon Environmental, Emigsville, Pennsylvania) in water, with sticker and thickener, from ground level to a height of 7 m. Twenty trees sprayed with just water, sticker and thickener served as controls. All trees were baited immediately after spraying with mountain pine beetle *Dendroctonus ponderosae* aggregation pheromone lures, and lures were refreshed after 4 weeks.
  - 2 Trees treated with verbenone had significantly lower attack density by *D. ponderosae* than controls at 2, 4, 6 and 8 weeks after application of flakes.
  - 3 None of the treated trees was attacked by red turpentine beetle *Dendroctonus valens*, whereas control trees averaged nearly two *D. valens* attacks per tree, 8 weeks after treatment.
  - 4 A dry frass index, used to predict ultimate tree mortality, was significantly higher in control trees than treated trees for all four sampling intervals. This index proved to be a significant predictor of ultimate tree mortality.
  - 5 Ten months after application, treated trees showed significantly lower mortality than control trees.

**Keywords** Individual tree protection, lodgepole pine, mountain pine beetle, pheromones, Pinaceae, red turpentine beetle, Scolytinae, Scolytidae, semiochemicals, 4,6,6-trimethylbicyclo(3.1)hept-3-en-2-one.

## Introduction

The mountain pine beetle *Dendroctonus ponderosa* Hopkins is the most important pest of lodgepole pine *Pinus contorta* Douglas ex Loudon throughout its range in North America (Furniss & Carolin, 1977; Wood *et al.*, 2003). *Dendroctonus ponderosae* typically erupts in large, episodic outbreaks. A current infestation in British Columbia, Canada has already reached over 11 million hectares in size and has breached the Continental Divide, spilling over into interior North America (Wilent, 2005). There is concern that *D. ponderosae* will

colonize millions of acres of jack pine *Pinus banksiana* Lamb. in the vast boreal forests of Canada and the north central U.S.A., and that global climate change will favour *D. ponderosae* range extensions into this habitat (Logan & Powell, 2001; Wilent, 2005). Indeed, jack pine and another boreal pine, *Pinus flexilis* James, have been shown to be adequate hosts for *D. ponderosae* (Cerezke, 1995). *Dendroctonus ponderosae* also attacks other economically important pines, such as ponderosa pine *Pinus ponderosa* Douglas ex P. & C. Lawson (Furniss & Carolin, 1977; Wood *et al.*, 2003) and ecologically important pines such as whitebark pine *Pinus albicaulis* Engelm. (Keane, 2001; Tomback *et al.*, 2001; Schwandt & Kegley, 2004). It has been suggested that factors such as elevated stand density, fire, drought and air pollution contribute to tree stress and thereby increase levels of bark

Correspondence: Nancy E. Gillette, USDA Forest Service, PSW Research Station, PO Box 245, Berkeley, CA 94701, U.S.A. Tel.: +1 510 559 6474; fax: +1 510 559 6499; e-mail: ngillette@fs.fed.us

beetle attack (Waters, 1985a; Amman & Logan, 1998; McHugh *et al.*, 2003; Elkin & Reid, 2004; Jones *et al.*, 2004; Safranyik *et al.*, 2004).

Models and stand hazard rating systems have been developed to predict *P. contorta* susceptibility to *D. ponderosae* (Amman & Logan, 1998; Shore *et al.*, 2000; Smith *et al.*, 2002; Perkins & Roberts, 2003; Negron & Popp, 2004; Wulder *et al.*, 2004), and many attempts have been made to develop silvicultural (Fiddler *et al.*, 1989; Johnstone, 2002) and chemical (Taylor *et al.*, 1997; Haverty *et al.*, 1998; Naumann & Rankin, 1999) means to manage *D. ponderosae* damage in pine stands. Although maintenance of stand health through silviculture is arguably the most durable and effective method of protecting trees from bark beetle outbreaks (Wood *et al.*, 1985; Whitehead & Russo, 2005; Zausen *et al.*, 2005), such treatments can be difficult to implement in remote, steep or environmentally sensitive pine stands. Silvicultural treatments, such as thinning to reduce basal area in heavily stocked stands, are labour-intensive and can present regulatory hurdles (Waters, 1985a; Wood *et al.*, 1985). Proximity of infested stands to human activities, such as parks, campgrounds and wildland–urban interfaces, can severely limit the use of pesticides. Because silvicultural and pesticide treatments can be used only under somewhat limited conditions, considerable effort has been made to develop semiochemical approaches for the control of bark beetles in pine stands.

S-(–)-verbenone, [(1S,5S)-4,6,6-trimethylbicyclo[3.1.1]hept-3-en-2-one], a beetle-produced anti-aggregation pheromone also found in pines (Kainulainen & Holopainen, 2002) and a wide variety of angiosperms (Molyneux *et al.*, 1980; Guillen & Cabo, 1996; Fournier *et al.*, 1997; Buttery *et al.*, 2000; Umamo *et al.*, 2000; Pintore *et al.*, 2002; Sefidkon *et al.*, 2002; Ghannadi & Zolfaghari, 2003; Robles *et al.*, 2003), can be effective in limiting damage to pines by *Dendroctonus* spp. bark beetles (Payne & Billings, 1989; Payne *et al.*, 1992; Salom *et al.*, 1995; Borden, 1997; Clarke *et al.*, 1999; Lindgren & Miller, 2002; Kegley *et al.*, 2003; Progar, 2003; Gibson & Kegley, 2004; Kegley & Gibson, 2004; Bentz *et al.*, 2005; Progar, 2005). Although verbenone has often been shown to be effective, the results obtained have not been entirely consistent. Progar (2005) reported that verbenone pouches initially reduced bark beetle-induced pine mortality in central Idaho, U.S.A., but that the effect did not last for the entire 5-year treatment period. However, in that study, the same plots were treated every year, and stand basal area was dramatically reduced by beetle activity in both the controls and surrounding untreated stands relative to treated plots, rendering the comparison problematic. Bentz *et al.* (2005) found that deployment of verbenone pouches in lodgepole and whitebark pine stands significantly reduced rate of attack by mountain pine beetle for up to three consecutive years, but they reported that some treated plots, particularly those with large emerging beetle populations, showed higher attack rates than controls. In the two latter cases, greater concentrations of verbenone may have been needed to overcome high concentrations of aggregation pheromones released from large beetle populations. In other earlier studies, failures may have resulted from photoisomerization of

verbenone to chrysanthenone (Kostyk *et al.*, 1993), but more recent formulations have stabilizers to prevent isomerization. Lister *et al.* (1990) found that verbenone bubblecaps were ineffective for single tree protection, but the bubblecap release devices used in their study may have release properties, such as rate and distribution of release, that are not optimal for protection from scolytid attack. For example, an earlier polyolefin bead formulation was shown to release too rapidly to provide protection throughout the beetle flight periods (Holsten *et al.*, 2000), probably explaining the inconsistent results seen with that formulation in field trials (Shea *et al.*, 1992). We speculated that many small, point-source, reservoir-type releasers with longer-lasting release periods, such as flakes, might provide better pheromone dispersal and could better simulate natural beetle release in a forest stand, giving better efficacy than larger releasers such as pouches and bubble-caps. In addition to potentially superior release characteristics, the flakes represent a less concentrated source of the active ingredient than do plastic pouch and bubble cap formulations, and may present less of a risk from tampering. The latex sticker used in the application provides excellent adhesion over several months (N. E. Gillette *et al.*, unpublished data), so there is extremely little risk of accidental removal by wildlife brushing up against the treated trees. In some cases, discrete, removable devices may have regulatory advantages over small, highly dispersed release devices such as flakes. However, most discrete, removable devices that have been developed for use in managing bark beetles require placement at a height of 5 m above ground, making removal (as well as deployment) an operational challenge. Although the flakes, unlike pouches and bubble caps, are not discrete removable devices, each flake contains only a small quantity of active ingredient and they are firmly glued to the bark, making it difficult to conceive that wildlife or park visitors would remove enough of them to present a hazard. If desired, sprays could even be applied above the reach of children. We used a hydroseeder for our applications to achieve a high level of consistency, which limited our study sites to roadside treatments. However, in an operational setting, other application methods, such as paint-balls and aerial sprays, would open up the possibility of this type of treatment for steep and remote areas.

Other chemicals, such as green leaf volatiles and nonhost angiosperm volatiles, may increase the efficacy of verbenone in protecting pines from attack by bark beetles (Wilson *et al.*, 1996; Huber & Borden, 2001; Borden *et al.*, 2003; Pureswaran & Borden, 2004; Fettig *et al.*, 2005). However, in these studies, at least eight additional components were required to achieve significant control of mountain pine beetle (Huber & Borden, 2001), and only a nine-component blend added to verbenone performed better than high-release verbenone alone for control of western pine beetle *Dendroctonus brevicomis* LeConte (Fettig *et al.*, 2005). Each additional component must undergo regulatory evaluation, including an array of fish, avian, invertebrate, plant and mammalian toxicity tests, for it to be registered for use in pest control. That process might be prohibitive for such complex blends.

We hypothesized that a single-component pheromone could be effective if deployed in a release system with

the proper release characteristics (i.e. sustained, high release lasting throughout the beetle flight period). Such a single-component release system has obvious regulatory and marketing advantages, which are often crucial for product development for forestry uses. We selected verbenone as the active ingredient because of its demonstrated efficacy as an anti-aggregation pheromone for a wide variety of species in the genus *Dendroctonus*, and in particular *D. ponderosae* (Borden, 1997). We chose a laminated flake release device with the expectation that such a formulation, in which the active ingredient is held largely within the release device, might have superior release characteristics that would provide sustained levels of verbenone in a stand throughout the beetle flight period, normally late May to August in interior northern California (Furniss & Carolin, 1977; Waters, 1985a). In the summer of 2004, we field-tested a laminated 'flake' formulation of verbenone (DISRUPT Micro-Flake® Verbenone Bark Beetle Anti-Aggregant, Hercon Environmental, Emigsville, Pennsylvania) similar to one used for decades in the Gypsy Moth *Lymantria dispar* Linnaeus 'Slow-the-Spread' program (Sharov *et al.*, 2002).

We report the results of tests of the verbenone flake formulation applied to individual pine trees for protection from attack by *D. ponderosae*. The red turpentine beetle *Dendroctonus valens* LeConte, a secondary pest that sometimes attacks trees weakened by previous *D. ponderosae* attacks (Wood *et al.*, 2003), was also present in our study site and we therefore exploited the opportunity to assess efficacy against this pest too. This is the first report of a laminated flake formulation for protection of forest trees from bark beetle attack.

## Materials and methods

### Study location

The study was installed on 12 August 2004 on industrial forestry lands in Section 31, Siskiyou County, California (122°7'24.88W, 41°31'42.24N, approximately 5.6 km north-east of the peak of Mt. Shasta), T43N, R3W, Section 36, at an elevation of 2240 m. The site is in the rain shadow of the southern Cascade Range on the edge of the Basin and Range Province, and is characterized by cool, dry summers and very cold winters. The forest consists of secondary growth natural stand composed of interior lodgepole pine with a small component of Shasta red fir *Abies magnifica* var. *shastensis* Lemm. The site was chosen because it has a history of *D. ponderosae* infestations and is characteristic of much of industrial forestry land in the western U.S.A. (i.e. basal area and competing vegetation levels are lower than in most publicly managed lands). The flight periods of both *D. ponderosae* and *D. valens* typically begin 2–3 months before applications were made (Wood *et al.*, 2003); thus, we baited trees with aggregation pheromones to ensure high levels of attack by *D. ponderosae*. Earlier tests (N. E. Gillette *et al.*, unpublished data) indicated that trees under attack by *D. ponderosae* are likely to be attacked by *D. valens* as well. Because there is no known aggregation pheromone for *D. valens*, we relied on existing host volatiles cues to attract *D. valens*.

### Pheromone formulation

(S)-(-)-verbenone (Bedoukian Research, Connecticut) was formulated as the active ingredient (AI) in a polyvinyl chloride resin layer sandwiched between two impermeable polymer layers (Hercon Environmental) at the rate of 15% wt/wt AI (i.e. 150 g AI/kg flake material). The sheets of laminate were cut into 3 × 3 mm squares, which were suspended in a solution of distilled water with 4.0% (v/v) Gelva sticker (Hercon Environmental) and 5.72 g/L guar gum (Hercon Environmental) as thickener. A solution of the formulation, identical except for the absence of flakes, was prepared for application to control trees.

### Application

Application was made on 12 August 2004 using a modified hydroseeder seed/mulch spraying apparatus (Hercon Environmental) at the rate of 11.4 L/tree carrying 100 g/tree of flakes and 15 g/tree of AI. Flake and control formulations were sprayed onto trees in a single strip, approximately 30 cm wide, on a single side of each tree from ground level to a height of 7 m. Laboratory pheromone release data provided by the manufacturer showed that the flakes release verbenone at the rate of approximately 1 mg/day per gram of product, and we estimate that each tree therefore released approximately 100 mg verbenone/day during beetle flight. Immediately after treatment, each tree was baited at breast height with a mountain pine beetle lure (Phero Tech, Inc., Canada) containing *trans*-verbenol and *cis*-verbenol, in a ratio of 87 : 13, plus *exo*-brevicomin. Each lure was taped over half of its surface with duct tape to reduce the release rate of lure components because previous experience indicated that full strength aggregation pheromone lures were probably too strong a cue. Lures were refreshed 4 weeks after treatment, and were left in place over the winter because early snowstorms blocked access to the site after the 8-week sampling interval. The site remained inaccessible for 9 months after the last sampling interval.

### Experimental design, sampling and statistical analysis

Forty trees of similar size (as measured by diameter at breast height; d.b.h.) and live crown ratio were selected from among all accessible (i.e. roadside) trees in the stand. Hydroseeders must be towed by an all-terrain vehicle, and therefore not all trees within the stand were accessible for treatment. Trees selected for the study were a minimum of 50 m apart. Twenty trees were selected at random for treatment from among the 40 trees, and the remainder served as controls. Both d.b.h. and live crown ratio are thought to be correlated with tree susceptibility to *D. ponderosae* attack, with larger trees and trees with lower ratios of live crown to total tree height being considered more susceptible to attack (Waters, 1985b). Competition among trees, as measured by basal area of tree surface per hectare, is also thought to contribute to susceptibility to bark beetle attack (Wood *et al.*, 1985). Trees were therefore measured for d.b.h., live crown ratio and basal area, and these traits were subjected to a chi-square test to ensure

that there was no inadvertent bias in the random selection of trees for treatment vs. control categories. There were no significant differences in d.b.h., basal area, or live crown ratio between treated and control trees (Table 1).

A 30-cm band was demarcated on the bole of each tree, with the base of the band centred at breast height. At 2, 4, 6 and 8 weeks after treatment, counts were made of the number of resinous pitch tubes (*D. ponderosae* attacks) within each of these 30-cm bands. At the same time intervals, counts were also made below the bottom of the 30-cm base of the number of *D. valens* attacks per tree and the number *D. ponderosae* attacks producing dry frass. The latter number was coded as a dry frass index (0 = no dry frass; 1 = one to ten attacks with dry frass; 2 = more than ten attacks with dry frass). These categories were intended to provide a prediction of ultimate tree mortality because trees that have many attacks with dry frass (as opposed to highly resinous pitch tubes) are considered likely to have depleted their defensive resources and are more likely to succumb to bark beetle attack. We assumed that no trees in category 0 would ultimately die, some in category 1 would die, and all in category 2 would die. At 11 months after treatment, trees were inspected and ranked into one of three categories: (1) obviously dead, (2) obviously alive, or (3) uncertain status. The obviously living trees had deep green foliage overall, whereas the obviously dead trees had lime-green ('fading') to yellow, orange or brown foliage. Trees placed in the 'uncertain' category still had deep green foliage but had evidence of reduced resin flow (many dry frass tubes), or individual branches with brown foliage. There can be some spatial overlap on tree boles between attacks by *D. ponderosae* and *D. valens*, but *D. valens* normally produces much more resinous pitch tubes that are located lower on the bole than *D. ponderosae* pitch tubes. However, when the species of attacking beetle was uncertain, we later removed bark in the area of the pitch tubes to examine beetle galleries, which are clearly diagnostic for the two species (Furniss & Carolin, 1977).

The numbers of *D. ponderosae* and *D. valens* attacks per unit area of bark (attack density) were analysed with generalized linear models for over-dispersed Poisson distributed responses for counts with repeated measures (McCulloch & Searle, 2001) offset by the log of the area of tree trunk measured for rate of attack. The dry frass index was analysed with the Pearson's chi-squared test for independence for a three-way contingency table (Agresti, 1996) and graphically displayed using Mosaic plots (Friendly, 1994). The final dry frass indices were fitted with the proportional odds model (Collett, 1991). The score chi-square statistic was used to compare treatment effect vs. control for each sampling interval. This test is based on the generalized score function (Rotnitzky & Jewell, 1990; Boos, 1992). A GENMOD procedure (SAS Institute, 1997) was used for fitting over-dispersed Poisson generalized linear model. A LOGISTIC procedure was used for fitting proportional odds model to the frass index, and R XTABS routine with R MOSAICPLOTS routine (Development Core Team, 2004) were used for the Pearson's chi-squared test. The tree mortality status was also assessed by the R XTABS routine using the Pearson chi-squared test for independence.

**Table 1** Stand characteristics of treated and control trees

Stand	d.b.h. (cm)	Live crown ratio (%)	Basal area (m <sup>2</sup> /ha)
Treated	31.25 ± 4.20 <sup>a</sup>	70.75 ± 15.58 <sup>a</sup>	29.02 ± 15.50 <sup>a</sup>
Control	34.30 ± 5.68 <sup>a</sup>	68.25 ± 17.04 <sup>a</sup>	28.69 ± 15.17 <sup>a</sup>

Means ± SD in the same columns followed by the same superscript letter are not significantly different at  $\alpha = 0.05$ , Chi-square test


d.b.h., Diameter at breast height; live crown ratio, ratio of height of tree to live crown to total height of tree, expressed as a percentage; basal area, square meters of tree trunk per ha ( $P > 0.05$  for each of the three comparisons).

## Results

Treated trees had significantly lower attack rates by *D. ponderosae* at 2, 4, 6 and 8 weeks after application of flakes (Fig. 1), with mean attack densities reduced to near zero for treated trees at each sampling interval ( $P < 0.0001$ ). No single treated tree was attacked by *D. valens*, whereas control trees averaged nearly two *D. valens* attacks per tree, 8 weeks after treatment, resulting in a significant treatment effect for this pest species too (Fig. 2) ( $P < 0.0001$ ). The dry frass index, used to predict ultimate tree mortality, was significantly higher in control trees than treated trees for all sampling intervals ( $P < 0.0001$ ) (Fig. 3) and, at week 8, the proportional odds model showed that the control trees had a significantly poorer prognosis than treated trees ( $P < 0.0001$ ). Ten months after application, treated trees showed significantly lower mountain pine beetle mortality than control trees (Fig. 4) ( $P < 0.0001$ ), with no mortality in treated trees and no definite survivors in untreated trees. The dry frass index was a highly reliable indicator of ultimate tree mortality (Fig. 5) ( $P < 0.0001$ ), indicating that this measurement may provide a useful short-term predictor for long-term survival of trees under attack by bark beetles.

## Discussion


Several previous studies have demonstrated efficacy of other verbenone formulations for protection of both individual trees and forest stands from attack by *D. ponderosae* (Lindgren & Miller, 2002; Kegley *et al.*, 2003; Progar, 2003; Gibson & Kegley, 2004; Kegley & Gibson, 2004). Those studies utilized pouch or bubble cap release devices applied singly to trees and resulting in a rate of 800 mg AI/tree for bubble caps and 4.65 g/AI/tree for pouches, compared with 15 g/AI/tree for the flake formulation. The reduction in rate of attack by *D. ponderosae* and *D. valens* in the present study was rather striking and was significant at each sampling interval, lasting for at least 8 weeks (Figs 1, 2). This result was considerably better than in some previous studies using both bubble caps and pouches (Lister *et al.*, 1990; Progar, 2005). The improved efficacy observed with the flake formulation in the present study may be simply a result of higher application rates/tree using flakes, or may result from improved pheromone distribution with release of pheromone from



**Figure 1** Mean density of attack by *Dendroctonus ponderosae* (number of attacks/cm<sup>2</sup> of bark) at 2-week intervals after treatment. Shaded boxes represent the interquartile range; whiskers represent the range excluding outliers and extreme values; outliers are shown as open circles, and extreme values are shown as asterisks. Rate of attack on treated trees was significantly different from controls ( $P < 0.0001$ ) for every sampling interval (score chi-square statistic).

many point-sources over a much larger surface area (i.e. 2.1 m<sup>2</sup> for flakes vs. 0.070 m<sup>2</sup> for bubble caps). Bentz *et al.* (2005) suggest that higher emerging beetle populations may overwhelm the standard verbenone pouch treatment, but further research will be needed to determine whether the increased verbenone release rate or the more dispersed pheromone release pattern was responsible for the improved efficacy observed in the present study.

The increased rate of attack by *D. valens* in untreated trees vs. treated trees may have resulted from: (1) an increase in host attractants released by *D. ponderosae* attacks in those untreated trees (Hobson *et al.*, 1993); (2) pheromones released by attacking *D. ponderosae*; or (3) the anti-aggregation effects of the verbenone treatment itself. Verbenone was




**Figure 2** Mean density of attack by *Dendroctonus valens*/tree (number of attacks/cm<sup>2</sup> of bark) at 2-week intervals after treatment. Shaded boxes represent the interquartile range; whiskers represent the range excluding outliers, and outliers are shown as open circles. Rate of attack on treated trees was significantly different from controls ( $P < 0.0001$ ) for each sampling period (score chi-square statistic).




**Figure 3** Contingency table representation of dry frass indices for four sampling intervals after treatment. White, no dry frass; grey, one to ten attacks with dry frass; black, more than ten attacks with dry frass. Treatment had a significant effect on dry frass indices (Pearson's chi-squared test,  $P < 0.0001$ ).

previously shown to inhibit attraction of *D. valens* to its hosts (Rappaport *et al.*, 2001; Sun *et al.*, 2003), and we believe that this outcome is probably a result of both the verbenone treatment and the induction of other semiochemicals, such as frontalin, exo-brevicommin and *cis*- and *trans*-verbenol, by attacking congeners (Pureswaran *et al.*, 2000), as has been shown for *D. valens* after attacks by *D. brevicomis* LeConte (D. R. Owen, unpublished data). Indeed, Sánchez Martínez & Wagner (2002) reported that *D. valens* was an indicator of unmanaged stands in a ponderosa pine ecosystem in northern Arizona, U.S.A., suggesting that the effect may be reflective of increased tree stress and subsequent changes in the semiochemical environment.

The proportion of trees with 10 or more attacks with dry frass increased slightly over time in both the control and treated trees, but reached a plateau at four weeks in only treated trees (not controls) (Fig. 3), suggesting that an effective dose of verbenone was released from flakes during this



**Figure 4** Frequency distribution of trees in mortality categories. Categories are Alive, Uncertain and Dead (as defined in the text). There was a significant treatment effect on mortality categories (Pearson's test,  $P < 0.0001$ ).



**Figure 5** Numbers of trees in mortality categories plotted against dry frass index (dry frass categories: 0 = no dry frass; 1 = one to ten attacks with dry frass; 2 = more than attacks with dry frass) (Pearson's chi-squared test of independence,  $P < 0.0001$ ).

eight week portion of beetle flight. A related study of aerially applied Verbenone DISRUPT® flakes, conducted nearby during the same year, indicated that the flakes continued to emit behaviorally significant amounts of verbenone for at least 14 weeks following treatment (Gillette et al., unpublished data), lending support to that conclusion. The dry frass index may prove to be very useful as an early indicator for assessing the effects of other tree-killing bark beetle species, as well. This method facilitates estimates of ultimate tree mortality shortly following bark beetle attack, without waiting a year or more for obvious tree death and without resorting to destructive sampling methods such as removal of tree bark to examine the status of bark beetle galleries.

The final tree mortality counts (Fig. 4) revealed a striking treatment effect, with treatment dramatically improving the chance of survival for a tree. Ninety percent of treated trees were alive, 10% of treated trees were of uncertain status and none were dead, whereas 70% of control trees were clearly dead, 30% of control trees were of uncertain status and none were alive. The treated trees clearly suffered a few beetle attacks (Figs 1, 2), but were able to defend themselves by pitching out the beetles with resin. These results, indicating that verbenone DISRUPT® flakes were able to inhibit aggregation to a very strong cue, are quite promising. By the time of the final tree mortality count, the flakes were apparently depleted because there was evidence of fresh (spring or early summer of 2005) beetle attacks on treated trees, but the trees were producing copious resin and were effectively repelling the attacks. In addition to the immediate tree mortality data, which show a clear and strong treatment effect, we noted that there were four 'spill-over' attacks on trees adjacent to trees baited with aggregation pheromone. All four of the spill-over attacks occurred adjacent to control trees, and none of them occurred next to treated trees. There is thus some evidence of an analogous spill-over protective effect of the verbenone flakes, which is encouraging in view of the desirability of achieving an area-wide protective effect.

Pouch and bubble cap release devices, when effective, are particularly useful in situations where access is easy and terrain smooth. On the other hand, flakes are amenable to aerial

application, and thus have advantages in remote areas. The flake formulation is one of the few new release devices for verbenone in the last decade. Shea *et al.* (1992) reported one success using a polyolefin bead formulation of verbenone, but they report that they were unable to replicate that success. Holsten *et al.* (2000) later showed that the bead formulation released verbenone for only 2 weeks after application to forest stands, which may explain the inconsistent results achieved with that formulation. Sun *et al.* (2003) showed efficacy of a sprayable, microencapsulated formulation of verbenone for protection of Chinese pine *Pinus tabuliformis* Carrière for protection from *D. valens* attack, but that formulation is no longer commercially available.

The results of the present study suggest that either a higher release rate or a larger release surface may provide better protection than some previous formulations, but further studies must be conducted, especially considering the inconsistent results obtained in some previous studies with verbenone (Shea *et al.*, 1992; Progar, 2005). It is promising that this single-component release system has shown efficacy for one important forest pest, *D. ponderosae*, and one secondary pest, *D. valens*, which occasionally acts as a primary tree-killer (Sun *et al.*, 2003; Eaton & Rodriguez Lara, 1967; Rappaport *et al.*, 2001). Although multiple component formulations using verbenone with ipsdienol (Bertram & Paine, 1994) or angiosperm or green-leaf volatiles may provide enhanced efficacy over verbenone alone for some beetle species (Huber & Borden, 2001; Borden *et al.*, 2003; Pureswan & Borden, 2004), each additional component poses further regulatory and manufacturing obstacles that may limit product development in what is widely considered to be a 'niche' market. On the other hand, many of the nonhost volatiles are considerably cheaper than verbenone, a factor that may outweigh the costs of nontarget testing and maintenance of regulatory approval for multiple component formulations. Single component products, to the extent that they provide effective control with minimal regulatory effort, are nevertheless obviously advantageous. The evidence presented herein suggests that the efficacy of verbenone for management of bark beetles may be considerably enhanced simply by deploying it at higher release rates and/or from different types of release devices. However, it would be a simple matter to include adjuvant flakes in cases where verbenone alone does not provide sufficient protection. Verbenone is an interruptant for a wide range of *Dendroctonus* spp. (Skillen *et al.*, 1997; Rappaport *et al.*, 2001), and this broad activity makes it especially promising as a candidate for managing bark beetles. On the other hand, it should be noted that we conducted two similar trials of the flake formulation for individual tree protection against attack by *D. brevicomis*, with no treatment effect whatsoever (N. E. Gillette *et al.*, unpublished data). Although the individual tree protection tests using verbenone flakes with *D. brevicomis* were not promising, recent aerial application tests of the flakes for *D. brevicomis* control in *P. ponderosa* stands were very encouraging (N. E. Gillette *et al.*, unpublished data). The verbenone flake formulation is one of only two new bark beetle pheromone formulations in more than a decade with the potential for aerial as well as ground application (Sun *et al.*, 2003), and the release device is already

registered for use with pheromones of forest Lepidoptera (Sharov *et al.*, 2002). Although more testing is clearly required to confirm the efficacy shown in this study, changing forest stand conditions (Hessburg *et al.*, 2000) and climate trends (Logan & Powell, 2001; Breshears *et al.*, 2005; Wilent, 2005) suggest a continuing need for large, area-wide treatments for bark beetle management.

## Acknowledgements

We thank Charlie Brown (Fruit Growers Supply, Hilt, California, U.S.A.) for providing access to private lands to conduct this study. Thomas Young and Kelly Conner (Fruit Growers Supply, Hilt, California, U.S.A.) and James Nelson (Hercon Environmental, Wenatchee, Washington, U.S.A.) assisted with installation of the study. Bob Carlson (USDA Forest Service, Redding, California, U.S.A.) assisted with post-treatment assessments. James Heath (Hercon Environmental, Emigsville, Pennsylvania, U.S.A.) provided verbenone flakes to our specifications and provided helpful advice on release curves, application methods, and formulation procedures. Steven Burke (S. E. Burke Consulting, Surrey, British Columbia, Canada) provided valuable assistance with aspects of formulation. Dan Miller (USDA Forest Service, Athens, Georgia, U.S.A.) and Eric Smith (USDA Forest Service, Fort Collins, Colorado, U.S.A.) provided very helpful reviews of the manuscript. Mention of trade names does not constitute endorsement by the USDA Forest Service.

## References

- Agresti, A. (1996) *An Introduction to Categorical Analysis*. John Wiley & Sons Inc., New York, New York.
- Amman, G.D. & Logan, J.A. (1998) Silvicultural control of mountain pine beetle: prescriptions and the influence of microclimate. *American Entomologist*, **44**, 166–177.
- Bentz, B.J., Kegley, S., Gibson, K. & Thier, R. (2005) A test of high-dose verbenone for stand-level protection of lodgepole and whitebark pine from mountain pine beetle (Coleoptera: Curculionidae: Scolytinae) attacks. *Journal of Economic Entomology*, **98**, 1614–1621.
- Bertram, S.L. & Paine, T.D. (1994) Influence of aggregation inhibitors (verbenone and ipsdienol) on landing and attack behavior of *Dendroctonus brevicomis* (Coleoptera: Scolytidae). *Journal of Chemical Ecology*, **20**, 1617–1629.
- Boos, D. (1992) On generalized score tests. *American Statistician*, **46**, 327–333.
- Borden, J.H. (1997) Disruption of semiochemical-mediated aggregation in bark beetles. *Insect Pheromone Research: New Directions* (ed. by R.T. Cardé and A.K. Minks), pp. 421–438. Chapman & Hall, New York, New York.
- Borden, J.H., Chong, L.J., Earle, T.J. & Huber, D.P.W. (2003) Protection of lodgepole pine from attack by the mountain pine beetle, *Dendroctonus ponderosae* (Coleoptera: Scolytidae) using high doses of verbenone in combination with nonhost bark volatiles. *Forestry Chronicle*, **79**, 685–691.
- Breshears, D.D., Cobb, N.S., Rich, P.M. *et al.* (2005) Regional vegetation die-off in response to global-change-type drought. *Proceedings of the National Academy of Sciences of the U.S.A.*, **102**, 15144–15148.
- Buttery, R.G., Light, D.M., Nam, Y.L., Merrill, G.B. & Roitman, J.N. (2000) Volatile components of green walnut husks. (2000). *Journal of Agricultural and Food Chemistry*, **48**, 2858–2861.
- Cerezke, H.F. (1995) Egg gallery, brood production, and adult characteristics of mountain pine beetle, *Dendroctonus ponderosae* Hopkins (Coleoptera: Scolytidae), in three pine hosts. *The Canadian Entomologist*, **127**, 955–965.
- Clarke, S.R., Salom, S.M., Billings, R.F., Berisford, C.W., Upton, W.W., McClellan, Q.C. & Dalusky, M.J. (1999) A scentsible approach to controlling southern pine beetles: two new tactics using verbenone. *Journal of Forestry*, **97**, 26–31.
- Collett, D. (1991) *Modelling Binary Data*. Chapman & Hall, New York, New York.
- Eaton, B. & Rodriguez Lara, R. (1967) Red turpentine beetle *Dendroctonus valens* LeConte. *Important Forest Insects and Diseases of Mutual Concern to Canada, the United States, and Mexico* (ed. by A.G. Davidson and R.M. Prentice), pp. 21–24. Canadian Department of Forest and Rural Development, Canada.
- Elkin, C.M. & Reid, M.L. (2004) Attack and reproductive success of mountain pine beetles (Coleoptera: Scolytidae) in fire-damaged lodgepole pines. *Environmental Entomology*, **33**, 1070–1080.
- Fettig, C.J., McKelvey, S.R. & Huber, D.P. (2005) Nonhost angiosperm volatiles and verbenone disrupt response of western pine beetle, *Dendroctonus brevicomis* (Coleoptera: Scolytidae), to attractant-baited traps. *Journal of Economic Entomology*, **98**, 2041–2048.
- Fiddler, G.O., Hart, D.R., Fiddler, T.A. & McDonald, P.M. (1989) *Thinning Decreases Mortality and Increases Growth of Ponderosa Pine in Northeastern California*. Research Paper PSW-94. Pacific Southwest Forest and Range Experiment Station, Berkeley, California.
- Fournier, G., Hadjiakhoondi, M., Leboeuf, M., Cave, A. & Charles, B. (1997) Essential oils of Annonaceae. Part VII. Essential oils of *Monanthotaxis diclina* (Sprague) Verdcourt and *Unonopsis guaterioides* R.E. Fries. *Flavour and Fragrance Journal*, **12**, 95–98.
- Friendly, M. (1994) Mosaic displays for multi-way contingency tables. *Journal of the American Statistical Association*, **89**, 190–200.
- Furniss, R.L. & Carolin, V.M. (1977) *Western Forest Insects*. USDA Forest Service Miscellaneous Publication 273. USDA Forest Service, Washington, District of Columbia.
- Ghannadi, A. & Zolfaghari, B. (2003) Compositional analysis of the essential oil of *Lallemantia royleana* (Benth. in Wall.) Benth. from Iran. *Flavour and Fragrance Journal*, **18**, 237–239.
- Gibson, K. & Kegley, S. (2004) *Testing the Efficacy of Verbenone in Reducing Mountain Pine Beetle Attacks in Second-Growth Ponderosa Pine*. Forest Health Protection Report, Northern Region. USDA Forest Service, Missoula, Montana.
- Guillen, M.D. & Cabo, N. (1996) Characterisation of the essential oils of some cultivated aromatic plants of industrial interest. *Journal of the Science of Food and Agriculture*, **70**, 359–363.
- Haverty, M.I., Shea, P.J., Hoffman, J.T., Wenz, J.M. & Gibson, K.E. (1998) *Effectiveness of Esfenvalerate, Cyfluthrin, and Carbaryl in Protecting Individual Lodgepole Pines and Ponderosa Pines from Attack by Dendroctonus spp.* Research Paper, Pacific Southwest Research Station PSW-RP-237. USDA Forest Service, Berkeley, California.
- Hessburg, P.F., Smith, B.G., Slater, R.B., Ottman, R.D. & Alvarado, E. (2000) Recent changes (1930s–1990s) in spatial patterns of interior northwest forests, USA. *Forest Ecology and Management*, **136**, 53–83.
- Hobson, K.R., Wood, D.L., Cool, L.G., White, P.R., Ohtsuka, T., Kubo, I. & Zavarin, E. (1993) Chiral specificity in responses by

- the bark beetle *Dendroctonus valens* to host kairomones. *Journal of Chemical Ecology*, **19**, 1837–1846.
- Holsten, E.H., Webb, W., Shea, P.J. & Werner, W.A. (2000) *Release Rates of Methylcyclohexenone and Verbenone From Bubblecap and Bead Releasers Under Field Conditions Suitable for the Management of Bark Beetles in California, Oregon, and Alaska*. USDA Forest Service Research Paper PNW-RP-544. USDA Forest Service, Portland, Oregon.
- Huber, D.P.W. & Borden, J.H. (2001) Protection of lodgepole pines from mass attack by mountain pine beetle, *Dendroctonus ponderosae*, with nonhost angiosperm volatiles and verbenone. *Entomologia Experimentalis et Applicata*, **99**, 131–141.
- Johnstone, W.D. (2002) *Thinning Lodgepole Pine in Southeastern British Columbia: 46-Year Results*. Working Paper, Ministry of Forests, Forest Science Program, British Columbia, Vol. 63, p. 12. Ministry of Forests, Canada.
- Jones, M.E., Paine, T.D., Fenn, M.E. & Poth, M.A. (2004) Influence of ozone and nitrogen deposition on bark beetle activity under drought conditions. *Forest Ecology and Management*, **200**, 67–76.
- Kainulainen, P. & Holopainen, J.K. (2002) Concentrations of secondary compounds in Scots pine needles at different stages of decomposition. *Soil Biology and Biochemistry*, **34**, 37–42.
- Keane, R.E. (2001) Can the fire-dependent whitebark pine be saved? *Fire Management Today*, **61**, 17–20.
- Kegley, S. & Gibson, K. (2004) *Protecting Whitebark Pine Trees From Mountain Pine Beetle Attack Using Verbenone*. Forest Health Protection Report, Northern Region. USDA Forest Service, Missoula, Montana.
- Kegley, S., Gibson, K., Schwandt, J. & Marsden, M. (2003) *A Test of Verbenone to Protect Individual Whitebark Pine From Mountain Pine Beetle Attack*. Forest Health Protection Report, Northern Region. USDA Forest Service, Missoula, Montana.
- Kostyk, B.C., Borden, J.H. & Gries, G. (1993) Photoisomerization of antiaggregation pheromone verbenone: biological and practical implications with respect to the mountain pine beetle, *Dendroctonus ponderosae* Hopkins (Coleoptera: Scolytidae). *Journal of Chemical Ecology*, **19**, 1749–1759.
- Lindgren, B.S. & Miller, D.R. (2002) Effect of verbenone on five species of bark beetles (Coleoptera: Scolytidae) in lodgepole pine forests. *Environmental Entomology*, **3**, 759–765.
- Lister, C.K., Schmid, J.M., Mata, S.A., Haneman, D., O'Neil, C., Pasek, J. & Sower, L. (1990) *Verbenone Bubble Caps Ineffective as a Preventive Strategy Against Mountain Pine Beetle Attacks in Ponderosa Pine*. Research Note RM-501, Rocky Mountain Forest and Range Experiment Station. USDA Forest Service, Ogden, Utah.
- Logan, J.A. & J.A. Powell (2001) Ghost forests, global warming, and the mountain pine beetle (Coleoptera: Scolytidae). *American Entomologist*, **47**, 160–173.
- McCulloch, C.E. & Searle, S.R. (2001) *Generalized, Linear, and Mixed Models*. John Wiley & Sons, Inc., New York, New York.
- McHugh, C.W., Kolb, T.E. & Wilson, J.L. (2003) Bark beetle attacks on ponderosa pine following fire in Northern Arizona. *Environmental Entomology*, **32**, 510–522.
- Molyneux, R.J., Stevens, K.L. & James, L.F. (1980) Chemistry of toxic range plants: volatile constituents of broomweed (*Gutierrezia sarothrae*). *Journal of Agricultural and Food Chemistry*, **28**, 1332–1333.
- Naumann, K. & Rankin, L.J. (1999) Pre-attack systemic applications of a neem-based insecticide for control of the mountain pine beetle, *Dendroctonus ponderosae* Hopkins (Coleoptera: Scolytidae). *Journal of the Entomological Society of British Columbia*, **96**, 13–19.
- Negron, J.F. & Popp, J.B. (2004) Probability of ponderosa pine infestation by mountain pine beetle in the Colorado Front Range. *Forest Ecology and Management*, **191**, 17–27.
- Payne, T.L. & Billings, R.F. (1989) Evaluation of (S)-verbenone applications for suppressing southern pine beetle (Coleoptera: Scolytidae) infestations. *Journal of Economic Entomology*, **82**, 1702–1708.
- Payne, T.L., Billings, R.F., Berisford, C.W., Salom, S.M., Grosman, D.M., Dalusky, M.J. & Upton, W.W. (1992) Disruption of *Dendroctonus frontalis* (Col., Scolytidae) infestations with an inhibitor pheromone. *Journal of Applied Entomology*, **114**, 341–347.
- Perkins, D.L. & Roberts, D.W. (2003) Predictive models of white-bark pine mortality from mountain pine beetle. *Forest Ecology and Management*, **174**, 495–510.
- Pintore, G., Usai, M., Bradesi, P. et al. (2002) Chemical composition and antimicrobial activity of *Rosmarinus officinalis* L. oils from Sardinia and Corsica. *Flavour and Fragrance Journal*, **17**, 15–19.
- Progar, R.A. (2003) Verbenone reduces mountain pine beetle attack in lodgepole pine. *Western Journal of Applied Forestry*, **18**, 229–232.
- Progar, R.A. (2005) Five-year operational trial of verbenone to deter mountain pine beetle (*Dendroctonus ponderosae*; Coleoptera: Scolytidae) attack of lodgepole pine (*Pinus contorta*). *Environmental Entomology*, **34**, 1402–1407.
- Pureswaran, D.S. & Borden, J.H. (2004) New repellent semiochemicals for three species of *Dendroctonus* (Coleoptera: Scolytidae). *Chemecology*, **14**, 67–75.
- Pureswaran, D.S., Gries, R., Borden, J.H. & Pierce, J.D. Jr (2000) Dynamics of pheromone production and communication in the mountain pine beetle, *Dendroctonus ponderosae* Hopkins, and the pine engraver, *Ips pini* (Say) (Coleoptera: Scolytidae). *Chemecology*, **10**, 153–168.
- R. Development Core Team. (2004) *R: a language and environment for statistical computing*. <http://www.r-project.org/> [accessed on 3 August 2005].
- Rappaport, N.G., Owen, D.R. & Stein, J.D. (2001) Interruption of semiochemical-mediated attraction of *Dendroctonus valens* (Coleoptera: Scolytidae) and selected nontarget insects by verbenone. *Environmental Entomology*, **30**, 837–841.
- Robles, C., Bousquet-Melou, A., Garzino, S. & Bonin, G. (2003) Comparison of essential oil composition of two varieties of *Cistus ladanifer*. *Biochemical Systematics and Ecology*, **31**, 339–343.
- Rotnitzky, A. & Jewell, N.P. (1990) Hypothesis testing of regression parameters in semiparametric generalized linear models for cluster correlated data. *Biometrika*, **77**, 485–497.
- Safranyik, L., Shore, T.L., Carroll, A.L. & Linton, D.A. (2004) Bark beetle (Coleoptera: Scolytidae) diversity in spaced and unmanaged mature lodgepole pine (Pinaceae) in southeastern British Columbia. *Forest Ecology and Management*, **200**, 23–38.
- Salom, S.M., Grosman, D.M., McClellan, Q.C. & Payne, T.L. (1995) Effect of an inhibitor-based suppression tactic on abundance and distribution of the southern pine beetle (Coleoptera: Scolytidae) and its natural enemies. *Journal of Economic Entomology*, **88**, 1703–1716.
- Sánchez Martínez, G. & Wagner, M.R. (2002) Bark beetle community structure under four ponderosa pine forest stand conditions in northern Arizona. *Forest Ecology and Management*, **170**, 145–160.
- SAS Institute. (1997) *SAS/STAT Software: Changes and Enhancements Through Release 6.12*. SAS Institute, Inc., Cary, North Carolina.
- Schwandt, J. & Kegley, S. (2004) *Mountain pine beetle, blister rust, and their interaction on whitebark pine at Trout Lake and Fisher Peak in northern Idaho from 2001 to 2003*. Forest Health Protection Report, Northern Region. USDA Forest Service, Missoula, Montana.
- Sefidkon, F., Jalili, A. & Mirhaji, T. (2002) Essential oil composition of three *Artemisia* spp. from Iran. *Flavour and Fragrance Journal*, **17**, 150–152.

- Sharov, A.A., Leonard, D., Liebhold, A.M., Roberts, E.A. & Dickerson, W. (2002) 'Slow the spread': a national program to contain the gypsy moth. *Journal of Forestry*, **100**, 30–35.
- Shea, P.J., McGregor, M.D. & Daterman, G.E. (1992) Aerial application of verbenone reduces attack of lodgepole pine by mountain pine beetle. *Canadian Journal of Forest Research*, **22**, 436–441.
- Shore, T.L., Safranyik, L. & Lemieux, J.P. (2000) Susceptibility of lodgepole pine stands to the mountain pine beetle: testing of a rating system. *Canadian Journal of Forest Research*, **30**, 44–49.
- Skillen, E.L., Berisford, C.W., Camann, M.A. & Reardon, R.C. (1997) *Semiochemicals of Forest and Shade Tree Insects in North America and Management Applications*. Forest Health Technology Enterprise Team, FHTET 96–15. USDA Forest Service, Morgantown, West Virginia.
- Smith, E.L., McMahan, A.J. & Wang, G.Z. (2002) *Eco-Physiology Approach to Projecting Tree Stress and Vigor in FVS*. Proceedings of the Rocky Mountain Research Station, RMRS-P-25, pp. 146–150. USDA Forest Service, Ogden, Utah.
- Sun, J.H., Gillette, N.E., Miao, Z.W., Kang, L., Zhang, Z.N., Owen, D.R. & Stein, J.D. (2003) Verbenone inhibits attraction to host volatiles and reduces attack by *Dendroctonus valens* LeConte (Coleoptera: Scolytidae) on *Pinus tabulaeformis* in China. *Canadian Entomologist*, **135**, 721–732.
- Taylor, S.P., Wilson, I.M. & White, K.J. (1997) Topical application of carbon dioxide and liquid nitrogen against the mountain pine beetle, *Dendroctonus ponderosae* (Coleoptera: Scolytidae). *Journal of the Entomological Society of British Columbia*, **94**, 63–66.
- Tomback, D.F., Arno, S.F. & Keane, R.E. (2001) *Whitebark Pine Communities: Ecology and Restoration*. Island Press, Washington, District of Columbia.
- Umamo, K., Hagi, Y., Nakahara, K., Shoji, A. & Shibamoto, T. (2000) Volatile chemicals identified in extracts from leaves of Japanese mugwort (*Artemisia princeps* Pamp.). *Journal of Agricultural and Food Chemistry*, **48**, 3463–3469.
- Waters, W.E. (1985a) The pine-bark beetle ecosystem: A pest management challenge. *Integrated Pest Management in Pine-Bark Beetle Ecosystems* (ed. by W.W. Waters, R.W. Stark and D.L. Wood), pp. 49–60. John Wiley & Sons, Inc., New York, New York.
- Waters, W.E. (1985b) Monitoring bark beetle populations and beetle-caused damage. *Integrated Pest Management in Pine-Bark Beetle Ecosystems* (ed. by W.W. Waters, R.W. Stark and D.L. Wood), pp. 141–176. John Wiley & Sons, Inc., New York, New York.
- Whitehead, R.J. & Russo, G.L. (2005) 'Beetle-Proofed' Lodgepole Pine Stands in Interior British Columbia Have Less Damage From Mountain Pine Beetle. Information Report BC-X-402. Pacific Forestry Centre, Canadian Forest Service, Canada.
- Wilent, S. (2005) Mountain pine beetles threaten Canadian, US forest. *The Forestry Source*. [http://www.safnet.org/archive/0505\\_beetle.cfm](http://www.safnet.org/archive/0505_beetle.cfm) [accessed on 28 February 2006].
- Wilson, I.M., Borden, J.H., Gries, R. & Gries, G. (1996) Green leaf volatiles as antiaggregants for the mountain pine beetle, *Dendroctonus ponderosae* Hopkins (Coleoptera: Scolytidae). *Journal of Chemical Ecology*, **22**, 1861–1875.
- Wood, D.L., Stark, R.W., Waters, W.W., Bedard, W.D. & Cobb, F.W. Jr (1985) Treatment tactics and strategies. *Integrated Pest Management in Pine-Bark Beetle Ecosystems* (ed. by W.W. Waters, R.W. Stark and D.L. Wood), pp. 121–140. John Wiley & Sons, Inc., New York, New York.
- Wood, D.L., Koerber, T.W., Scharpf, R.F. & Storer, A.J. (2003) *Pests of the Native California Conifers*. University of California Press, Berkeley, California.
- Wulder, M.A., Seemann, D., Dymond, C.C., Shore, T. & Riel, B. (2004) *Arc/Info Macro Language (AML) Scripts for Mapping Susceptibility and Risk of Volume Losses to Mountain Pine Beetle in British Columbia*. Technology Transfer Note. Pacific Forestry Centre, Canadian Forest Service, Canada.
- Zausen, G.L., Kolb, T.E., Bailey, J.D. & Wagner, M.R. (2005) Long-term impacts of stand management on ponderosa pine physiology and bark beetle abundance in northern Arizona: a replicated landscape study. *Forest Ecology and Management*, **218**, 291–305.

Accepted 4 May 2006