

**First occurrence of the goldspotted oak borer parasitoid,  
*Calosota elongata* (Hymenoptera: Eupelmidae), in California**

Author(s): Laurel J. Haavik, Tom W. Coleman, Yigen Chen, Michael I. Jones,  
Robert C. Venette, Mary L. Flint, and Steven J. Seybold

Source: Pan-Pacific Entomologist, 88(2):274-276. 2012.

Published By: Pacific Coast Entomological Society

DOI: <http://dx.doi.org/10.3956/2012-08.1>

URL: <http://www.bioone.org/doi/full/10.3956/2012-08.1>

---

BioOne ([www.bioone.org](http://www.bioone.org)) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/page/terms\\_of\\_use](http://www.bioone.org/page/terms_of_use).

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

## Scientific Note

### **First occurrence of the goldspotted oak borer parasitoid, *Calosota elongata* (Hymenoptera: Eupelmidae), in California**

*Calosota elongata* Gibson (Hymenoptera: Eupelmidae) is a gregarious, ectoparasitic larval parasitoid that was described recently (Gibson 2010) in association with the goldspotted oak borer, *Agrilus coxalis* Waterhouse [now considered to be *Agrilus auroguttatus* Schaeffer (Coleoptera: Buprestidae)] in its native range in southeastern Arizona (Coleman & Seybold 2011). We have discovered this parasitoid outside of its presumptive native range on 1 December 2010 near the Pine Creek Trailhead, Cleveland National Forest in San Diego Co., California, beneath the bark of coast live oak, *Quercus agrifolia* Née (Fagaceae), in association with *A. auroguttatus*. Six pupae of *C. elongata* were collected from an *A. auroguttatus* pupal cell, which had been exposed during destructive sampling of *Q. agrifolia* phloem tissue with a hatchet. The pupae were brought to the laboratory where three specimens matured into adult females. Additional collections of *C. elongata* pupae were made in the same manner and in the same general vicinity on: 1) 19 January 2011—3 pupae (Noble Canyon Trailhead, Cleveland National Forest); and 2) 19 December 2011—6 pupae (Pine Creek Trailhead). No host remains were found during the three aforementioned collections, suggesting that the parasitoids had completely consumed the immature *A. auroguttatus*. Typically, host head capsules are found following these bouts of feeding by gregarious parasitic Hymenoptera (Quicke 1997), however in this instance, the head capsules may have been lost during the destructive sampling of the bark. Subsequent flight trapping in the same general vicinity yielded nine adult female specimens of *C. elongata* that flew between 19 May and 12 October 2011. The wasps were trapped primarily on yellow sticky cards on the north and south sides of two mature *Q. agrifolia*, however, one specimen of *C. elongata* was trapped between 8 September and 12 October 2011 on a purple prism-style trap located at the dripline of the foliage of *Q. agrifolia*. The latter traps have been used routinely to trap *Agrilus* species throughout North America (Francesca et al. 2008). The flight period that we have established from the California specimens contains the emergence period that we noted from some of the type material from Arizona. Bark samples collected on 1–2 May 2009 from Box Canyon (Santa Rita Mountains, Pima Co., Arizona) yielded 12 adults emerging between 6 and 14 July 2009 and 7 adults emerging between 14 July and October 2009 (Coleman & Seybold 2011). Of the 19 specimens from that collection two were males. There were no males among the 12 specimens collected in California, suggesting that this may be a thelytokous population or simply that no males were taken in this small sample from California. We suspect that *C. elongata* was introduced to California with the original population of *A. auroguttatus* because these first California collection records for the wasp are in the area that we consider as the hypothetical point of introduction of the California population of *A. auroguttatus* from Arizona (Coleman et al. 2012a). Also supporting some level of founder effects are preliminary observations of the rate of parasitism, which is higher (15% vs. < 1%) in Arizona than in California (Coleman et al. 2012b).

Record—CALIFORNIA: San Diego Co., Pine Creek Trailhead, Cleveland National Forest, Descanso Ranger District, approx. 1085 m elev., 32.83469° N,

116.54345° W, XII-1-2010, coll. L. J. Haavik, ex: outer bark of coast live oak tree, *Quercus agrifolia*. Collected as pupae in the pupal cells of goldspotted oak borer, *Agrilus auroguttatus*, and reared to adults in laboratory.

Record—CALIFORNIA: San Diego Co., Pine Creek Trailhead, Cleveland National Forest, Descanso Ranger District, 1088 m elev., 32.833031° N, 116.547049° W, V-19- to VI-14-2011, coll. T. W. Coleman, S. J. Seybold, ex: yellow sticky trap placed near coast live oak trees, *Quercus agrifolia*.

We thank G. Gibson [Canadian National Collection (CNC), Ottawa, Ontario, Canada] for verifying the identification of *C. elongata*; L. Walker (UC-Davis, Department of Entomology) for assistance with lab work; and J. C. Lee (USDA ARS, Corvallis, OR) and G. E. Heimpel (University of Minnesota, St. Paul) for pre-submission reviews of the manuscript. Voucher specimens of the adults were placed at the CNC and the California Academy of Sciences, San Francisco, California. We acknowledge the contributions of R. L. (“Dick”) Penrose to the discovery of *A. auroguttatus* in California. Dick coordinated the statewide bark beetle and woodborer survey that in June 2004 yielded the first specimen of *A. auroguttatus* (then *A. coxalis*) captured in the state (Westcott 2005). This work was supported by an STDP grant (#R5-2010-02) from the USDA Forest Service, Research and Development Washington Office, by the USDA Forest Service Northern and Pacific Southwest Research Stations, by the USDA Forest Service Forest Health Protection, Region 5 and by the University of California, Davis.

Laurel J. Haavik, *Department of Entomology, University of California Davis, One Shields Ave., Davis, California 95616*; Tom W. Coleman, *USDA Forest Service Forest Health Protection, 602 S. Tippecanoe St., San Bernardino, California 92408*; Yigen Chen, *Department of Entomology, University of California Davis, One Shields Ave., Davis, California 95616*; Michael I. Jones, *Department of Entomology, University of California Davis, One Shields Ave., Davis, California 95616*; Robert C. Venette, *USDA Forest Service Northern Research Station, 1561 Lindig Ave., St. Paul, Minnesota 55108*; Mary L. Flint, *Department of Entomology, University of California Davis, One Shields Ave., Davis, California 95616*; and Steven J. Seybold, *USDA Forest Service Pacific Southwest Research Station, Chemical Ecology of Forest Insects, 720 Olive Dr., Suite D, Davis, California 95616*; e-mail: [sjseybold59@gmail.com](mailto:sjseybold59@gmail.com)

#### LITERATURE CITED

- Coleman, T. W. & S. J. Seybold. 2011. Collection history and comparison of the interactions of the goldspotted oak borer, *Agrilus auroguttatus* Schaeffer (Coleoptera: Buprestidae), with host oaks in southern California and southeastern Arizona. *The Coleopterists Bulletin* 65:93–108.
- Coleman, T. W., A. D. Graves, M. S. Hoddle, Z. Heath, Y. Chen, M. L. Flint & S. J. Seybold. 2012a. Forest stand composition and impacts associated with *Agrilus auroguttatus* Schaeffer (Coleoptera: Buprestidae) and *Agrilus coxalis* Waterhouse in oak woodlands. *Forest Ecology and Management* 276:104–117.
- Coleman, T. W., V. Lopez, R. Stouthamer, P. F. Rugman-Jones, S. J. Seybold, R. Reardon & M. S. Hoddle. 2012b. Can the destruction of California’s oak woodlands be prevented? Potential for biological control of the goldspotted oak borer, *Agrilus auroguttatus*. *BioControl* 57:211–225.
- Francese, J. A., J. B. Oliver, I. Fraser, D. R. Lance, N. Youssef, A. J. Sawyer & V. C. Mastro. 2008. Influence of trap placement and design on capture of emerald ash borer (Coleoptera: Buprestidae). *Journal of Economic Entomology* 101:1831–1837.

- Gibson, G. A. P. 2010. *Calosota* Curtis (Hymenoptera, Chalcidoidea, Eupelmidae) - review of the New World and European fauna including revision of species from the West Indies and Central and North America. *ZooKeys* 55:1–75.
- Quicke, D. L. J. 1997. *Parasitic Wasps*. Chapman & Hall, London, U.K., 470 pp.
- Westcott, R. L. 2005. A new species of *Chrysobothris* Eschscholtz from Oregon and Washington, with notes on other Buprestidae (Coleoptera) occurring in the United States and Canada. *Zootaxa* 1044:1–15.

*Received 18 Feb 2012; Accepted 20 Mar 2012 by A. H. Smith-Pardo; Publication date 14 Sep 2012.*