

Abronia umbellata ssp. *breviflora* on the Oregon coast: Reintroduction and population monitoring



2018

Report to the Bureau of Land Management, and USDA Forest Service

Report prepared by Denise E.L. Giles and
Thomas N. Kaye
Institute for Applied Ecology



PREFACE

This report is the result of a cooperative agreement between the Institute for Applied Ecology (IAE) and a federal agency. IAE is a non-profit organization dedicated to natural resource conservation, research, and education. Our aim is to provide a service to public and private agencies and individuals by developing and communicating information on ecosystems, species, and effective management strategies and by conducting research, monitoring, and experiments. IAE offers educational opportunities through 3-4 month internships. Our current activities are concentrated on rare and endangered plants and invasive species.



Questions regarding this report or IAE should be directed to:

Matt A. Bahm

Conservation Research Program Director

Institute for Applied Ecology

563 SW Jefferson Avenue

Corvallis, Oregon 97333

phone: 541-753-3099

fax: 541-753-3098

email: mattab@appliedeco.org

ACKNOWLEDGEMENTS

This project would not have been possible without the assistance of Tim Rodenkirk, Jennie Sperling, Kip Wright, (Coos Bay District, BLM), and Matthew Smith, Armand Reibischke (Siuslaw National Forest) and Marty Stein (Retired, Siuslaw National Forest). In 2018, work was supported by IAE staff: Michelle Allen, Matt Bahm, Meaghan Petix, Lisa Schomaker, Michel Wiman, Jennifer Thornhill and Kristina Lopez. A special thank you is also extended to all volunteers who helped to clean and distribute seed including Kenai Kauffman, Boone Kauffman, Dian Cummings, Philip San Romani and Valerie Petersen.

Cover photograph: Pink sand verbena (*Abronia umbellata* ssp. *breviflora*).

Suggested Citation

Giles, D.E.L. and T.N. Kaye. 2018. *Abronia umbellata* ssp. *breviflora* on the Oregon coast: Reintroduction and population monitoring. Report prepared by Institute for Applied Ecology, Corvallis, Oregon for USDA Forest Service, Siuslaw National Forest and USDI Bureau of Land Management, Coos Bay District. x + 52 pp.

TABLE OF CONTENTS

PREFACE.....	II
ACKNOWLEDGEMENTS.....	III
LIST OF FIGURES	VI
LIST OF TABLES	VII
EXECUTIVE SUMMARY.....	IX
INTRODUCTION	1
METHODS	3
Seed Source.....	3
Direct Seeding.....	3
Population Surveys	3
Coos Bay North Spit.....	4
Seeded Transects.....	4
Plant Measurements	4
Overwintering Plants.....	6
Substrate Assessment	Error! Bookmark not defined.
RESULTS.....	10
Population Surveys	10
General Trend	10
Siuslaw National Forest	10
Bureau of Land Management.....	16
Plant Measurements	25
Plant Size and Reproductive Effort	25
Plant Density	25
Plant Community.....	25
Substrate Assessments.....	Error! Bookmark not defined.
DISCUSSION	27
Population trends, plant size and habitat management	27
Substrate Assessments.....	Error! Bookmark not defined.
Transplanting	28
ADDITIONAL RESEARCH.....	28
Importance of a seedbank.....	29
Population modeling to assess recovery objectives	29
Restoration with additional beach species.....	29
Hypotheses for Future Research.....	30

RECOMMENDATIONS FOR 2019	32
Seeding and Habitat Management	32
Population surveys.....	33
LITERATURE CITED	34
APPENDIX I. MATERIALS REQUIRED FOR POPULATION MONITORING.....	36
APPENDIX II. SURVEY RESULTS FOR SEEDED AND NATURAL POPULATIONS	37
APPENDIX III. MAPS OF SELECTED SITES- SEEDED OR MONITORED	49
New River.....	49
Overlook North and South.....	50
Siltcoos.....	51
Tahkenitch.....	52

LIST OF FIGURES

Figure 1. Pink sand-verbena (<i>Abronia umbellata</i> ssp. <i>breviflora</i> ; (from Hitchcock et al 1964).....	1
Figure 2. Historic and current distribution of pink sand-verbena (<i>Abronia umbellata</i> subsp. <i>breviflora</i>) in Oregon.....	2
Figure 3. In 2009, the populated area was divided into three section, 1) the original habitat (southeast), 2) northeast and 3) west of the Foredune Road. Each area was subsampled using the methods described in the text.....	5
Figure 4. Population trends at Siltcoos Creek, 1996 – 2018. An erosion event in the winter of 2012/2013 resulted in significant loss of habitable habitat, thus the decrease in seeding effort and the number of plants observed at the site.....	11
Figure 5. Population trends and seeding history at Tahkentich Creek of pink sand from 2001-2018. This site was not monitored in 2007, 2008 or 2010.....	12
Figure 6. Population trends and seeding history at Overlook North and South of pink sand verbena from 2000-2018. Overlook South was not monitored in 2008.....	13
Figure 7. Population trends at the Coos Bay North Spit reintroduction site. The area was seeded in January 1997 with 50,000 seeds from Port Orford. Population size estimates for 1997-1999 are complete censuses of all plants in the population, while those from 2000-2008 are estimated from a subsample of a reference area and census of the remainder of the population. The error bars for 2000-2008 estimates represent 95% confidence intervals. *In 2008, an estimate of the entire population size was not possible because over 50% of the population had been disced prior to our surveys. See text for a discussion of population trends. In 2009, the sampling methods were modified so that the entire population was sub-sampled (and no areas censused). In this chart the CI for 2009-2019 are not reported because the population estimate is a combination of 3 different populations, each with their own CI.....	17
Figure 8. Estimated number of plants at the coos bay north Spit in each of the three population areas. Note that because the population is small on the west side, the scale differs than that for the north and south. Error bars represent 95% Confidence intervals.....	18
Figure 9. Density of plants per m ² at the Coos Bay North Spit from 2009-2018. the decrease in all areas from 2012-2013 coincided with lack of habitat management. Density measurements for the West side are not available for 2017. Error bars represent 95% Confidence Intervals.....	19
Figure 10. Population trends and seeding history of pink sand verbena at the New River ACEC reintroduction site. Seeding has been conducted in all years except 1998.....	24
Figure 11. Nearly all plants in 2015 at New River were found against the back of the dune in an area that had been overwashed. The area circled in red had hundreds of small pink sand verbena plants, as compared to the single, large reproductive plant in the foreground.	24
Figure 13. Average Length of longest branch vs. pH of substrate samples collected in 2012 and 2015.	Error! Bookmark not defined.
Figure 14. In 2018, Plants were larger with more flowers at Overlook South (left) where new areas had been treated for Beachgrass removal than at the nearby Overlook North site (Right).....	27
Figure 15. The percent of seeds stored from 0-9 years. The equation for the linear regression is $y = -0.03x + 0.955$, $R^2 = 0.81$, $P = 0.015$	31

LIST OF TABLES

Table 1. Reintroduction sites for the Coos Bay BLM District, methods used, and habitats for seeding and transplanting efforts discussed in this report. Seedlings reported here all involved the use of 50,000 seeds per site, except where noted.	7
Table 2. Reintroduction sites of the Siuslaw National Forest, methods used, and habitats for seeding and transplanting efforts discussed in this report. Seedlings reported here all involved the use of 50,000 seeds per site except where noted.	8
Table 3. Number of pink sand verbena at Overlook South and Overlook North 2000-2018. *Overlook South was not monitored in 2008.	14
Table 4. Number of plants observed at Sutton Creek and Baker Beach from 2003 to present. Sites were not monitored in each year.	16
Table 5. Estimated number of pink sand verbena at the Coos Bay North Spit from 2009-2018.	20
Table 6. Average size of plants in 2012-2018 at select sites. “-“ indicates data was not collected in that year. Values in () represent 95% confidence intervals. Systematic Measurements on overwintering plants were only conducted in 2012.	21
Table 7. Average number of flowers in 2012-2018 at Select sites. “-“ indicates data was not collected in that year. Values in () represent 95% confidence intervals. Systematic measurement on overwintering plants were not conducted in every year.	22
Table 8. Count of Plant Species within 0.5m of Randomly Selected pink sand-verbena plants, including percent of pink sand-verbena monitored that had species present, in 2012-2018 at the Coos Bay North Spit. Names in Bold indicate native species. *Because it is difficult to Distinguish between individuals of both American and European Beachgrass, Presence/Absence only was noted.	26
Table 9. Range and average pH at select sites monitored in 2012 and 2015. Note that the sand from the areas with cover of shell greater than 5% had the highest pH.....	Error! Bookmark not defined.
Table 10. Plant species native to Oregon beaches and dunes that are candidates for use in ecosystem restoration projects. More information is needed on techniques for germination and propagation of these taxa.....	Error! Bookmark not defined.
Table 11. Current and suggested seeding rates for pink sand-verbena sites.	32
Table 12. Sites with pink sand-verbena documented in the past 10 years, with sites monitored in 2018 listed in bold.....	33
Table 13. Results of surveys for pink sand-verbena at Coos Bay District Bureau of Land Management sites along the Oregon Coast. IAE conducted surveys from 2000-2018. Population data for years prior to 2000 are from various sources.....	37
Table 14. Results of surveys for pink sand-verbena sites managed by the Siuslaw National Forest along the Oregon Coast. IAE conducted surveys from 2000-2012. Population data for years prior to 2000 are from various sources.....	39
Table 15. Results of surveys for pink sand-verbena sites managed by the Oregon Department of Parks and Recreation along the Oregon Coast. IAE conducted surveys from 2000-2012. Population data for years prior to 2000 are from various sources.....	42
Table 16. Results of surveys for pink sand-verbena sites managed by the USFWS along the Oregon Coast. IAE conducted surveys from 2009-2012.	44

Table 17. Results of surveys for pink sand-verbena sites along the Oregon Coast managed by various land managers including the State Parks Department. IAE conducted surveys from 2000-2012. Population data for years prior to 2000 are from various sources.....45

EXECUTIVE SUMMARY

Pink sand-verbena (*Abronia umbellata* ssp. *breviflora*) is listed as endangered by the Oregon Department of Agriculture, and a Species of Concern by the U.S. Fish and Wildlife Service. Since the late 1970's, the number of natural populations in Oregon has decreased from approximately 10 populations to less than 5, though the current status of populations not included in this report are unknown. The 2006 Conservation Strategy cites surveys conducted from 1986-2003 to establish the number and size of populations present in Oregon. The date of the last rangewide survey in Oregon is not clear, but occurred prior to 2004. The primary threats to the species include competition from European beachgrass (*Ammophila arenaria*) and other weedy beach species, and other human-influenced habitat changes.

This report includes information about pink sand-verbena research along the Oregon coast since 1997, including seeding and transplantation experiments, and population monitoring at several beach and dune habitats.

In 2018 our actions and observations included activities at sites managed by the USDA Forest Service and Bureau of Land Management.

1. USDA Forest Service, Siuslaw National Forest sites

- a. Tahkenitch: ~90,000 seeds were distributed at Tahkenitch in 2018, resulting in a total of 361 plants (117 reproductive). This was the eighth seeding at the site since 2005. In 2014-2018, seeding was focused in the southern 1/3 of the site, as this was the area that had been most successful in previous years.
- b. Siltcoos Creek: ~50,000 seeds were distributed in the habitat still present at the site and 292 plants (147 reproductive) were found in the fall of 2018. This is likely an underestimate as surveying occurred later in the season when some plants had already senesced or been buried by shifting sands.
- c. Overlook: Despite substantial decreases in this population in 2014, this population is now the second largest in Oregon and we strongly recommend continued beachgrass control and seed addition. In 2018 South Overlook was expanded and many robust plants were present in the new habitat.
 - i. North: ~75,000 seeds were distributed, resulting in 497 plants (284 reproductive).
 - ii. South: ~90,000 seeds were distributed resulting in 3,476 plants (1,850 reproductive).
- d. Baker Beach: In 2018, Forest Service staff spread a total of ~35,000 seeds at Baker Beach and Sutton Creek resulting in 28 plants in the area near Sutton Creek. No plants were found at Baker Beach. Prior to 2018, this area had not been seeded since 2009.

2. USDI Bureau of Land Management, Coos Bay District sites

- a. New River: ~100,000 seeds were distributed by BLM staff in four overwashes in the New River ACEC resulting in 196 plants (124 reproductive). This represents a substantial increase from 2017 when only 9 plants were present at the site. As in previous years most plants were found in areas where breaks in the foredune had allowed overwash to occur - highlighting the affinity of pink sand-verbena for areas with repeated disturbance (and possibly nutrient input). Continued seeding at the site to augment the existing seed bank, and management in the form of continued European beachgrass control, as well as control of other non-native species including *Hypochaeris radicata*, *Cakile maritima*, and *Rumex acetosella*, will be vital in maintaining suitable coastal dune habitat for this and other dune species.

- b. Coos Bay North Spit: ~100,000 seeds were distributed by BLM staff. In 2018, it is estimated that there are approximately 46,961 reproductive pink sand-verbena plants on the Coos Bay North Spit. This is the lowest number observed at this site since 2005. Over the course of this study, we have seen an increase in the number and diversity of weed species at this site. This is likely due to soil formation related to sand stabilization and the lack of overwash and salt-spray in the area that contribute to natural soil formation processes on this coastal dune. There was a precipitous drop in the population when beach grass management was discontinued for just one year in 2013 in the southern portion of the population; this portion of the population rebounded when discing resumed in 2014-2018.

Abronia umbellata ssp. breviflora on the Oregon coast: Reintroduction and population monitoring

REPORT TO THE BUREAU OF LAND MANAGEMENT AND USDA FOREST SERVICE
REPORT TO THE BUREAU OF LAND MANAGEMENT AND USDA FOREST SERVICE

INTRODUCTION

Pink sand-verbena (*Abronia umbellata* ssp. *breviflora*; Figure 1) is listed by the Oregon Department of Agriculture as endangered, and it is considered a Species of Concern by the U.S. Fish and Wildlife Service. Historically, the species was known from beaches along the Pacific Coast from Vancouver Island (British Columbia) south to northern California. Until two plants were discovered in Willapa Bay in 2006, the species had been believed to be extinct in Washington. In 2000, a small population (two plants) was rediscovered on Vancouver Island, but this population did not re-establish in subsequent years. Since the late 1970's, the number of natural populations in Oregon has dwindled from around ten down to about five (Figure 2). . The 2006 Conservation Strategy cites surveys conducted from 1986-2003 to establish the number and size of populations present in Oregon. The date of the last rangewide survey in Oregon is not clear, but occurred prior to 2004.

The primary threats to the species include, the stabilization of dunes across the Pacific Northwest coast, competition from non-native species including (but not limited to) European beachgrass (*Ammophila arenaria*) and to a lesser extent European searocket (*Cakile maritima*), American searocket (*Cakile edentula*), as well as habitat disturbance by off road vehicles and the unknown effects of climate change. Shifts in dune vegetation, and substrate chemistry associated with dune stabilization may contribute to observed declines in populations of this rare species, and prevent new populations from establishing in the fragmented dune system of the Oregon Coast.




FIGURE 1. PINK SAND-VERBENA (*ABRONIA UMBELLATA* SSP. *BREVIFLORA*); (FROM HITCHCOCK ET AL 1964).

The decline of pink sand-verbena along the coast is correlated with the reduction in number of many beach species, including native plants and wildlife. For example, the Western Snowy Plover has suffered a well-documented decline (Oregon Department of Fish and Wildlife 1994) that parallels that of pink sand-verbena. There is some historical evidence that snowy plovers used native vegetation, specifically sand-verbenas, for foraging and cover (Gabrielson and Jewett 1970, p.238). Recovery efforts for a variety of these dune species may be most effective if coordinated or combined.

Since the mid-1990's, research has been conducted on methods for reintroduction of pink sand-verbena. Information is still needed on the reintroduction potential of the species in various beach habitats, and an understanding of 'successful' introduction or augmentation sites developed in order to develop a range-wide conservation and recovery plan. The ultimate goal of these reintroduction efforts is to support the requirements of a Conservation Strategy (Kaye 2006) and remove the species from the endangered list.

In this report, we summarize recent seeding and monitoring efforts at several beaches and dune habitats along the Oregon coast managed by the US Forest Service and BLM. This report focuses on recent efforts at the Forest Service sites: Baker Beach Sutton Creek, Tahkenitch Creek, Siltcoos Creek, North and South Overlook); and the Coos Bay District Bureau of Land Management sites, Coos Bay North Spit and the New River Area of Critical Environmental Concern. Information on additional populations, previous seeding and transplanting efforts at various sites can be found in previous year's reports (Giles-Johnson and Kaye 2014).




FIGURE 2. HISTORIC AND CURRENT DISTRIBUTION OF PINK SAND-VERBENA (*ABRONIA UMBELLATA* SUBSP. *BREVIFLORA*) IN OREGON.

METHODS

General seeding and monitoring information is reviewed here. Site-specific information is included with the results of each site, below.

Seed Source

All of the seed materials manually dispersed to sites in Oregon in 1997 through 2018 were collected from populations at Port Orford (1995-1997) or Coos Bay, Oregon (1999-2018). The pink sand-verbena population at Coos Bay was established from seeds originally collected at Port Orford and are therefore of the same genetic lineage (McGlaughlin et al. 2002). Due to the decline in population size at Port Orford, in 1999 we began collecting seeds at the introduced site at Coos Bay North Spit. Seeds have been collected annually with the exception of 2008 when seed collection was not possible as the Coos Bay North Spit was bulldozed and disced in mid-October prior to fruit set for most individuals. These seeds were also the source for transplants grown in greenhouses at Oregon State University and IAE in Corvallis, Oregon. We use screens to sieve mature fruits from the sand. Fruits are collected from at least 20 areas distributed throughout the population. Immature seeds and areas where non-native species are present are avoided. In 2012 and 2013, seed was collected by BLM staff at the North Spit. In 2014-2016, BLM and IAE staff collected seed and BLM staff conducted seed collection in 2017 and 2018. In 2014-2018, seed cleaning included the removal of seeds of weedy species including *Hypochaeris radicata* and *Cakile* spp.

The seeds of pink sand-verbena occur in single-seeded fruits. After collection, fruits were cleaned of residual sand and other foreign material (such as twigs, shells and leaves) and dried. In 1999, groups of 100 fruits were weighed to determine average fruit weight, and groups of 10 fruits were clipped open to determine average seed set. In subsequent years, this information was used to weigh out bags of ~10,000 seeds (154 g) each for dispersal to beaches.

Direct Seeding

More than 15 sites have been seeded since this project began in 1997 (for information about sites not listed in Table 17, see Kaye and Cramer 2005). Seeded sites were revisited the following late summer/early fall to survey for pink sand-verbena plants.

Pink sand-verbena fruits were scattered in February through early April of each year by hand by walking along the beach and slowly releasing handfuls of material until all were dispersed. Seeding rates vary by site and year depending on many factors including the amount of suitable habitat, seed availability and time constraints.

Population Surveys

Beginning in 2000 IAE started surveying known pink sand-verbena habitat. Sites were selected based on population and reintroduction history, as well as time limitations. When populations were found, we documented the number of reproductive and vegetative individuals, and occasionally, measurements of plant size [length of the longest stem, longest width, and perpendicular width (see below)]. In large populations we only counted reproductive plants. Because pink sand-verbena is an annual, vegetative plants do not contribute to future populations. In 2009, due to the increase in the population size at the Coos Bay North Spit, sampling was modified so that the habitable area is divided into three sections and

each is sub-sampled and the population size estimated; these methods are discussed in more detail below.

Coos Bay North Spit

In 2009, the population at Coos Bay North Spit had expanded such that a census of the habitat (as performed from 2000-2008) was no longer feasible. We divided the occupied habitat into three sections and sub-sampled within each area counting only reproductive plants. The three areas included 1) the original habitat (Southeast) which includes the reference area used in population counts from 2000-2008 as well as part of the areas censused, 2) the area West of the foredune road, and 3) the area to the north of the original reference area. Starting points for 100m transects were randomly selected within the 3 areas, and sampled as in 2000-2008, counting only reproductive individuals within 1 meter of either side of the tape. (To see details of sampling methods in previous years, see Thorpe 2009). The total population size was estimated by multiplying the average number of plants per 100 m x 2 m plot by the total number of possible plot locations.

In 2012-2018, plant measurements including length of longest branch and the number of inflorescences were collected on 3 plants selected at random along each transect. This data can be used to track changes in plant size and reproductive vigor, and is discussed below. To investigate the substrate of the population, the percent cover of shell within a 0.5 meter radius as well as the number and species of associated plants were recorded.

Seeded Transects

In the spring of 2016, multiple 5m transects were established at Overlook North and Siltcoos, and both *Abronia umbellata* ssp. *breviflora* (~1,000) and *Glehnia littoralis* (~500) were planted in a narrow 5-10 cm deep trench with sand compacted over the seeds. These transects were monitored in the fall of 2016 for survivorship.

Plant Measurements

In 2012-2018, additional plant measurements were recorded on selected sand-verbena plants on the North Spit. At Coos Bay North Spit, three plants along each survey transect were randomly selected to be monitored. In addition, at the sites listed in Table 6, 20 points were randomly selected using GIS and uploaded points to handheld GPS units. In the field, the closest flowering plant to the preselected GPS point was monitored. Plant measurements included length of longest branch, number of infructescences on the entire plant, and the number and species of plants within 0.5m. Notes about substrate and habitat were recorded.



FIGURE 3. IN 2009, THE POPULATED AREA WAS DIVIDED INTO THREE SECTION, 1) THE ORIGINAL HABITAT (SOUTHEAST), 2) NORTHEAST AND 3) WEST OF THE FOREDUNE ROAD. EACH AREA WAS SUBSAMPLED USING THE METHODS DESCRIBED IN THE TEXT.

Overwintering Plants

At Overlook North and South and Siltcoos, plant characteristics were measured on selected overwintering plants. At the time of seeding in March 2012, a number of robust plants were found apparently overwintering, and 18 of these plants were marked with wooden stakes, GPS coordinates and spatial notes relative to the marking stake were taken. Six plants at Overlook North, South and Siltcoos were marked (although more overwintering plants were present at all sites). It is not uncommon for a few of the usually annual plants to perennate. In the fall these plants were monitored and the longest branch and number of inflorescences counted in 2013, 2015 and 2016 (if the plant was present).

In the spring of 2016, additional overwintering plants were marked at Overlook North and Overlook South and were monitored for presence/absence as well as plant measurements in the fall of 2016. While overwintering plants have been noted since 2016, resources have not been available for targeted monitoring of these plants.

TABLE 1. REINTRODUCTION SITES FOR THE COOS BAY BLM DISTRICT, METHODS USED, AND HABITATS FOR SEEDING AND TRANSPLANTING EFFORTS DISCUSSED IN THIS REPORT. SEEDINGS REPORTED HERE ALL INVOLVED THE USE OF 50,000 SEEDS PER SITE, EXCEPT WHERE NOTED.

Land Manager	Site	Location	Method	Habitat
<i>Coos Bay District Bureau of Land Management</i>				
	Coos Bay, North Spit	North of Coos Bay	<u>Seeding:</u> Dec. '96, March '06 (100k; newly restored area), March '07 (100k), March '08 (90k), Feb '09 (90k), Feb '10 (100k), Feb '11 (100k), Feb '12 (100k), Mar '13 (100k), Mar '14 (100k), Mar '15 (100k), Mar '16 (100k), Mar '17 (100k), Mar '18 (100k)	Open beach and beachgrass treatment area (sand spit)
	New River ACEC	sand spit near the mouth of Fourmile Creek about 8 miles south of Bandon	<u>Seeding:</u> Jan. '97, March '99, '00, '01 (70k), '02 (100k), March '03 (80k), March '04 (100k), March '05 (95k), March '06 (100k), March '07 (150k), March '08 (80k), Feb. '09, (80k) Feb. '10 (100k), Feb '11 (100k), Feb '12 (120k)*, Mar '13 (120k), Mar '14 (100k), Mar '15 (100k), Mar '16 (100k), Mar '17 (110k), Mar '18 (100k)**)	Foredune beachgrass treatment area and open sand * In 2012, 20k seeds were distributed in an area ~1.5 miles to the south near Butte Creek. It is recommended that additional beach grass removal occur at the site prior to any future seeding efforts. ** In 2018, seeding efforts were focused on the 4 areas to the south of the boat ramp that experience overwash during large winter storms.

TABLE 2. REINTRODUCTION SITES OF THE SIUSLAW NATIONAL FOREST, METHODS USED, AND HABITATS FOR SEEDING AND TRANSPLANTING EFFORTS DISCUSSED IN THIS REPORT. SEEDINGS REPORTED HERE ALL INVOLVED THE USE OF 50,000 SEEDS PER SITE EXCEPT WHERE NOTED.

Land Manager	Site	Location	Method	Habitat
<i>Siuslaw National Forest</i>				
Baker Beach		<i>Siuslaw National Forest</i> Lily Lake, north side of Berry Creek, approximately 7 miles north of Florence	<u>Seeding</u> : Feb. '03, March '04, March '05, March '06, March '07, March '08, March '09, March '18 35k split between Baker and Sutton	Beach, beachgrass treatment area.
Sutton Creek		<i>Siuslaw National Forest</i> 6 miles north of Florence, Lane County	<u>Seeding</u> : March '04, March '06, March '18 35k split between Baker and Sutton	Mouth of creek, foredune beachgrass treatment area and open sand
Overlook (N and S)		Douglas County about 2.5 miles south of the Lane/Douglas County line (in T20S R12W Sec 17 NW)	<u>Transplanting</u> : '00 <u>Seeding</u> : March '01 (N 50k and S 50k), March '02 (S only), March '05 (S only), March '07 (N 80k and S 50k), March '08 (N 100k), March '09 (N 50k and S 50k), March '10 (N 50k and S 50k), March '11 (N 60k and S 60k), March '12 (N 60k and S 60k), March '13 (N 60k and S 60k), March '14 (N 60k and S 60k), March '15 (N 60k and S 60k), March '16 (N 60k and S 60k), March '17 (N 60k and S 60k), March '18 (N 75k and S 90k)	Foredune beachgrass treatment area and open sand
Siltcoos Creek		10 miles north of Reedsport	<u>Transplanting</u> : '97 and '01 <u>Seeding</u> : Feb. '98, March '00, March '02 (60K), March '04, March '05, March '06, March '07 (15k to small beach north of creek and 85k to large plover area), March '08 (60k), March '09 (55k), March '10, March '11, March '12 (12.5k), Mar '13	Foredune beachgrass treatment area and open beach, mouth of creek

(12.5k), Mar '14 (10k), Mar '15 (50k),
March '16 (53k), March '17 (50k),
March '18 (50k)

Tahkenitch Creek

7 miles north of
Reedsport

Transplanting: '97-98, '00-01 Seeding:
Feb. '98, March '01, March '02, March
'04, March '05, March '11 (80k),
March '12 (100k), Mar '13 (100k),
Mar '15 (80k in southern portion- focus in
southern portion), Mar '16 (80k - focus in
southern portion), Mar '17 (80k - focus in
southern portion), Mar '18 (90k - focus in
southern portion)

Foredune beachgrass
treatment area and open
beach, mouth of creek

RESULTS

Population Surveys

General Trend

The population of pink sand-verbena increased at all sites except Coos Bay North Spit and North Overlook. (Totals increased at New River, Tahkenitch, Baker Beach and South Overlook.)

Despite increases observed in 2018, most populations of pink sand-verbena in Oregon remain fewer than 1,000 individuals. Substantial interannual variability in population counts observed over the course of this study are cause for increasing management efforts and continuing monitoring (Figure 7, Figure 10). Two of the largest drivers appear to be amount of disturbance (e.g. the exposed beach at New River and Siltcoos) and competition by European beach-grass. For example, in 2013 at the North Spit, there was a clear difference in the density, size and reproductive effort between plants in areas that had (or had not) received discing (and beach grass removal) (Figure 7). Climate is also likely an important driver. On a local scale, coastal climate may also be influenced by landscape features such as headlands and nearshore rocks, which may result in differences in wave action between sites in the same year.

For a historical summary of inventories for pink sand-verbena conducted by IAE from 2000-2015 on the central and southern Oregon Coast see Giles-Johnson and Kaye 2015.

Siuslaw National Forest

Monitoring activities for pink sand verbena in 2018 occurred on five Forest Service Sites, Siltcoos, Overlook North, Overlook South, Tahkenitch and Baker Beach/Sutton Creek.

SILTCOOS CREEK

Pink sand-verbena restoration and reintroduction was initiated at the mouth of Siltcoos Creek in 1997, when 27 plants were outplanted at the site. The population has fluctuated dramatically over time, from zero in 1999, to a high of 989 (nearly all of which were reproductive) in 2005 (Figure 4, Table 14). In the winter of 2012/2013 a substantial portion of the previously occupied habitat eroded leaving less than 20% of the original habitat. In 2014 only ten plants were noted, however since that time the population has increased to as many as 427 plants (2016, Figure 4). In 2018, 292 plants were observed 147 reproductive. This is also likely an underestimate of the plants at the site, as monitoring occurred later than at other sites when some plants may have already senesced or been covered by shifting sands. Despite the apparent loss of habitat in 2012, Siltcoos has consistently had one of the highest reintroduction success rates, and we recommend continued seed addition efforts and beach-grass removal.

In 2011, 2015, 2016 and 2018 plants at this site were generally larger and had more fruits per plant compared to the other sites monitored (Table 6, Table 7). No plant measurements were recorded at Siltcoos in 2017.




FIGURE 4. POPULATION TRENDS AT SILCOOS CREEK, 1996 – 2018. AN EROSION EVENT IN THE WINTER OF 2012/2013 RESULTED IN SIGNIFICANT LOSS OF HABITABLE HABITAT, THUS THE DECREASE IN SEEDING EFFORT AND THE NUMBER OF PLANTS OBSERVED AT THE SITE.

TAKHENITCH CREEK

In 1996, 34 transplanted pink sand-verbenas and 5,000 seeds were dispersed at Tahkenitch Creek as part of an early effort to establish plants on Oregon beaches. This effort has been followed with two additional transplanting efforts and seeding. Prior to 2005, the area was divided into two sections, North Tahkenitch and South Tahkenitch. The Tahkenitch stream channel has shifted to the south and now most of the habitable area is north of the creek. In 2005 no plants were found, and in 2006, only one vegetative plant was present.

An informal survey in 2009 by Marty Stein, located 4 plants, 2 reproductive and 2 vegetative, suggesting that this site may be suitable for future reintroduction efforts. In 2011, 80,000 seeds were distributed resulting in 95 plants (57 vegetative and 38 reproductive). In 2012, 100,000 seeds were distributed resulting in 381 plants (246 vegetative and 135 reproductive.) Seeding continued in 2013-2018 resulting in 13-361 plants (Figure 5). Despite the relatively small number of plants, we recommend continued seeding in the area combined with beach grass control to build the seed bank at this site.




FIGURE 5. POPULATION TRENDS AND SEEDING HISTORY AT TAHKENTICH CREEK OF PINK SAND FROM 2001-2018. THIS SITE WAS NOT MONITORED IN 2007, 2008 OR 2010.

OVERLOOK NORTH AND SOUTH

European beach-grass control was initiated at the Overlook sites in the fall of 1999 through the use of bulldozers operated by the Oregon National Guard in coordination with the Siuslaw National Forest. Pink sand-verbena reintroduction at the Overlook sites began in spring 2000 with the planting of 50 individuals. In 2001, both the north and south sites received 50,000 seeds which resulted in the establishment of 482 and 658 plants, respectively.

At Overlook North from 2002-2006 no seeding occurred. Seeding resumed in 2007 and has continued annually (Table 14). The population remained relatively stable from 2008-2012. In 2012, North Overlook had 4,860 plants (3,233 vegetative and 1,627 reproductive). This is the second highest number of plants found at the site however the ratio of reproductive to vegetative plants was lower. At South Overlook seeding has occurred annually with the exceptions of 2002, 2003, and 2006. In 2012 South Overlook had 2,247 plants (1,619 vegetative and 928 reproductive). In 2013, as with other sites, the number of plants decreased at both Overlook North and Overlook South. This pattern continued into 2015, where only 124 and 158 plants were noted, respectively. In 2017, numbers slightly increased at both Overlook sites, however remain among the lowest observed (Figure 6, Table 14).

Despite the precipitous drop in the population from 2012-2015, the reintroduced population at the Overlook sites is now the second largest population in Oregon (Coos Bay North Spit is the largest), and has gradually increased in the last two years to more than 1,500 plants (Figure 6, Table 3). Without continual beach-grass control, the substantial population growth at this site will likely reverse and the population will decline. We recommend that seeding continue at both sites, with higher priority given to the Overlook South treatment area with the potential addition of other native dune species (see Recommendations section). We also recommend continued observations of the spread and potential impact of *Leymus mollis* on the other native dune species.




FIGURE 6. POPULATION TRENDS AND SEEDING HISTORY AT OVERLOOK NORTH AND SOUTH OF PINK SAND VERBENA FROM 2000-2018. OVERLOOK SOUTH WAS NOT MONITORED IN 2008.

TABLE 3. NUMBER OF PINK SAND VERBENA AT OVERLOOK SOUTH AND OVERLOOK NORTH 2000-2018. *OVERLOOK SOUTH WAS NOT MONITORED IN 2008.

	Overlook South					Overlook North				
	Vegetative	Reproductive	Total	% reproductive	Amount seeded (k)	Vegetative	Reproductive	Total	% reproductive	Amount seeded (k)
2000	-	-	-	-	-	0	2	2	100%	transplants
2001	231	427	658	65%	50	92	390	482	81%	50
2002	291	1435	1726	83%	50	517	574	1091	53%	
2003	153	1954	2107	93%		119	1359	1478	92%	
2004	3229	6349	9580	66%		1111	2706	3817	71%	
2005	1759	2581	4340	59%	50	2019	1191	3210	37%	
2006	6	4	10	40%		5849	1976	7825	25%	
2007	22	76	98	78%	50	655	2790	3445	81%	80
2008*	Not monitored				50	1584	811	2395	34%	100
2009	555	1518	2073	73%	50	323	466	789	59%	50
2010	628	424	1052	40%	50	492	238	730	33%	50
2011	1361	988	2349	42%	60	1194	744	1938	38%	60
2012	1619	628	2247	28%	60	3233	1627	4860	33%	60
2013	395	87	482	18%	60	783	211	994	21%	
2014	43	66	109	61%	60	132	149	281	53%	
2015	37	121	158	77%	60	53	71	124	57%	60
2016	239	375	614	61%	60	111	158	269	59%	60
2017	348	618	966	64%	60	241	375	616	61%	60
2018**	1626	1850	3476	53%	90	213	284	616	46%	70
Max. value	3229	6349	9580	93%		5849	2790	7825	100%	
Min. value	6	4	10	18%		0	2	2	21%	
Average	738	1147	1885	59%		985	849	1840	55%	
Median	348	618	1052	61%		492	466	994	55%	

Seeded Transects

In the spring of 2016, a total of six transects were planted, each with approximately 500 *Glehnia littoralis* and 1,000 *A. umbellata* ssp. *breviflora* (three transects at Overlook South and three transects at Siltcoos.) In the fall of 2016, there were no plants found along the transects at Overlook, and few vegetative plants were present at Siltcoos. At Overlook (where no plants were found on the transect), the seeded trench was excavated, and no seeds were recovered in the area where seeds had been placed. Shifting sands and winds, seed predation, or decay of seeds in the sand could have been factors contributing to the lack of success of these transects, and the apparent absence of seeds in the fall.

Overwintering Plants

At both North and South Overlook nearly all overwintering plants were still present in the fall of 2012. At Siltcoos, none of the overwintering plants noted in the spring of 2012 were found in the fall of 2012. In that year, the average length of the longest stem on overwintering plants at South Overlook was 96 cm, with an average inflorescence count of 620. This is substantially higher than the average for the rest of the population at South Overlook which had an average longest branch of 24 cm and only 6.3 inflorescences. North Overlook followed a similar pattern for overwintering plants with the average longest branch of 57 cm and 97 inflorescences compared to 44 cm and 25 inflorescences. It was also noted that the overwintering plants generally had more branches per plant than other plants.

In the fall of 2013 measurements were repeated on any overwintering plants that were still present, (none at Siltcoos and three at Overlook.) In 2015, a single plant remained, and was reproductive, from those originally marked in 2012, but was not recorded in 2016. Of the 12 overwintering plants marked at Overlook North in spring 2016, only 3 were still alive during the fall sample. At the time of monitoring in 2017, no previously overwintering plants were noted.

BAKER BEACH AND SUTTON CREEK

In 2018, approximately 35,000 seeds were spread at Baker Beach and Sutton Creek. Previously, the Baker Beach site had been seeded with 50k seeds from 2003-2009 (Table 2). No plants were present at the site in 2018. The nearby Sutton Creek, was seeded in 2004 and 2006 and again in 2018. A total of 28 plants were observed at Sutton Creek, 18 reproductive. Measurements of plant size were not conducted at this site in 2018.

TABLE 4. NUMBER OF PLANTS OBSERVED AT SUTTON CREEK AND BAKER BEACH FROM 2003 TO PRESENT. SITES WERE NOT MONITORED IN EACH YEAR.

Year	Sutton Creek				Baker Beach			
	Veg.	Repro.	Total	% Repro.	Veg.	Repro.	Total	% Repro.
2003	Not monitored				28	27	55	49%
2004	122	28	150	19%	56	37	93	40%
2005	0	0	0		61	11	72	15%
2006	0	0	0		6	6	12	50%
2007	0	0	0		12	30	42	71%
2008	Not monitored				38	32	70	46%
2009	Not monitored				5	1	6	17%
2010	Not monitored				Not monitored			
2011	Not monitored				Not monitored			
2012	Not monitored				Not monitored			
2013	Not monitored				Not monitored			
2014	Not monitored				Not monitored			
2015	Not monitored				Not monitored			
2016	Not monitored				Not monitored			
2017	Not monitored				Not monitored			
2018	10	18	28	64%	0	0	0	

Bureau of Land Management

COOS BAY NORTH SPIT

Efforts to eliminate European beachgrass from a roughly 170 acre Habitat Restoration Area (HRA) on the Coos Bay North Spit (Figure 7) have included machine ripping and discing (with a tractor), herbicide (Rodeo), burning, saltwater irrigation, hand pulling, and various combinations of these techniques. Ten thousand pink sand-verbena seeds were sown in each of five 30 x 30 m plots randomly placed in areas with different beachgrass management histories on January 15, 1997 (50,000 seeds total). In the winter of 2005-2006 an area closer to the beach on the west side of the access road was cleared of European beach grass. One hundred thousand seeds were added to this area in both 2006 and 2007, 90,000 seeds were added in 2008 and 2009, and 100,000 seeds were added in 2010-2012. All habitat at the North Spit was disced in October (2008) or November (2009-2018).

Population surveys

Following re-introduction in 1997, pink sand-verbena populations grew steadily to an estimated peak in 2012 of 349,423 (95% CI range: 196,366-502,480), and then 470,092 (296,255-643,929) in 2014 (Figure 7, Table 13). In 2009, we modified our monitoring technique so that the entire population was subsampled. The increase in population size from 2007 to 2009 likely is a result of both population growth and improved monitoring techniques initiated in 2009. Despite the large increase in population size from 2011 to 2012, and again in 2014, the size of reproductive plants seems to be diminishing as does seed yield (Table 6, Table 7, Figure 7, Figure 9). Additionally, there was a significant decline in the size of the population from 2012-2013 that coincided with a year of no discing treatments in the

southern portion of the occupied habitat. In 2018, the population declined to an estimated 37,084 plants (95% CI range: 14,012-60,156) which is the lowest observed since 2005.

In 2012-2018, plant measurements were taken on 3 randomly selected plants along each transect (Table 6, Table 7). These data allow us to track changes in plant size and reproductive vigor at the Coos Bay North Spit, and compare the size of the plants to other occupied sites (Table 6, Table 7). In addition to plant measurements, the number and species of plants within 0.5m were recorded as well as the percent cover of oyster shell.

In 2018 plant density in the Northeast and Southeast portions of the habitat decreased (). Density of plants in the north habitat was the highest at 0.11 plants/m² (down from 2.0 in 2017) and lowest on the west side with only 0.05 plants/m² (Figure 8, Figure 9). In the southern portion of the habitat, density decreased from 0.2 plants/m² to 0.03 plants/m² from 2017 to 2018. Only densities in 2013 have been lower for this portion of the habitat. In 2013, the density of plants decreased in the southern portion of the habitat that did not receive discing treatments, which highlights the tenuous hold that the species has and the importance of continued management. In 2014, after management resumed in the southern portion of the habitat, plant density returned to levels observed in previous years (when management occurred). The cause of the declines in density and estimated population size at the North Spit in 2018 is not clear.




FIGURE 7. POPULATION TRENDS AT THE COOS BAY NORTH SPIT REINTRODUCTION SITE. THE AREA WAS SEEDED IN JANUARY 1997 WITH 50,000 SEEDS FROM PORT ORFORD. POPULATION SIZE ESTIMATES FOR 1997-1999 ARE COMPLETE CENSUSES OF ALL PLANTS IN THE POPULATION, WHILE THOSE FROM 2000-2008 ARE ESTIMATED FROM A SUBSAMPLE OF A REFERENCE AREA AND CENSUS OF THE REMAINDER OF THE POPULATION. THE ERROR BARS FOR 2000-2008 ESTIMATES REPRESENT 95% CONFIDENCE INTERVALS. *IN 2008, AN ESTIMATE OF THE ENTIRE POPULATION SIZE WAS NOT POSSIBLE BECAUSE OVER 50% OF THE POPULATION HAD BEEN DISCED PRIOR TO OUR SURVEYS. SEE TEXT FOR A DISCUSSION OF POPULATION TRENDS. IN 2009, THE SAMPLING METHODS WERE MODIFIED SO THAT THE ENTIRE POPULATION WAS SUB-SAMPLED (AND NO AREAS CENSUSED). IN THIS CHART THE CI

FOR 2009-2019 ARE NOT REPORTED BECAUSE THE POPULATION ESTIMATE IS A COMBINATION OF 3 DIFFERENT POPULATIONS, EACH WITH THEIR OWN CI.



FIGURE 8. ESTIMATED NUMBER OF PLANTS AT THE COOS BAY NORTH SPIT IN EACH OF THE THREE POPULATION AREAS. NOTE THAT BECAUSE THE POPULATION IS SMALL ON THE WEST SIDE, THE SCALE DIFFERS THAN THAT FOR THE NORTH AND SOUTH. ERROR BARS REPRESENT 95% CONFIDENCE INTERVALS.

The Coos Bay North Spit is now the largest pink sand-verbena population in Oregon and serves as the primary seed source for reintroduction efforts in Oregon. Successful plant establishment and population growth has followed successful control measures for European beach-grass on the spit (described in Pickart and Sawyer 1998 and Kaye 1998). No beach-grass control was conducted in 2004 and 2005, and the pink sand-verbena population declined in both of those years. Although the seedbank appears large enough for continual recruitment, the success of this site appears dependent on sustained management activities, particularly annual discing to control plant invasion and additional vegetation control north of the HRA. This pattern was again highlighted in 2013, when the density of pink sand verbena plants decreased from 1.5 to 0.02 /m² in just one year without discing treatment (Figure 9).

However, we have also learned that second year plants can become substantially larger than first year plants if left undisturbed. In 2013, in the southern area where discing did not occur, the few remaining plants were larger than in previous years (Table 6, Table 7). Therefore, in areas with no European beachgrass and abundant pink sand verbena, we recommend that discing be postponed to allow for some individual plants to reach larger sizes and produce larger amounts of seed (Table 6, Table 7). This could be accomplished by lifting the disc on the tractor to skip areas with no beach grass.




FIGURE 9. DENSITY OF PLANTS PER M² AT THE COOS BAY NORTH SPIT FROM 2009-2018. THE DECREASE IN ALL AREAS FROM 2012-2013 COINCIDED WITH LACK OF HABITAT MANAGEMENT. DENSITY MEASUREMENTS FOR THE WEST SIDE ARE NOT AVAILABE FOR 2017. ERROR BARS RESPRESENT 95% CONFIDENCE INTERVALS.

TABLE 5. ESTIMATED NUMBER OF PINK SAND VERBENA AT THE COOS BAY NORTH SPIT FROM 2009-2018.

	Area	Population Estimate	95% Confidence Interval	
			Lower	Upper
2009	North	78,080	26,008	130,152
	South	113,613	73,082	154,144
	West	9,684	4,922	14,446
	Total	201,377	104,011	298,743
2010	North	79,437	47,327	111,547
	South	154,502	71,459	237,545
	West	6,549	2,137	10,961
	Total	240,488	120,923	360,053
2011	North	104,030	74,775	133,285
	South	71,371	35,006	107,736
	West	5,376	2,723	8,029
	Total	180,777	112,504	249,050
2012	North	198,063	113,862	282,264
	South	139,614	79,309	199,919
	West	11,746	3,195	20,297
	Total	349,423	196,366	502,480
2013	North	64,840	44,570	85,110
	South	5,301	3,248	7,354
	West	4,663	1,247	8,079
	Total	74,804	49,065	100,543
2014	North	276,349	189,456	363,242
	South	188,345	104,210	272,480
	West	5,398	2,589	8,207
	Total	470,092	296,255	643,929
2015	North	169,537	68,290	270,784
	South	91,960	53,146	130,774
	West	3,260	-125	6,645
	Total	264,757	121,311	408,203
2016	North	191,512	113,787	269,237
	South	107,184	59,158	155,210
	West	7,480	2,495	12,465
	Total	306,176	175,440	436,912
2017	North	139,340	77,650	201,030
	South	55,192	20,905	89,479
	West	5000*	*	*
	Total	194,532	98,555	290,509
2018	North	14,806	10,533	19,079
	South	10,720	2,669	18,771
	West	11,558	810	22,306
	Total	37,084	14,012	60,156

*Values for 2017 on the west side are estimates.

TABLE 6. AVERAGE SIZE OF PLANTS IN 2012-2018 AT SELECT SITES. “-“ INDICATES DATA WAS NOT COLLECTED IN THAT YEAR. VALUES IN () REPRESENT 95% CONFIDENCE INTERVALS. SYSTEMATIC MEASUREMENTS ON OVERWINTERING PLANTS WERE ONLY CONDUCTED IN 2012.

Site		Average of Longest stem (cm)						
		2012	2013	2014	2015	2016	2017	2018
Tahkenitch		-	-	-	34.2 (10.2)	-	-	54.2 (14.2)
Siltcoos		-	-	-	66.1 (15.2)	90 (29.6)	-	72.2 (13.4)
Coos Bay North Spit	North	18.6 (5.2)	20.7 (4.8)	16 (4.4)	16.4 (8.2)	14.7 (5.0)	15.3 (3.5)	20.1 (5.5)
	South	14.6 (2.3)	33.1 (12.7)	16.7 (3.0)	13.2 (3.1)	17 (5.1)	11.1 (1.8)	17.2 (3.7)
	West	31.7 (7.1)	29.2 (8.1)	24.2 (5.5)	28 8.4)	22.6 (4.9)	9.9 (5.6)	31.1 (6.7)
New River		18.8 (6.4)	24.5 (9.2)	-	34.5 (10.4)	68.0 (16.9)	-	39.2 (10.3)
Overlook Overwintering plants	North	56.8 (35.5)	-	-	-	-	-	-
	South	96.5 (20.8)	-	-	-	-	-	-
Overlook	North	44.2 (21.8)	24.2 (9.1)	-	24.9 (7.4)	37.2 (9.8)	-	30.0 (6.9)
	South	23.1 (4.6)	33.9 (13.7)	-	49.5 (13.6)	50.3 (14.5)	-	68.5 (38.4)

TABLE 7. AVERAGE NUMBER OF FLOWERS IN 2012-2018 AT SELECT SITES. “-“ INDICATES DATA WAS NOT COLLECTED IN THAT YEAR. VALUES IN () REPRESENT 95% CONFIDENCE INTERVALS. SYSTEMATIC MEASUREMENT ON OVERWINTERING PLANTS WERE NOT CONDUCTED IN EVERY YEAR.

Site	Average of # Flowers							
	2012	2013	2014	2015	2016	2017	2018	
Tahkenitch	-	-	-	6.6 (3.5)	-	-	50.3 (29.1)	
Siltcoos	-	-	-	46.3 (20.2)	23.2 (8.8)	-	71.0 (33.9)	
Coos Bay North Spit	North	8.6 (2.5)	7.8 (2.6)	13.1 (5.8)	-	9.5 (5.4)	6.6 (1.7)	25.7 (23.5)
	South	8.7 (1.4)	11.5 (4.9)	14.5 (4.2)	6.6 (2.7)	11 (7.4)	4.3 (1.1)	8.5 (4.8)
	West	11.7	12.6	9.7	-	14.2 (4.9)	7.3 (11.9)	18.6 (7.3)
New River	14.7 (9.1)	6.5 (5.5)	-	14.1 (9.8)	70.4 (25.7)	-	135.1 (44.8)	
Overlook Overwintering plants	North	97.5 (86.4)	-	-	-	-	-	
	South	620.2 (515.4)	-	-	-	-	-	
Overlook	North	24.6 (24.6)	7.2 (4.5)	-	9.3 (4.3)	26.5 (10.6)	--	11.7 (7.0)
	South	6.3 (2.9)	14.6 (11.0)	-	33.7 (20.6)	47.5 (26.2)	-	68.5 (38.3)

NEW RIVER ACEC

At New River, 50,000-150,000 seeds were dispersed within swales through the foredune almost every year from 1997 to 2017 (Figure 10). The spit at New River has been breached in various places by storm waves, creating swales that have low vegetative cover but are bordered on two sides by established plants, primarily European beachgrass. From 1998 to 2005, the BLM made aggressive attempts each fall to reduce the abundance of European beachgrass and lower the foredune using heavy machinery. There was no disturbance in winter 2006 – 2008 and European beachgrass rapidly reinvaded the dunes. Discing resumed winter 2009. Although beachgrass remained throughout the site, the habitat was much more open in 2010 compared to previous years. Discing occurred on a portion of the area in 2010, and was repeated at least bi-annually through 2017, reducing the cover of beach grass in the disced area. Population surveys occurred in September or October of each year since 1996. Immediately prior to the 1999 site visit, much of the area that had been seeded with pink sand-verbena was worked by a bull dozer to remove European beachgrass.

Population surveys

The pink sand verbena population at New River has been sustained by direct seeding with 50,000 - 150,000 seeds from 1997-2018 (Figure 10). In the first year of seeding (1997), 118 plants established. Since that time, the population has ranged from 0- 2,172 plants (in 1998 and 2005 respectively) (Figure 10). We recommend that seeding and beachgrass removal efforts continue at this site using the current approach of seeding following disturbance in the fall. In 2012 -17 most plants at the site were found in areas that had been overwashed in winter/spring storms as well as on the shore side of the foredune, highlighting the species adaptation to disturbed habitats. In 2018, seeding efforts were focused in overwashes at the New River ACEC.

In the fall of 2015, a large proportion of the plants (1,630 of the 1,824 plants observed) were found in an area that had received extensive overwash and temporary pool formation- as evidenced by the presence of fine sediment materials (Figure 11). Increased nutrient and water availability to the disturbed seed bank likely contributed to the high numbers of plants in the area, which also included large amounts of *Rumex acetosella* (as well as the native *Rumex maritimus*) (Figure 11). Although total numbers were up for the site in 2015, the number of reproductive plants remained relatively low with only 287 flowering (18%); only 198 plants were found in the remaining portion of the population in 2015. In 2017 only 9 plants were found, and as in previous years, the few plants were found in and along the edges of overwashes.. In 2018, when seeding efforts were focused in overwashes at the New River ACEC, a total of plants 196 plants were observed in three of the five overwashes seeded in 2018 (72 vegetative, 126 reproductive) (Figure 10).




FIGURE 10. POPULATION TRENDS AND SEEDING HISTORY OF PINK SAND VERBENA AT THE NEW RIVER ACEC REINTRODUCTION SITE. SEEDING HAS BEEN CONDUCTED IN ALL YEARS EXCEPT 1998.



FIGURE 11. NEARLY ALL PLANTS IN 2015 AT NEW RIVER WERE FOUND AGAINST THE BACK OF THE DUNE IN AN AREA THAT HAD BEEN OVERWASHED. THE AREA CIRCLED IN RED HAD HUNDREDS OF SMALL PINK SAND VERBENA PLANTS, AS COMPARED TO THE SINGLE, LARGE REPRODUCTIVE PLANT IN THE FOREGROUND.

Plant Measurements

Plant Size and Reproductive Effort

Since 2012, plant size has been measured on randomly selected reproductive plants at selected sites (Table 6, Table 7). The average length of the longest branch and the average number of inflorescences per plant for sites monitored in 2012 is listed in Table 6 and Table 7. In 2012, measurements were also conducted on plants that had overwintered. These plants were larger with more flowers than other plants at the same sites that had not overwintered (Table 6, Table 7).

In 2018, the largest plants, (as measured by the length of the longest stem), were found at Siltcoos, New River and Overlook South (Table 6). Plants at the Coos Bay North Spit have been consistently smaller than plants measured at other sites with fewer flowers (Table 6). In 2018, only Overlook North had plants as small or smaller as those observed at the North Spit (Coos Bay range: 8.5 ± 4.8 - 25.7 ± 23.5 flowers, 17.2 ± 3.7 - 31.1 ± 6.7 cm, and Overlook South: 11.7 ± 7.0 flowers, 30.0 ± 6.9 cm). Plants at Coos Bay North Spit are generally larger on the west side (where density is also much lower), than in the other portions of the habitat. Plant size and number of flowers has shown a general decrease since 2012 at the North Spit, with the smallest average plants and least number of flowers observed in 2017. Plant size at the North Spit increased slightly in 2018; despite increases in plant size from 2017, plants at this site continue to be smaller and with fewer flowers than plants at other sites (Table 6, Table 7).

Plant Density

In addition to measurements of plant size and reproductive effort, plant density was also estimated for all monitored populations in 2012, and at the Coos Bay North Spit annually since that date. The estimated density of plants/m² at the sites monitored in 2012 was 0.001-1.5 plants/m². Coos Bay North Spit had the highest density of plants overall with a high of 1.5 plants/m² in the northern portion of the population. In contrast, Overlook North (what we consider a fairly successful site) had only 0.02 plants/m². In 2013, plant density was lower at all sites, but the general pattern of density remained the same with Coos Bay North Spit the most dense portion of the population, and other populated sites with densities as low as 0.001 plants/m². This general pattern for density was the same in 2014-2018 (Figure 9, Table 6, Table 7).

Plant Community

The presence(s) and count of plants within a 0.5 m radius of each measured sand-verbena plant was also noted at the North Spit from 2012-2018, and at most other sites in 2012-2018. There was a total of 19 different plant species found growing adjacent to sand-verbena, these species are listed Table 8. The most common plant species found within a 0.5 m radius of pink sand-verbena were *Cakile* sp. (sea rocket) followed by *Ammophila arenaria*, (European beachgrass), *Abronia umbellata* (pink sand-verbena) and *Camissonia cheiranthifolia* (Beach suncup) (Table 8). *Leymus mollis* (American beachgrass) was also common at both Overlook sites as well as in the western portion at Coos Bay North Spit.

TABLE 8. COUNT OF PLANT SPECIES WITHIN 0.5M OF RANDOMLY SELECTED PINK SAND-VERBENA PLANTS, INCLUDING PERCENT OF PINK SAND-VERBENA MONITORED THAT HAD SPECIES PRESENT, IN 2012-2018 AT THE COOS BAY NORTH SPIT. NAMES IN BOLD INDICATE NATIVE SPECIES. *BECAUSE IT IS DIFFICULT TO DISTINGUISH BETWEEN INDIVIDUALS OF BOTH AMERICAN AND EUROPEAN BEACHGRASS, PRESENCE/ABSENCE ONLY WAS NOTED.

Year	Count (and percent) of times species observed						
	2012	2013	2014	2015	2016	2017	2018
Number of ABUM monitored	178	60	158	83	115	107	109
Species							
<i>Ammophila arenaria*</i>	112 (63%)	31 (52%)	81 (51%)	42 (51%)	51 (44%)	47 (41%)	63 (58%)
<i>Leymus mollis*</i>	14 (8%)	13 (22%)	23 (15%)	8 (10%)	23 (20%)	1 (1%)	18 (17%)
<i>Abronia umbellata</i> ssp. <i>breviflora**</i>	124 (70%)	37 (62%)	88 (56%)	68 (82%)	74 (64%)	47 (44%)	41 (38%)
<i>Achillea millefolium</i>	0 (0%)	0 (0%)	1 (1%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
<i>Anaphalis margaritacea</i>	3 (2%)	0 (0%)	2 (1%)	1 (1%)	0 (0%)	0 (0%)	0 (0%)
<i>Cakile edentula/maritima</i>	132 (74%)	29 (48%)	97 (61%)	58 (70%)	54 (47%)	34 (32%)	60 (55%)
<i>Camissonia cheiranthifolia</i>	66 (37%)	33 (55%)	25 (16%)	26 (31%)	38 (33%)	20 (19%)	9 (8%)
<i>Convolvulus soldanella</i>	1 (1%)	0 (0%)	1 (1%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
<i>Fragaria chiloensis</i>	1 (1%)	2 (3%)	0 (0%)	0 (0%)	6 (5%)	0 (0%)	0 (0%)
<i>Glehnia littoralis</i>	6 (3%)	2 (3%)	14 (9%)	19 (23%)	7 (6%)	16 (15%)	19 (17%)
<i>Hypochaeris radicata</i>	45 (25%)	21 (35%)	41 (26%)	24 (29%)	24 (21%)	16 (15%)	11 (10%)
<i>Juncus lesueurii</i>	4 (2%)	1 (2%)	2 (1%)	0 (0%)	2 (2%)	0 (0%)	0 (0%)
<i>Lathyrus japonicus</i>	5 (3%)	3 (5%)	5 (3%)	1 (1%)	1 (1%)	0 (0%)	0 (0%)
<i>Lathyrus littoralis</i>	1 (1%)	0 (0%)	0 (0%)	2 (2%)	0 (0%)	0 (0%)	0 (0%)
<i>Lupinus littoralis</i>	22 (12%)	18 (30%)	5 (3%)	8 (10%)	13 (11%)	23 (21%)	18 (17%)
<i>Rumex acetosella</i>	9 (5%)	10 (17%)	9 (6%)	9 (11%)	11 (10%)	18 (17%)	2 (2%)
Mean % cover shell	2.6	2	3.2	2.1	1.4	5.1	2
Mean % cover wood	-	0.3	0.3	1	0.4	0.1	0.2

** Counts of pink sand verbena are in addition to the plant being monitored.

DISCUSSION

Population trends, plant size and habitat management

We have observed that seeding is often successful in establishing a pink sand-verbena population the year of seed addition, but that these populations may not persist in subsequent years without either seed additions and/or beachgrass control. Repeated seeding efforts to build up a local seedbank (see seedbank discussion) may be necessary. At four reintroduction sites that have received multiple seeding or transplants attempts, medium to relatively large numbers of plants have been observed despite no seeding in those years at those sites (Siltcoos Creek 2003, Bastendorff Beach 2003, Floras Lake 2004 - 2006, Tahkenitch 2000 – 2004, 2009 and Bandon Beach 2003).

In 2018, at Overlook South, where beachgrass control had occurred and the potential habitat expanded in the previous fall/winter, large numbers of plants were found in the recently cleared habitat. Habitat management actions included the use of herbicide as well as discing of large areas of European beachgrass. Plants in this area were vigorous, with many flowers, while plants at the nearby Overlook North, were smaller with fewer flowers (Figure 13, Table 6, Table 7). This is consistent with observations at other sites including New River ACEC, where in the first few years after initial beachgrass removal, more plants were observed than in subsequent years despite seeding annually (with similar, or sometimes larger amounts of plant materials). Similarly, it is not uncommon to find plants in overwashes, breaches, and in areas that have received natural disturbance in the preceding winter season. Seeding at New River ACEC in 2018 was targeted in overwashes, and more plants were found at this site than in previous years when seeding efforts were focused in the beachgrass management area, which includes large portions of the beach which have not recently been breached.



FIGURE 12. IN 2018, PLANTS WERE LARGER WITH MORE FLOWERS AT OVERLOOK SOUTH (LEFT) WHERE NEW AREAS HAD BEEN TREATED FOR BEACHGRASS REMOVAL THAN AT THE NEARBY OVERLOOK NORTH SITE (RIGHT).

Since the initiation of the project on the Coos Bay North Spit, there has been an increase in the number of plants in the original habitat and in the northern area and an expansion of the habitat to include an area west of the foredune road, however, the size of the plants has decreased since initiation of this study, and in 2018, the estimated population size also decreased. This could be due to a number of factors including; annual variability in the populations due to climactic factors, intra-specific competition,

competition with other annuals including *Cakile edentula*, *Cakile maritima* and *Hypochaeris radicata*, a decrease in the availability of nutrients, the timing and intensity of discing and plowing, a change in the pH caused by the oyster shells and the possible (if unlikely) presence of a contaminant on the oyster shells brought in for the plover habitat.

The populations at Coos Bay North Spit and the Overlook sites are examples of reintroductions into beachgrass control areas that have done reasonably well. In 1996, 50,000 seeds were distributed at the Coos Bay North Spit. The population quickly grew from over 1700 plants in 1997, to 4111 in 1998 and at least 50,000 in 1999. Since 2000, the plants were so abundant that a reference area was subsampled with randomly placed plots and the abundance of the species in the rest of the spit was counted. Since vegetative plants do not contribute to the population (pink sand-verbena is usually an annual) and counting all plants would be infeasible because of time constraints, we counted only reproductive individuals at this site since 2000. It is important to note that in 2004 and 2005 and again in 2013 and 2018, the Coos Bay North Spit population declined significantly, which may be due to the lack of beachgrass management during the prior fall and winter, or other unknown factors. Without repeated discing to reduce invading vegetation such as European beachgrass, pink sand-verbena may fail to increase in numbers and expand its area at this site.

The success of seeding may be related to habitat quality and susceptibility to the scouring effects of winter storms. For example, the presence of competing vegetation, especially in foredune habitats, may result in small plants that produce few or no seeds. Also, winter storms on the Oregon coast can have dramatic effects on beaches, removing large amounts of sand eroding foredunes, and causing shifts in the location of stream channels carrying sediment and nutrients to the sea. The habitats at the Coos Bay North Spit and the Overlook sites are protected from these winter storms, and it is possible that the lack of salt spray and the relatively stable nature of these dunes is causing shifts in the soil/sand chemistry at the site.

At the North Spit, the population is located in the interior of the sand spit and at the Overlook site, populations are on the elevated foredune. This protection allows all seeds produced from the pink sand-verbena plants that grow there to remain on site, maximizing their chances of establishing seedlings the following year. Disturbance at these sites included discing and manual removal of beachgrass and other species in an attempt to keep competing plant species from revegetating the sites. Initial plant establishment at the Coos Bay North Spit in 1997 was highly negatively correlated with the local abundance of European beachgrass (Kaye 1998). Competition studies at Port Orford and Gold Beach on dredge material also suggest that competing vegetation is a major factor affecting establishment and survival of pink sand-verbena (Kaye 1999).

Transplanting

Previous tests of transplant success, and factors effecting survivorship of transplants are discussed in Kaye 1995 and Kaye 1996. Due to the efforts involved in propagation of plant materials for transplanting, and the short life-cycle of this species, further transplanting efforts have not been implemented, with a focus on seeding in order to maximize resources allocated to this species.

ADDITIONAL RESEARCH AND ACTIONS

Population Assessment

At this time, the current status of natural (and some reintroduced) populations of pink sand verbena in Oregon is not known. A comprehensive survey for the species has not been conducted since prior to the publication of the 2006 Conservation Strategy. A comprehensive survey should be conducted to identify the current extent and status of existing populations and to identify potential areas for reintroductions, augmentations, and habitat management actions.

Importance of a seedbank

Long-term persistence of pink sand-verbena at sites on the Oregon coast may depend on the development and maintenance of a long-lived persistent seedbank. Results from germination tests with seeds of various ages (stored in paper sacks at room temperature) show that seeds of this species can remain viable for long periods of time. Loss of viability over time appears to be very slow; seeds stored for nine years retained over 80% viability (). Data collected in 2012-2016 indicates that overwintering plants are both significantly larger and produce more seed than annual plants and may play an important role in developing a natural seedbank (Table 6, Table 7).

The stochastic behavior of natural and reintroduced pink sand-verbena populations suggests that seedbanks buffer populations from sharp declines and can allow re-colonization of a site. At Otter Point, a population decline to zero plants in 2000 was followed by the emergence of three plants in 2001; seven plants were counted at Cape Blanco in 2004 although none had been observed there in 2003. Seeding in 2002 at three reintroduction sites, Siltcoos Creek, Bastendorff Beach, and Bandon Beach, resulted in low plant establishment (fewer than 20 plants each) that year, but much higher populations (>100 plants) in 2003 without additional seeding. The appearance of some populations of pink sand-verbena long distances from natural populations, such as at Tenmile in 1995, Gearhart in 1993, and Vancouver Island in 2000 (after nearly 60 years of absence), suggest that some populations may establish from seed many years after the presence of any adult plants. Two new sightings (Tish-A-Tang and Arizona Beach) in 2012, and one new observation in 2018 at Humbug Creek indicate that seeds may establish from the seedbank. Taken together, these lines of evidence provide strong support for the notion that pink sand-verbena is capable of maintaining a persistent seedbank, and that buried seeds may play an important role in the population dynamics of this species and should be fostered in reintroduction attempts.

Population modeling to assess recovery objectives

The Conservation Strategy for pink sand-verbena (Kaye 2006) calls for the development of recovery objectives that take into account our ability to successfully reintroduce populations of the species. However, we have no estimate of the number of populations needed to establish a successful pink sand-verbena meta-population. Data gathered from population monitoring on natural population dynamics, population establishment success, and estimations of dispersal rates can be used to simulate population dynamics over time. Through the use of matrix models, the viability of individual populations may be estimated based on observations of individuals in populations through time. Pink sand-verbena may be a suitable 'model system' for structuring reintroduction programs and setting objectives, especially for rare beach plant species.

Restoration with additional beach species

Pink sand-verbena is only one of several beach species in decline on the Oregon coast. Efforts to control European beachgrass to improve habitat conditions for native species such as pink sand-verbena and the western snowy plover create large, open areas devoid of vegetation (if successful), similar to the type of habitats that were plentiful prior to beachgrass introduction. These areas would also likely provide good habitat for other native beach and dune plant species. Unfortunately, propagation methods for many beach species have not been developed. A preliminary list of 11 native plant species that could be useful in restoration projects of beach ecosystems is provided in . Additionally, the identification of pollinator species, as well as fauna that utilize the pink sand verbena for food or shelter could inform habitat management and restoration activities in dune habitats.

TABLE 9. PLANT SPECIES NATIVE TO OREGON BEACHES AND DUNES THAT ARE CANDIDATES FOR USE IN ECOSYSTEM RESTORATION PROJECTS. MORE INFORMATION IS NEEDED ON TECHNIQUES FOR GERMINATION AND PROPAGATION OF THESE TAXA.

Common name	Latin name
yellow sand-verbena	<i>Abronia latifolia</i>
beach pea	<i>Lathyrus littoralis</i>
maritime pea	<i>Lathyrus japonicus</i>
beach morning-glory	<i>Convolvulus soldanella</i>
silver bursage	<i>Ambrosia chamissonis</i>
American beach-cup	<i>Camissonia cheiranthifolia</i>
black knotweed	<i>Polygonum paronychia</i>
American dunegrass	<i>Elymus mollis</i>
American glehnia	<i>Glehnia littoralis</i> var. <i>leiocarpa</i>
silvery phacelia*	<i>Phacelia argentea</i>
Wolf's evening primrose*	<i>Oenothera wolfii</i>
seaside dock	<i>Rumex maritimus</i>

*Listed as Threatened with the Oregon Dept. of Agriculture; Species of Concern with the U.S. Fish and Wildlife Service.

Hypotheses for Future Research

The results of attempts to reintroduce pink sand-verbena to beach and dune habitats in Oregon, combined with observations made during field visits, have led to several hypotheses that require further testing before they can be used to guide additional reintroduction efforts.

Recently deposited or disturbed sand is a more suitable substrate for pink sand-verbena growth than older substrates.

1. Is soil formation having an impact on size or reproductive effort of pink sand verbena, and the surrounding plant community?
2. What are the relationships between pink sand verbena and sand pH?
3. What are relationships with salt spray?
4. What are management actions that encourage the number and size of pink sand verbena?

Natural and transplanted plants at Port Orford and Gold Beach in the 1990's consistently thrived on freshly deposited dredge material, but tended to decline in vigor two to three years after the sand was deposited. In 2018 at Overlook South in the newly expanded habitat, plants were larger with more flowers than in the nearby Overlook North which has been treated for beachgrass control since 2004. Furthermore, natural populations in southern Oregon and northern California often occur near the mouths of rivers and creeks where fresh sand is consistently deposited and disturbance is frequent.

A persistent seedbank is a necessary component of a viable population; repeated seedings or transplantings may be required for successful reintroduction and recovery.

1. How long do seeds remain viable when buried?
 - a. Does seed lose dormancy when stored?
2. What is the viability of seed collected in each year?
3. Does the viability of seed vary by site?

Pink sand-verbena seeds are long-lived and appear to be capable of persistence in beach sands for long periods of time (up to several decades). Therefore, buried seeds may play an important role in viable populations of pink sand-verbena, acting as a mechanism for population re-establishment after catastrophes or stochastic swings in population size; similar to what was observed at Siltcoos after major shifts in sand during the 2010/2011 winter season. They may also serve to provide some genetic stability to populations by containing samples of genetic variability that are occasionally lost from the above-ground population. Successful reintroduction of this species may require the development of a large pool of buried seeds, so that introduced populations can rebound after periods of population decline. This suggests that at each site, repeated seedings or transplantings over several years may be required to build-up a seedbank.




FIGURE 13. THE PERCENT OF SEEDS STORED FROM 0-9 YEARS (FROM 1991-2000). THE EQUATION FOR THE LINEAR REGRESSION IS $Y = -0.03X + 0.955$, $R^2 = 0.81$, $P = 0.015$.

Overwintering plants play an important role in the creation of a self-sustaining population. Large overwintering plants are larger and produce significantly more seed than annual plants.

1. How do over-wintering plants differ in size and reproductive effort from annual plants?
2. What percentage of plants overwinter?
3. Are there management actions that can be taken to encourage plants to over-winter?

Mechanical methods of discing and bulldozing commonly used for invasive beach grass control at many of the sand-verbena sites may need to be balanced with the need to allow at least some plants to overwinter and grow for a second year. Manual removal (where feasible), targeted herbicide use, or staggered discing treatments are possible ways to allow for both the control of beach grass and for some sand verbena plants to perennate. Especially in habitat patches where beachgrass is not established in any given year, lifting the disc equipment temporarily may allow more second year plants to flourish.

RECOMMENDATIONS FOR 2019

Seeding and Habitat Management

In 2018 seeds were collected from Coos Bay by BLM staff, for distribution in 2019.

Based on the success of seeding and transplant efforts in previous years, we recommend that the following sites (Table 11) be the focus of seeding and restoration efforts in 2019.

TABLE 10. CURRENT AND SUGGESTED SEEDING RATES FOR PINK SAND-VERBENA SITES.

Management Agency	Site	Seeding Conducted		Notes
		in 2018	Recommended Action in 2019	
US Forest Service, Siuslaw N.F.	Siltcoos Creek	50,000	50,000	
	Sutton Creek/Baker Beach	35,000	50,000	Prior to 2018, site has not been seeded since 2009
	Overlook, North	75,000	60,000	
	Overlook, South	90,000	120,000	Focus seeding in newly treated habitat
	Tahkenitch	90,000	100,000	
Bureau of Land Management, Coos Bay	New River	110,000	120,000	Continue to focus seeding in overwashes across the entire ACEC

Coos Bay North Spit

Continued monitoring and seed collection for redistribution at other restoration sites.

The small number of plants at nearly all monitored populations highlights the need for continued management efforts, and identification of areas where augmentations or reintroductions could be appropriate is recommended. In addition to mechanical and chemical control of European beach grass, continued management at occupied sites could also include seeding of other associated dune species, and efforts to re-establish natural dune conditions (sand deposition, salt spray/overwash, etc.).

Population surveys

With the exception of the sites monitored by IAE in the recent past, the status of this species along the rest of the Oregon coast is unknown. It is recommended that population surveys across Oregon be conducted in the near future, as recommended in the 2006 Conservation Strategy in order to accurately assess the status of this rare species. The sites listed in Table 12 have had plants observed in the last 10 years; it is possible that other isolated populations exist up and down the Oregon Coast. Survey efforts should also include identification of potential habitat, and areas which could benefit from seeding or habitat restoration activities.

TABLE 11. SITES WITH PINK SAND-VERBENA DOCUMENTED IN THE PAST 10 YEARS, WITH SITES MONITORED IN 2018 LISTED IN BOLD.

County (Listed North – South)	Site
Lincoln County	Salmon River
Lane County	Baker Beach Siltcoos Creek Sutton Creek
Douglas County	Overlook North and South Tahkenitch
Coos County	Bandon Beach China Creek Bandon Beach South Coos Bay North Spit
Curry County	Cape Blanco Elk River Euchre Creek (Ophir) Floras Lake New River Otter Point Port Orford Pistol River

LITERATURE CITED

- Gabrielson, I.N. and S.G. Jewett. 1970. *Birds of the Pacific Northwest with special reference to Oregon*. Dover Publishing Inc., New York.
- Giles, D.E.L. and Kaye, T.N. 2015. *Abronia umbellata* subsp. *breviflora* on the Oregon coast: Reintroduction and population monitoring. Institute for Applied Ecology, Corvallis, Oregon, USDA Forest Service, Siuslaw National Forest, USDI Bureau of Land Management, Coos Bay District, and Oregon Department of Parks and Recreation. xi + 49 pp.
- Giles-Johnson, D.E.L. and T. N. Kaye. 2014. *Abronia umbellata* var. *breviflora* on the Oregon coast: Reintroduction and population monitoring. Institute for Applied Ecology, Corvallis, Oregon, USDA Forest Service, Siuslaw National Forest, USDI Bureau of Land Management, Coos Bay District, and Oregon Department of Parks and Recreation. xi + 63 pp.
- Hitchcock, C.L., A. Cronquist, M. Owenby, and J.W. Thompson. 1964. *Vascular Plants of the Pacific Northwest. Part 2: Salicaceae to Saxifragaceae*. University of Washington Press, Seattle.
- Kaye, T.N. 1995. Experimental reintroduction of the endangered pink sand-verbena on Oregon beaches. 1995 Report. Oregon Department of Agriculture, Salem, Oregon.
- Kaye, T.N. 1996. Experimental reintroduction of the endangered pink sand-verbena on Oregon beaches: direct seeding. Oregon Department of Agriculture, Salem, Oregon.
- Kaye, T.N. 1998. Experimental reintroduction of pink sand-verbena at four sites on the Oregon Coast: Coos Bay, New River, Tahkenitch, and Siltcoos. Oregon Department of Agriculture, Salem, Oregon.
- Kaye, T.N. 1999. Reintroduction of pink sand-verbena (*Abronia umbellata* var. *breviflora*) on dredge material and natural habitat. Final report. Oregon Department of Agriculture, Salem, Oregon.
- Kaye, T.N. 2006. Conservation strategy for pink sand-verbena (*Abronia umbellata* var. *breviflora*). Prepared by the Institute for Applied Ecology for Coos Bay District BLM, USFS Siuslaw National Forest, and Oregon Parks and Recreation Department. iv +44pp.
- Kaye, T.N. and J. Cramer. 2005. Experimental Reintroduction of Pink Sand-Verbena on the Oregon Coast: Transplanting and Seeding. Prepared by the Institute for Applied Ecology for Coos Bay District BLM, USFS Siuslaw National Forest, and Oregon Parks and Recreation Department.
- Kaye, T.N., M. Kirkland, and N. Testa. 1999. Growing endangered plants to save them: germinating, propagating and restoring pink sand-verbena. In, *Native plants: propagating and planting*. R. Rose and D. Haase, editors. Forest Research Lab, Oregon State University. Pp. 100-107.

- Kaye, T.N. 1999. Propagation of endangered species: Variable germination of pink sand-verbena from Pacific Coast beaches. Combined Proceedings of the International Plant Propagators Society 49:617-621.
- Kaye, T.N., K. Amsberry, S.D. Gisler, and R.J. Meinke. 1998. Back from the brink: the challenges of endangered plant re-introduction. Hortus West 9:4-9, 90-91.
- McGlaughlin, M., K. Karoly, and T.N. Kaye. 2002. Genetic variation and its relationship to population size in reintroduced populations of the pink sand verbena, *Abronia umbellata* subsp. *breviflora*. Conservation Genetics 3:411-420.
- Oregon Department of Fish and Wildlife. 1994. Oregon conservation program for the western snowy plover (*Charadrius alexandrinus nivosus*). Oregon Department of Fish and Wildlife, Portland, Oregon.
- Thorpe, A.S., D.E.L. Giles-Johnson, R.T. Massatti, and T.N. Kaye. 2009. *Abronia umbellata* var. *breviflora* on the Oregon coast: Reintroduction and population monitoring. Institute for Applied Ecology, Corvallis, Oregon, USDA Forest Service, Siuslaw National Forest, USDI Bureau of Land Management, Coos Bay District, and Oregon Department of Parks and Recreation

APPENDIX I. MATERIALS REQUIRED FOR POPULATION MONITORING

Population Monitoring:

pin flags

data sheets

GPS

previous year's report

6 100m tapes (Coos Bay only)

Flagging

Candy canes

Meter poles (1 per person)-- for measuring distance from tape, as well plant species within 0.5 meter of select plants)

Seeding:

GPS

Seed

Wooden Stakes

Mallet

Sharpie

APPENDIX II. SURVEY RESULTS FOR SEEDED AND NATURAL POPULATIONS

TABLE 12. RESULTS OF SURVEYS FOR PINK SAND-VERBENA AT COOS BAY DISTRICT BUREAU OF LAND MANAGEMENT SITES ALONG THE OREGON COAST. IAE CONDUCTED SURVEYS FROM 2000-2018. POPULATION DATA FOR YEARS PRIOR TO 2000 ARE FROM VARIOUS SOURCES.

Site	natural or reintroduced	date surveyed	population size	Reintroduction history and other comments
Coos Bay	reintro.	2018	~37,084 (repro.)	Seeded (100K seeds)
		2017	~199,503 (repro.)	Seeded (100K seeds)
		2016	~306,176 (repro.)	Seeded (100K seeds)
		2015	~264,757 (repro.)	Seeded (100K seeds)
		2014	~470,092 (repro.)	Seeded (100K seeds)
		2013	~74,804 (repro.)	Seeded (100K seeds)
		9/26/2012	~349,658 (repro.)	Seeded (100K seeds)
		10/6/2011	~185,623 (repro.)	Seeded (100K seeds)
		9/22/2010	~240,488 (repro.)	Seeded (100K seeds)
		10/14/2009	~203,959 (repro.)	Seeded (90K seeds)
		10/15/2008	>10,929 (repro.)*	Seeded (90K seeds)
		2007	~111,063 (repro.)	Seeded (100K seeds)
		9/27/2006	~99,354 (repro.)	Seeded (100K seeds)
		10/3/2005	~31,411 (repro.)	---
		10/12/2004	~66,697 (repro.)	---
10/2/2003	~111,496 (repro.)	---		
9/25/2002	~65,922 (repro.)	---		
9/14/2001	~45,257 (repro.)	---		
9/19/2000	~45,000 (repro.)	---		
1996		---	Seeded (50K seeds)	

New reintro. River	2018	196 plants(124 repro., 72veg.)	Seeding focused in overwashes (100 k seeds)
	2017	9 plants (6 repro., 3 veg.)	Seeded (110k seeds)
	2016	108 plants (30 repro., 78 veg.)	Seeded (100k seeds)
	2015	1,824 plants (287 repro., 1,537 veg.)	Seeded (100k seeds)
	2014	366 plants (160 repro., 206 veg.)	Seeded (100k seeds)
	2013	33 plants (8 repro., 25 veg.)	Seeded (120k seeds)
	9/27/2012	66 plants (26 repro., 40 veg.)	Seeded (120k seeds)
	10/5/2011	*237 plants (44 repro., 193 veg.)	Seeded (100k seeds)
	9/21/2010	810 plants (369 repro., 441 veg.)	Seeded (100K seeds)
	10/5/2009	62 plants (42 repro., 20 veg.)	Seeded (80K seeds)
	10/16/2008	180 plants (69 repro., 111 veg.)	Seeded (80K seeds)
	9/29/2007	480 plants (269 repro., 211 veg.)	Seeded (150K seeds)
	9/25/2006	616 plants (380 repro., 236 veg.)	Seeded (100K seeds)
	10/4/2005	2,174 plants (1,114 repro., 1,058 veg.)	Seeded (95K seeds)
	9/14/2004	1,628 plants (1,014 repro., 614 veg.)	Seeded (100K seeds)
	10/1/2003	917 plants (459 repro., 458 veg.)	Seeded (80K seeds)
	9/11/2002	524 plants (373 repro., 151 veg.)	Seeded (100K seeds)
	9/12/2001	145 plants	Seeded (70K seeds)
	8/15/2000	421 plants	Seeded (50K seeds)
	1999	---	Seeded (50K seeds)
1997	---	Seeded (50K seeds)	

TABLE 13. RESULTS OF SURVEYS FOR PINK SAND-VERBENA SITES MANAGED BY THE SIUSLAW NATIONAL FOREST ALONG THE OREGON COAST. IAE CONDUCTED SURVEYS FROM 2000-2012. POPULATION DATA FOR YEARS PRIOR TO 2000 ARE FROM VARIOUS SOURCES.

Site	natural or reintroduced	date surveyed	population size	Reintroduction history and other comments
Tahkenitch Cr.	reintro.	2018	361 plants (117 repro., 244 veg.)	Seeded (90k seeds)
		2017	160 plants (23 repro., 137 veg.)	Seeded (100k seeds)
		2016	141 plants (76 repro., 65 veg.)	Seeded (100k seeds)
		2015	100 plants (65 repro., 35 veg.)	Seeded (100k seeds)
		2014	100 plants (43 repro., 57 veg.)	Seeded (100k seeds)
		2013	13 plants (5 repro., 8 veg.)	Seeded (100k seeds)
		2012	381 plants (135 repro., 246 veg.)	Seeded (100k seeds)
		10/5/11	95 plants (38 repro., 57 veg.)	Seeded (80k seeds)
		10/09	4 plants (2 repro., 2 veg.) *briefly surveyed by Marty Stein	---
			1 veg.	---
		10/4/06	0 plants	---
		9/14/05	121 plants (72 repro., 49 veg.)	Seeded (50K seeds)
		9/29/04	0 plants	Seeded (50K seeds)
		9/18/03	22 plants (14 repro., 8 veg.)	---
9/25/02	50 plants (45 repro., 5 veg.)	Seeded (50K seeds)		
9/18/01		Seeded (50K seeds) +50 transplants		
9/20/00	32 plants (repro.)	50 transplants		
Overlook North	reintro.	2018	497 plants (284 repro., 213 veg.)	Seeded (75k seeds)
		2017	616 plants (375 repro., 241 veg.)	Seeded (60k seeds)
		2016	269 plants (158 repro., 111 veg.)	Seeded (60k seeds)
		2015	124 plants (71 repro., 53 veg.)	Seeded (60k seeds)
		2014	281 plants (149 repro., 132 veg.)	Seeded (60k seeds)
		2013	994 plants (211 repro., 783 veg.)	Seeded (60k seeds)
		10/1/12	4,860 plants (1,627 repro., 3,233 veg.)	Seeded (60k seeds)

		10/4/11	1,938 plants (794 repro., 1,194 veg.)	Seeded (60k seeds)
		9/26/10	730 plants (238 repro., 492 veg.)	Seeded (50K seeds)
		10/6/09	789 plants (466 repro., 323 veg.)	Seeded (50K seeds)
		10/2/08	2,395 plants (811 repro., 1,584 veg.)	Seeded (100K seeds)
		9/24/07	3,445 plants (2,790 repro., 655 veg.)	Seeded (80K seeds)
		10/4/06	7,825 plants (1,976 repro., 5,840 veg.)	---
		9/14/05	3,210 plants (1,191 repro., 2,019 veg.)	---
		9/29/04	3,741 plants (2,632 repro., 1,109 veg.)	---
		9/18/03	1,478 plants (1,359 repro., 119 veg.)	---
		9/25/02	1,091 plants (574 repro., 517 veg.)	---
		9/18/01	482 plants (390 repro., 92 veg.)	Seeded (50K seeds)
		9/20/00	2 plants (repro.)	50 transplants (most pulled by accident)
Overlook South	reintro.	2018	3,476 plants (1,850 repro., 1,626 veg.)	Seeded (90k seeds)
		2017	966 plants (618 repro., 348 veg.)	Seeded (60k seeds)
		2016	614 plants (375 repro., 239 veg.)	Seeded (60k seeds)
		2015	158 plants (121 repro., 37 veg.)	Seeded (60k seeds)
		2014	109 plants (66 repro., 43 veg.)	Seeded (60k seeds)
		2013	482 plants (87 repro., 395 veg.)	Seeded (60k seeds)
		10/1/12	2,247 plants (628 repro., 1,619 veg.)	Seeded (60k seeds)
		10/4/11	2,349 plants (988 repro., 1,361 veg.)	Seeded (60k seeds)
		9/26/10	1,052 plants (424 repro., 628 veg.)	Seeded (50K seeds)
		10/6/09	2,073 plants (1518 repro., 555 veg.)	Seeded (50K seeds)
		2008	Informal survey noted plants as present	Seeded (50K seeds)
		9/24/07	98 plants (76 repro., 22 veg.)	Seeded (50K seeds)
		10/4/06	10 plants (4 repro., 6 veg.)	---
		9/14/05	4,340 plants (2,581 repro., 1,759 veg.)	Seeded (50K seeds)
		9/29/04	9,554 plants (6,325 repro., 3,229 veg.)	---
		9/18/03	2,107 plants (1,954 repro., 153 veg.)	---
		9/25/02	1,726 plants (1,435 repro., 291 veg.)	Seeded (50K seeds)
		9/18/01	658 plants (427 repro., 231 veg.)	Seeded (50K seeds)
Baker Beach	reintro.	2018		0 Seeded (35K seeds with Sutton Creek)
		10/6/09	6 plants (5 repro., 1 veg.)	Seeded (40K seeds)
		10/2/08	70 plants (32 repro., 38 veg.)	Seeded (50K seeds)
		9/24/07	42 plants (30 repro., 12 veg.)	Seeded (50K seeds)
		9/5/06	12 plants (6 repro., 6 veg.)	Seeded (50K seeds)
		9/14/05	72 plants (11 repro., 61 veg.)	Seeded (50K seeds)
		10/28/04	93 plants (37 repro., 56 veg.)	Seeded (50K seeds)

		9/18/03	55 plants (27 repro., 28 veg.)	Seeded (50K seeds)
Siltcoos Cr.	reintro.	2018	292 plants (147 repro., 145 veg.)	Seeded (50K seeds)
		2017	116 plants (66 repro., 50 veg.)	Seeded (50K seeds)
		2016	427 plants (330 repro., 97 veg.)	Seeded (50K seeds)
		2015	298 plants (174 repro., 124 veg.)	Seeded (50K seeds)
		2014	10 plants (8 repro., 2 veg.)	Seeded (50K seeds)
		2013	38 plants (31 repro., 7 veg.)	Seeded (50K seeds)
		10/1/12	605 plants (135 repro., 470 veg.)	Seeded (50K seeds)
		10/5/11	415 plants (309 repro., 106 veg.)	Seeded (100K seeds)
		9/27/10	148 plants (28 repro., 120 veg.)	Seeded (50K seeds)
		10/6/09	36 plants (15 repro., 21 veg.)	Seeded (55K seeds)
		10/2/08	41 plants (14 repro., 27 veg.)	Seeded (60K seeds)
		9/24/07	66 plants (54 repro., 12 veg.)	Seeded (100K seeds)
		10/4/06	385 plants (202 repro., 183 veg.)	Seeded (50K seeds)
		9/14/05	989 plants (961 repro., 28 veg.)	Seeded (50K seeds)
		10/28/04	355 plants (311 repro., 44 veg.)	Seeded (50K seeds)
		9/18/03	215 plants (195 repro., 20 veg.)	---
		9/25/02	19 plants (12 repro., 7 veg.)	Seeded (60K seeds)
		9/18/01	7 plants (all repro.)	45 Transplants
		9/20/00	249 plants (135 repro., 114 veg.)	Seeded (50K seeds)
		1999		--- Seeded (50K seeds)
Sutton Creek	reintro.	2018	28 plants (18 repro., 10 veg.)	Seeded 35 k with Baker Beach
		9/24/07	0 plants	---
		9/5/06	0 plants	Seeded (50K seeds)
		2005	0 plants	Seeded (50K seeds)
		10/28/04	150 plants (28 repro., 122 veg.)	Seeded (50K seeds)
Tenmile Cr.	natural and reintro.	10/23/03	1 plant (repro.)	---
		9/25/02	0 plants	---
		9/18/01	0 plants	---
		9/20/00	0 plants	---
		1999		--- 150 transplants

TABLE 14. RESULTS OF SURVEYS FOR PINK SAND-VERBENA SITES MANAGED BY THE OREGON DEPARTMENT OF PARKS AND RECREATION ALONG THE OREGON COAST. IAE CONDUCTED SURVEYS FROM 2000-2012. POPULATION DATA FOR YEARS PRIOR TO 2000 ARE FROM VARIOUS SOURCES.

Site	natural or reintroduced	date surveyed	population size	Reintroduction history and other comments
Bandon Beach South	reintro.	9/27/2012	161 plants (84 repro., 77 veg.)	Seeded (100K seeds)
		10/2011	476 plants (173 repro., 303 veg.)	Seeded (100K seeds)
		9/21/2010	260 plants (91 repro., 149 veg.)	Seeded (120K seeds)
		10/14/2008	113 plants (52 repro., 61 veg.)	Seeded (50K seeds)
China Creek (Bandon Beach)		9/27/2012	6 plants (3 repro., 3 veg.)	---
		9/13/2007	173 plants (121 repro., 52 veg.)	Seeded (50K seeds)
		9/26/2006	452 plants (346 repro., 106 veg.)	Seeded (50K seeds)
		10/4/2005	139 plants (124 repro., 15 veg.)	Seeded (100K seeds)
Bastendorff Beach	reintro.	9/21/2010	2 plants (2 repro., 0 veg.)	---
		10/15/2008	9 plants (6 repro., 3 veg.)	---
		9/13/2007	2 plants (repro.)	---
		9/26/2006	11 plants (8 repro., 3 veg.)	---
		10/5/2005	536 plants (410 repro., 126 veg.)	Seeded (50K seeds)
		9/13/2004	371 plants (245 repro., 126 veg.)	Seeded (50K seeds)
		10/2/2003	110 plants (104 repro., 6 veg.)	---
		9/11/2002	13 plants (8 repro., 5 veg.)	Seeded (50K seeds)
		8/15/2000	0 plants	---
		1998	---	Seeded (50K seeds)
1997	---	Seeded (50K seeds)		
1996	---	Seeded & transplanted		
1995	---	Seeded & transplanted		

Harris Beach State Park	natural	9/15/2004	0 plants	Natural pop'n, but also seeded (5,000 seeds) in 1995 and transplanted
		9/10/2002	0 plants	
		9/13/2001	0 plants	
		8/16/2000	0 plants	
McVay Park	natural	8/16/2000	0 plants	appeared unsuitable
Pistol River	reintro.	10/5/2009	0 plants	---
		10/14/2008	1 veg.	Seeded (80K seeds)
Euchre Creek (Ophir)	reintro.	9/15/2004	2 veg. plants	---
		9/30/2003	7 plants (4 repro., 3 veg.)	---
		9/10/02 9/13/01	9 plants (all repro.)	Seeded (50K seeds)
		8/16/2000	0 plants	---
		1999	1 veg. plant	Seeded (50K seeds)
		---		Seeded (50K seeds)

TABLE 15. RESULTS OF SURVEYS FOR PINK SAND-VERBENA SITES MANAGED BY THE USFWS ALONG THE OREGON COAST. IAE CONDUCTED SURVEYS FROM 2009-2012.

Site	natural or reintroduced	date surveyed	population size	Reintroduction history and other comments
Elk River	reintro	9/25/2012	556 plants (384 repro., 172 veg.)	Seeded (100k seeds)
		10/4/2011	307 plants (105 repro., 202 veg.)	Seeded (100k seeds)
		9/21/2010	113 plants (86 repro, 27 veg.)	Seeded (100k seeds)
		9/20/2009	389 plants (353 repro., 36 veg.)	Seeded (100k seeds)
Crook Point	reintro	9/21/2010	2 (1 repro., 1 veg.)	Seeded (50k seeds)

TABLE 16. RESULTS OF SURVEYS FOR PINK SAND-VERBENA SITES ALONG THE OREGON COAST MANAGED BY VARIOUS LAND MANAGERS INCLUDING THE STATE PARKS DEPARTMENT. IAE CONDUCTED SURVEYS FROM 2000-2012. POPULATION DATA FOR YEARS PRIOR TO 2000 ARE FROM VARIOUS SOURCES.

Site	natural or reintroduced	date surveyed	population size	Reintroduction history and other comments
Charleston Beach (State and County Parks)	reintro.	9/21/2010	0 plants	---
		9/11/2002	0 plants	---
		8/15/2000	0 plants	---
		1996	---	Seeded and transplanted
		1995	---	Seeded and transplanted
Otter Point - Oregon Department of Parks and Recreation	natural	9/13/2007	0 plants	2000 was the first year since 1993 that no plants were observed at this site
		9/26/2006	0 plants	---
		9/15/2004	0 plants	
		9/30/2003	0 plants	
		9/10/2002	9 repro. plants	
		9/13/2001	3 repro. plants	
		8/16/2000	0 plants	
		1999	9 plants	
Heceta Beach Oregon Department of Parks and Recreation	reintro.	9/12/2001	0 plants	---
		10/18/2000	0 plants	---
		1995	---	Seeded (5K seeds)
	natural	9/12/2009	1 flowering plant	N42.8655 W124.53432

				(identified by D. & D. Bilderback)
N of Cape Blanco/ S of Blacklock pt. Oregon Department of Parks and Recreation	natural	9/26/2006	1 veg. plant	---
		9/15/2004	7 plants (3 repro., 4 veg.)	---
		10/1/2003	0 plants	---
		8/16/2000	2 repro., 1 N & 1 S of creek.	---
		1999	1 plant	---
		1984	several plants	---
Floras Lake - Oregon Department of Parks and Recreation/BL M	reintro.	9/25/2012	67 plants (8 repro., 59 veg.)	---
		10/6/2009	0 plants	---
		9/13/2007	0 plants	---
		9/26/2006	65 plants (41 repro., 24 veg.)	---
		10/4/2005	20 plants (all repro.)	---
		9/14/2004	220 plants (81 repro., 139 veg.)	---
		10/19/2000	10 plants (7 repro., 3 veg.)	Seeded (50K seeds)
		1996	7 plants	Seeded (5K seeds)
Hubbard Creek- Oregon Department of Parks and Recreation	reintro.	9/14/2004	4 repro. plants	---
		10/1/2003	0 plants	---
		9/11/2002	0 plants	---
		8/15/2000	0 plants	---

		1999		1 plant	---
		1996		---	Seeded & transplanted
		1995		---	Seeded & transplanted
Port Orford-Oregon Department of Parks and Recreation	natural & reintro	10/2011		146 reproductive plants	---
		9/21/2010		136 reproductive plants	---
		10/20/2009		269 plants (237 repro., 32 veg.)	---
		10/16/2008		226 plants (only repro. counted)	---
		9/13/2007		1,412 plants (846 repro., 566 veg.)	---
		9/26/2006		660 plants (454 repro., 206 veg.)	---
		10/4/2005		740 plants (392 repro., 348 veg.)	---
		9/14/2004		909 plants (556 repro., 353 veg.)	---
		9/30/2003		552 plants (159 repro. plants, 393 veg.)	---
		9/10/2002		1,146 plants (480 repro. and 666 veg.)	---
		9/13/2001		2,607 (1,467 repro. and 1,140 veg.)	---
		2000		1,834 plants	---
		1999		7,169 plants	---
Winchuck River - Oregon Department of Parks and Recreation	natural	9/10/2002		6 repro. plants	New site in 2000; N & S of creek mouth
		8/16/2000		5 repro., 1 veg.	
Salmon River - Camp Westwind (Westwind Stewardship Group)	reintro.	9/22/2012		0 plants	---
		9/26/2009		0 plants	---
		10/22/2005		92 plants (52 repro., 40 veg.)	---
		11/20/2004		1 repro. plant	---

10/24/2002	163 plants (129 repro., 34 veg.)	Seeded (50K seeds)
9/10/2001	741 plants (488 repro. 253 veg.)	Seeded (40K seeds)

APPENDIX III. MAPS OF SELECTED SITES- SEEDED OR MONITORED


New River



Overlook North and South



Siltcoos



Tahkenitch

