

Scientific Name: *Notropis jemezanus*

Common Name: Rio Grande shiner

BISON No.: 010435

Legal Status:

- | | | |
|---------------------------------------|------------------------------|------------------------------|
| ➤ Arizona, Species of Special Concern | ➤ ESA, Proposed Threatened | ➤ New Mexico-WCA, Threatened |
| ➤ ESA, Endangered | ➤ ESA, Threatened | ➤ USFS-Region 3, Sensitive |
| ➤ ESA, Proposed Endangered | ➤ New Mexico-WCA, Endangered | ➤ None |

Distribution:

- | | |
|---|---------------------------|
| ➤ Endemic to Arizona | ➤ Southern Limit of Range |
| ➤ Endemic to Arizona and New Mexico | ➤ Western Limit of Range |
| ➤ Endemic to New Mexico | ➤ Eastern Limit of Range |
| ➤ Not Restricted to Arizona or New Mexico | ➤ Very Local |
| ➤ Northern Limit of Range | |

Major River Drainages:

- | | |
|------------------------|-----------------------------|
| ➤ Dry Cimmaron River | ➤ Rio Yaqui Basin |
| ➤ Canadian River | ➤ Wilcox Playa |
| ➤ Southern High Plains | ➤ Rio Magdalena Basin |
| ➤ Pecos River | ➤ Rio Sonoita Basin |
| ➤ Estancia Basin | ➤ Little Colorado River |
| ➤ Tularosa Basin | ➤ Mainstream Colorado River |
| ➤ Salt Basin | ➤ Virgin River Basin |
| ➤ Rio Grande | ➤ Hualapai Lake |
| ➤ Rio Mimbres | ➤ Bill Williams Basin |
| ➤ Zuni River | |
| ➤ Gila River | |

Status/Trends/Threats (narrative):

Extirpated in the Rio Grande and stable in the Pecos River.

Distribution (narrative):

Native to the Rio Grande and Pecos river drainages. Historically, as far north as San Ildefonso in the Rio Grande. In the Pecos, once at far north as Santa Rose, but now restricted to the Pecos south of Sumner Lake.

Key Distribution/Abundance/Management Areas:

Panel key distribution/abundance/management areas:

Breeding (narrative):

No information on the breeding biology of this species. Spawns from mid May to late August.

Habitat (narrative):

Large, open rivers with laminar flows over sand-gravel-rubble bottoms with little aquatic vegetation. Silt if often present over substrates.

Key Habitat Components: laminar flow, variable substrates with sand and silt

Breeding Season:

- January
- February
- March
- April
- May
- June
- July
- August
- September
- October
- November
- December

Panel breeding season comments:**Aquatic Habitats:****Large Scale:**

- Rivers
- Streams
- Springs
- Spring runs
- Lakes
- Ponds
- Sinkholes
- Cienegas
- Unknown
- Variable

Small Scale:

- Runs
- Riffles
- Pools
- Open Water
- Shorelines

Panel comments on aquatic habitats:

Important Habitat Features (Water characteristics):

Current

- Fast (> 75 cm/sec)
- Intermediate (10-75 cm/sec)
- Slow (< 10 cm/sec)
- None
- Unknown
- Variable

Gradient

- High gradient (>1%)
- Intermediate Gradient (0.25-1%)
- Low Gradient (<0.25%)
- None
- Unknown
- Variable

Water Depth

- Very Deep (> 1 m)
- Deep (0.25-1 m)
- Intermediate (0.1-0.25 m)
- Shallow (< 0.1 m)
- Unknown
- Variable

Panel comments on water characteristics:

Important Habitat Features (Water Chemistry)

Temperature (general)

- Cold Water (4-15°C)
- Cool Water (10-21°C)
- Warm Water (15-27°C)
- Unknown
- Variable

Turbidity

- High
- Intermediate
- Low
- Unknown
- Variable

Conductivity

- Very High (> 2000 $\mu\text{S}/\text{cm}$)
- High (750-2000 $\mu\text{S}/\text{cm}$)
- Intermediate (250-750 $\mu\text{S}/\text{cm}$)
- Low (< 250 $\mu\text{S}/\text{cm}$)
- Unknown
- Variable

Panel comments on water chemistry:

Important Habitat Features (Structural elements):

Substrate

- Bedrock
- Silt/Clay
- Detritus
- Sand
- Gravel
- Cobble
- Boulders
- Unknown
- Variable

Cover

- Rocks, boulders
- Undercut banks
- Woody debris
- Aquatic vegetation
- Rootwads
- Not important
- Overhanging vegetation
- Unknown
- Variable

Panel comments on structural elements:

Diet (narrative):

Gut morphology suggests the species is carnivorous-omnivorous, typical of most species of the genus *Notropis*.

Diet category (list):

- Planktivore
- Herbivore
- Insectivore
- Piscivore (Fish)
- Omnivore
- Detritivore

Grazing Effects (narrative):

Based on habitats occupied, the species will probably not be affected by grazing. Because silt and sand are a component of substrates occupied by the species, increased input of fine materials would likely have no effect on breeding or feeding biology.

Panel limiting habitat component relative to grazing and comments:
<p>Panel assessment: Is this species a priority for selecting a grazing strategy? Throughout the species' distribution in New Mexico and Arizona YES NO UNKNOWN In key management area(s) YES NO UNKNOWN</p>

Principle Mechanisms Through Which Grazing Impacts This Species (list):

May be Revised

- | | | |
|---|---|--|
| <ul style="list-style-type: none"> ➤ Alteration of bank structures ➤ Alteration of substrate ➤ Alteration of water regimes ➤ Altered stream channel characteristics ➤ Altered aquatic vegetation composition | <ul style="list-style-type: none"> ➤ Altered bank vegetation structure ➤ Change in food availability ➤ Change in water temperature ➤ Change in water quality ➤ Habitat fragmentation | <ul style="list-style-type: none"> ➤ Increased turbidity ➤ Other biotic factors ➤ Parasites or pathogens ➤ Population genetic structure loss ➤ Range improvements ➤ Trampling, scratching ➤ Unknown |
|---|---|--|

Panel causal mechanisms comments:
--

Authors

- **Draft:**
- **GP 2001:**
- **GP 2002:**
- **Revision:**

Bibliography:

Koster, W. J. 1957. Guide to the fishes of New Mexico. Univ. of New Mexico Press, Albuquerque.

Sublette, J. E., M. D. Hatch, and M. Sublette. 1990. The Fishes of New Mexico. University of New Mexico Press. Albuquerque. 393 pp.