

THE FIRE EFFECTS INFORMATION SYSTEM — SERVING MANAGERS SINCE BEFORE THE YELLOWSTONE FIRES

Jane Kapler Smith,¹ Janet L. Fryer,² and Kristin Zouhar³

¹ Ecologist, USDA FS Rocky Mountain Research Station

² Ecologist, USDA FS Rocky Mountain Research Station

³ Ecologist, USDA FS Rocky Mountain Research Station

This presentation will describe the current status of the Fire Effects Information System (FEIS) and explore lessons learned from this 23-year-old project about the application of science to fire management issues. FEIS contains literature reviews covering biology and fire ecology for approximately 1,100 species in North America: plants and animals, native and nonnative. Established in 1985 and continually updated, the system has served managers, researchers, students, and the general public for >20 years. Species reviews in FEIS provide information that can help managers plan a prescribed fire, write a fire or fuel management plan, determine the need for post-fire rehabilitation, or assess the potential for increase of invasive species after fire. FEIS is cited in nearly half of EISs written by federal wildland fire managers. Research Project Summaries, a recent addition to FEIS, supplement species reviews and provide information on fire effects for an additional 250 species. To ensure that the latest science is within reach of the manager seeking information, FEIS reviews are clearly organized and widely available on the Internet. Reviews are updated depending on needs identified by users and support available. To use science syntheses effectively, managers must combine critical reading with understanding of the local ecosystem and its condition. FEIS reviews are constructed to convey general patterns of plant and animal response reported in the literature and also to describe exceptions to those patterns. Managers can use this information to infer the likelihood that local responses will fit the patterns and exceptions reported in the literature. FEIS reviews identify uncertainties, contradictions in research findings, and knowledge gaps so managers and planners can be aware of inconsistent findings and topics that are not well understood. FEIS reviews describe the location and results of individual studies to indicate their breadth of application, answering questions such as: Is a report based on a single observation or an extensive field study? Is it limited to a small geographic region or representative of a large area? Is it reported with a known level of confidence resulting from statistical analysis, or from anecdotal observation? The level of certainty conveyed in FEIS reviews may indicate to managers that a particular plant or animal response to fire is likely and chances of different results are low, or that a response is possible but uncertain, warranting post-fire monitoring and possibly an adaptive response by management.

SOME LIKE IT HOT: USING BURN INFORMATION TO BETTER PREDICT INVASIVE SPECIES POST-FIRE RESPONSES

Joel Silverman,¹ Monique Rocca,² Jeffrey Morissette,³ Peter Ma,⁴ and Nate Benson⁵

¹ Graduate Research Assist./Lead Fire Effects Monitor, Colorado State University/NPS

² Professor of Fire Science, Colorado State University

³ Research Scientist, NASA Goddard Space Flight Center

⁴ Remote Sensing/GIS Analyst, NASA Goddard Space Flight Center/Innovim

⁵ Fire Ecologist, National Park Service - National Interagency Fire Center

A growing body of literature links fire to the rapid colonization and proliferation of particular invasive plants. Despite this work, the potential for increased risk of invasion due to fire remains unknown for many exotic species at the regional level. We employed habitat suitability models (statistical models linking species responses at specific locations to environmental conditions) in order to 1) test species responses to burn variables, 2) create species habitat maps, and 3) infer relationships between fire and the species of interest based on parameter estimates. Whereas most habitat suitability models omit disturbance processes such as fire, we specifically incorporated burn characteristics (burn severity, occurrence, and time-since-fire), in addition to a standard suite of environmental variables used to predict species potential distributions (e.g., topographic, vegetative, and hydrologic data). We tested models using occurrence data for two exotic species that are currently invading the Greater Yellowstone Ecosystem: *Carduus nutans* (musk thistle) and *Linaria dalmatica* (Dalmatian toad-flax). Our results indicate that both species are influenced by burn characteristics and that for *C. nutans*, the effect of time-since-burn depends on the vegetation type. The relative odds of *C. nutans* presence continues to rise for the first decade following fires within several vegetation types, but the relative odds decreases during this range of years for other types. Each species also responded differently to burn severity (as measured through the Landsat-derived index, dNBR). Since each species responds to multiple environmental factors, habitat suitability models that include burn variables afford a greater degree of inference and potential predictive ability than either nonburn habitat suitability maps or direct burn-severity (dNBR) maps such as those used by Burned Area Emergency Rehabilitation teams. The habitat suitability maps we present can be used for demarcating areas of concern prior to prescribed fire as well as directing post-fire invasive species management at a regional to national level.