

Rutaceae—Rue family

Phellodendron amurense Rupr.

Amur corktree

Ralph A. Read and John C. Zasada

Dr. Read retired from the USDA Forest Service Rocky Mountain Research Station;
Dr. Zasada retired from the North Central Research Station

P

Growth habit, occurrence, and use. Amur corktree—*Phellodendron amurense* Rupr.—is native to northern China, Manchuria, Korea, and Japan. This small to medium deciduous tree—25 to 50 feet tall—has been cultivated in the Far East and eastern Europe. It was introduced into the United States around 1865, and its thick, corky bark and massive, irregular branches have made it of special interest for landscape and environmental plantings in the northern and western United States (Blackburn 1952; Everett 1964; Hoag 1965; Lewis 1957). In tests in Kansas, however, the tree did not perform well and was not recommended for general use (Hensley and others 1991). It is a potential source of industrial cork (Izmodenov 1972; Ota and others 1965), important as a nectar-bearing species in bee-keeping areas of the Russian Far East (Necaev and Pelemenev 1965), and of possible importance for the insecticidal properties of its fruit oils (Schechter 1943). In Byelorussia it is considered a “soil builder” when mixed with Scots pine—*Pinus sylvestris* L. (Letkovskij 1960). It tolerates a wide range of soil conditions, pH, drought, and pollution; it is easily transplanted and generally free of pests.


Flowering and fruiting. The species is dioecious, and female plants develop tend to have a bushier form than males (Hensley and others 1991). Small, yellowish green flowers, in large clusters of terminal panicles, appear in May and June (Krecetova 1960; Rehder 1940; Schechter 1943). Climate affects the time of day when flowers open, pollination, and the longevity of flowers (Starshova 1972).

Fruits are subglobose drupes about 1 cm in diameter (figure 1), green to yellowish green, turning black when ripe in September and October (Read 1974). They remain on the terminal panicles long after the leaves have dropped. Fruits are borne singly on short stalks (figure 1) and are very oily and aromatic. Each fruit usually contains 2 or 3 full-sized seeds and 3 or 4 aborted seeds (Read 1974). Seeds are brown to black, 5 mm long, 2 mm wide, and about 1 mm thick (figures 2 and 3); they have a moderately hard, stony coat (Gorokhova 1981; Read 1974).

Figure 1—*Phellodendron amurense*, Amur corktree: fruit cluster.




Figure 2—*Phellodendron amurense*, Amur corktree:



Minimum seed-bearing age is 7 to 13 years, both within the natural range and where the species has been introduced (Atkimockin 1960; Gorokhova 1981; Maljcev 1950; Read 1974; Starshova 1972). No data are available on the frequency of seedcrops. Severe drought had no marked effect on the

Figure 3—*Phellodendron amurense*, Amur corktree: longitudinal section through 2 planes of a seed.



morphology of fruits and seeds, but appeared to reduce seed quality (Gorokhova 1986).

Collection, extraction, and storage. The terminal panicles of fruit may be harvested with pruning shears in late September through October after leaf fall. After that, although many fruits remain tightly on the tree, some will have fallen. Fruits should be spread out in shallow layers to prevent heating and mildew during air-drying. Fruits may be soaked in water and seeds squeezed from the fleshy matter by hand; large lots can be run through a macerator and separated from the pulp by flotation. Fresh fruits weigh about 57 kg/hl (44 lb/bu) and yield about 0.9 kg (2 lb) of cleaned seeds (Read 1974). Based on seeds from 2 different lots, 1 kg contained 58,960 to 80,000 (26,800 to 36,363/lb) and 96,800 to 105,600 (44,000 to 48,000/lb) cleaned seeds (Read 1974; Swingle 1939). Seeds collected from plants growing in the natural range had similar seed weights (Gorokhova 1981).

Germination. Fresh seeds germinate well without pretreatment (Dirr 1990; Read 1974). However, there are a number of reports of greatly improved germination following stratification. Stratification is recommended for seeds stored any length of time (Dirr and Heuser 1987).

Germination for a seedlot (for which the handling and storage history was not described) was best following cold moist, underground stratification for 166 days (Timm 1989). In another study, seeds stratified for 8 weeks had a higher germination rate and the same germination percentage as seeds stratified 4 weeks; germination of unstratified seeds was less than half that of stratified seeds (Mukai and Yokoyama 1985). Based on the information available, it is recommended that seeds be stratified if the history of collection, handling, and storage is not documented. Seeds of other *Phellodendron* spp. vary in their requirements for stratification (Dirr and Heuser 1987; Lin and others 1994).

Seeds germinate best at alternating temperatures. Both Lin and others (1979) and Mukai and Yokoyama (1985) reported germination of 3% or less at constant temperatures and 75 to 90% at alternating temperatures. The best temperature regimes were 35/5 °C and 35/15 °C (day length and high temperature for 8 hours).

Nursery practice and natural regeneration. In its natural range, natural regeneration sometimes occurs in dense groups. Although the corktree has been reported to sucker from its roots (Dirr and Heuser 1987), this dense regeneration is believed to be from seeds present in the forest floor (Soludukin 1977). Light fire or disturbances that result in drying and warming of the forest floor are believed to promote this development. There was no indication regarding the longevity of the seeds in the forest floor environment, however the endocarp is moderately hard and might facilitate longevity under these conditions (Soludukin 1977).

In the nursery, untreated seeds may be sown in the fall (Read 1974) or stratified through winter for spring-seeding (Yerkes 1945). It is suggested that the best time for spring sowing is when the mean daily soil temperature has reached 8 to 10 °C (Antonyuk 1987). Trees may also be propagated vegetatively by root cuttings or shoot cuttings (Bailey 1947; Dirr and Heuser 1987).

References

- Antonyuk ED. 1987. Growth of seedlings in a polythene greenhouse in relation to sowing times. *Lesnoe Khozyaistvo* 4: 36–37.
- Atkimockin NG. 1960. [in Russian: Trials with *Phellodendron* spp. at the Forest-Steppe Experiment Station [abstract]. *Byulleten Glavnogo Botanicheskogo Sada [Moskva]* 37: 30–33.
- Bailey LH. 1947. *Standard cyclopedia of horticulture*. 2nd ed. New York: Macmillan. 1388 p.
- Blackburn BC. 1952. *Trees and shrubs of eastern North America*. New York: Oxford University Press. 358 p.
- Dirr MA. 1990. *Manual of woody landscape plants: their identification, ornamental characteristics, culture, propagation, and uses*. Champaign, IL: Stipes Publishing. 1007 p.
- Dirr MA, Heuser CW Jr. 1987. *The reference manual of woody plant cultivation: from seed to tissue culture*. Athens, GA: Varsity Press. 239 p.
- Everett TH. 1964. *New illustrated encyclopedia of gardening*. Volume 8. New York: Greystone Press: 1455–1456.
- Gorokhova GI. 1981. Fruiting of far-eastern species of plants under conditions of the forest steppe zone of western Siberia. *Soviet Journal of Ecology* 12 (4): 219–222 [translated and published 1982].
- Gorokhova GI. 1986. Effect of drought on the seed quality of exotic plants from the Soviet East [abstract]. *Byulleten Glavnogo Botanicheskogo Sada [Moskva]* 139: 85–89.
- Hensley DL, Wiest SC, Long CE, Pair JC, Gibbons FD III. 1991. Evaluation of 10 landscape trees for the Midwest. *Journal of Environmental Horticulture* 9(3): 149–155.

- Hoag DG. 1965. Trees and shrubs of the northern plains. Minneapolis: Lund Press. 376 p.
- Izmodenov AG. 1972. Multiple use of *Pinus koraiensis* forests [English abstract]. Moscow: Lenaya Promyshlennost. 122 p.
- Krecetova NV. 1960. [in Russian: The biology of flowering and fertilization in *P. amurensis*. abstract]. Botanicheski Zhurnal 45(9): 1336–1340.
- Letkovskij AI. 1960. [in Russian: Planting *Phellodendron amurense* to increase the productivity of Scots pine forest (in Belorussia.)] Lesnaya Hortus 12(1): 36–37 [Forestry Abstracts 23: 548, 1962].
- Lewis CE. 1957. American elm substitutes. American Nurseryman 106(10): 10–11, 50–51.
- Lin TP, TW Hu, Leu JJ. 1979. The effect of alternating temperatures on the germination of the seed of *Phellodendron* spp. [abstract]. Bulletin of the Taiwan Forest Research Institute 322: 1–10.
- Lin TP, Chen MH, Lin CH. 1994. Dormancy in seeds of *Phellodendron wilsonii* is mediated in part by abscisic acid. Plant Cell Physiology 35(1): 115–119.
- Mukai Y, Yokoyama T. 1985. Effect of stratification on seed germination of *Phellodendron amurense* [abstract]. Journal of the Japanese Forestry Society 67(3): 103–104.
- Necaev AD, Pelmenev VK. 1965. [in Russian: The importance of *Phellodendron amurense* as a nectar-bearing species.] Rastitel'nye Resursy [Moskva] 1(3): 419–423 [Forestry Abstracts 28: 621; 1967].
- Ota M and others. 1965. [in Japanese: A study of the manufacturing conditions and properties of SIS hardboard from the inner bark of Kihada (*Phellodendron amurense* Rupr.)] Wood Industries Tokyo 20(6): 13–16 [Forestry Abstracts 27: 1475; 1966].
- Read RA. 1974. *Phellodendron amurense* Rupr., Amur corktree. In: Schopmeyer CS, tech. coord. Seeds of woody plants in the United States. Agric. Handbk. 450. Washington, DC: USDA Forest Service: 578–579.
- Rehder A. 1940. Manual of cultivated trees and shrubs. 2nd ed. New York: Macmillan. 996 p.
- Schechter MS. 1943. The insecticidal principle in the fruit of Amur corktree (*Phellodendron amurense*). Journal of Organic Chemistry 8: 194–197.
- Soludukhin ED. 1977. Features of the regeneration of *Phellodendron amurense* [abstract]. Lesnoe Khozyaistvo 4: 32–33.
- Starshova NP. 1972. The 'anthecology' of *Phellodendron amurense* in the middle Volga region: 1. Ecology of flowering. Botanicheski Zhurnal 57(11): 1402–1412.
- Swingle CF, comp. 1939. Seed propagation of trees, shrubs, and forbs for conservation planting. SCS-TP-27, Washington, DC: USDA Soil Conservation Service. 198 p.
- Timm GT. 1989. Effect of stratification on the field germination rate of some tree seed. Journal of the Korean Forestry Society 78(1): 26–29.
- Yerkes GE. 1945. Propagation of trees and shrubs. Farm. Bull. 1567 (rev.) Washington, DC: USDA. 54 p.