

Rosaceae—Rose family
***Purshia* DC. ex Poir.**
 bitterbrush, cliffrose

D. Terrance Booth, Susan E. Meyer, and Nancy L. Shaw

Dr. Booth is a rangeland scientist at the USDA Agricultural Research Service's High Plains Grasslands Research Station, Cheyenne, Wyoming; Dr. Meyer is a research ecologist at the USDA Forest Service's Rocky Mountain Research Station, Shrub Sciences Laboratory, Provo, Utah; Dr. Shaw is a research botanist at the USDA Forest Service's Rocky Mountain Research Station, Forestry Sciences Laboratory, Boise, Idaho

Growth habit, occurrence, and use. The bitterbrush genus—*Purshia*—as presently circumscribed comprises 8 species of decumbent to arborescent shrubs of interior western North America. Three are common in the United States (table 1). The type species—antelope bitterbrush—has an essentially northern distribution, whereas cliffrose has an essentially southern distribution, and desert bitterbrush occurs in parts of the geographic area where the other 2 species have overlapping distributions. Cliffrose, along with the 5 Mexican species of the genus, has been traditionally referred to the genus *Cowania* D. Don. Cliffrose regularly forms hybrids with antelope bitterbrush, and desert bitterbrush could be interpreted as a stabilized hybrid between these species (Stutz and Thomas 1964). In fact, molecular genetics work by Jabbes (2000) indicates that *Purshia* was derived from *Cowania*. We follow Welsh and others (1987) in treating the group as congeneric under the name *Purshia*.

Members of the genus are erect, spreading or decumbent, freely branched shrubs up to 6 m in height. They have small, alternate, simple, apically lobed leaves that may be evergreen (cliffrose) to winter deciduous (antelope bitterbrush). Layering forms of bitterbrush (principally antelope bitterbrush) may resprout after fire, but erect forms are usually not fire tolerant. Because of their interesting habits, attractive foliage, and showy flowers, bitterbrush species

have potential as ornamentals in low-maintenance landscapes.

Bitterbrush species are hardy and drought tolerant. Antelope bitterbrush occurs mainly on well-drained soils over a wide elevational range and is often a principal component of mixed shrub, pinyon–juniper, ponderosa pine, and sometimes lodgepole pine communities, where it is notable as a nurse plant for conifer seedlings (Geier-Hayes 1987; McArthur and others 1983; Nord 1965; Tew 1983). It is valued as a high-protein browse for domestic and wild ungulates, being especially important on winter ranges (Bishop and others 2001; Scholten 1983). It also supplies high-quality forage during spring and summer months (Austin and Urness 1983; Ngugi and others 1992). Cliffrose grows primarily on rocky sites in blackbrush–joshua tree woodland, sagebrush–grassland, piñon–juniper woodland, mountain brush, and ponderosa pine communities, sometimes forming extensive stands on south-facing ridge slopes (McArthur and others 1983). It is also an important browse species, especially for mule deer (*Odocoileus hemionus*) (Plummer and others 1968). Desert bitterbrush is a component of blackbrush, chaparral, and piñon–juniper communities.

The bitterbrush species form actinorhizal root nodules that fix nitrogen when soil water is adequate (Bond 1976;

Table 1—*Purshia*, bitterbrush, cliffrose: common names and geographic distributions

Scientific name & synonym(s)	Common name	Geographic distribution
<i>P. glandulosa</i> Curran <i>P. tridentata</i> var. <i>glandulosa</i> (Curran) M.E. Jones	desert bitterbrush	SW Utah, S Nevada, & S California
<i>P. mexicana</i> (D. Don) Henrickson <i>Cowania mexicana</i> D. Don	cliffrose	S Colorado W through Utah to S California & S to New Mexico, Arizona, Sonora, & Chihuahua
<i>P. tridentata</i> (Pursh) DC.	antelope bitterbrush	British Columbia to W Montana, S to New Mexico, California, & N Arizona

Sources: Little (1979), Sargent (1965), Vines (1960).

Kyle and Righetti 1996; Nelson 1983; Righetti and others 1983). They readily function as pioneer species that colonize harsh, steep disturbances and have been used extensively in revegetation and disturbed-land reclamation. An ethanol extract of antelope bitterbrush aerial stems was found to inhibit reverse transcriptase of HIV-1 and to contain the cyanoglucosides pushianin and menisdaurin (Nakanishi and others 1994). Unfortunately, the cyanoglucosides lacked the inhibitory activity of the original extract. Cliffrose has also been examined for beneficial secondary products (Hideyuki and others 1995; Ito and others 1999). Specific populations of antelope bitterbrush with distinctive attributes are recognized and are commercially harvested and sold, although to date only two ('Lassen' and 'Maybell') have been formally named (Davis and others 2002; Shaw and Mosen 1995).

Flowering and fruiting. Most of the medium to large, perfect, cream to sulfur yellow flowers of this genus appear during the first flush of flowering in April, May, or June, depending on elevation. In areas where they co-occur, antelope bitterbrush usually flowers 2 to 3 weeks before cliffrose. The flowers are borne on lateral spurs of the previous year's wood (Shaw and Mosen 1983). In cliffrose, summer rains may induce later flowering on current-year leaders, but these flowers rarely set good seeds (Alexander and others 1974). The flowers have a sweet fragrance and are primarily insect-pollinated. Each has 5 sepals, 5 separate petals, numerous stamens, and 1 to 10 pistils borne within a hypanthium. Flowers of antelope and desert bitterbrushes usually contain a single pistil with a relatively short, non-plumose style, whereas those of cliffrose contain multiple pistils. The pistils develop into single-seeded achenes with papery pericarps. In cliffrose the achenes are tipped with persistent) plumose styles, 22 to 50 mm (1 to 2 in) in length, that give the plants a feathery appearance in fruit.

The main fruit crop ripens from June through August, depending on species and elevation. Plants begin to bear seeds as early as 5 years of age. At least some fruits are produced in most years, and abundant seedcrops are produced on average every 2 to 3 years (Alexander and others 1974; Deitschman and others 1974). Cliffrose seeds (figure 1) are apparently dispersed principally by wind (Alexander and others 1974). Scatter-hoarding rodents such as chipmunks (*Tamias* spp.), disperse bitterbrush seeds (figure 2) and seedlings from rodent caches appear to account for nearly all (99%) natural recruitment as survivors from seedling clumps containing 2 to >100 individuals (Evans and others 1983; Vander Wall 1994).

Seed collection, cleaning, and storage. Bitterbrush plants produce more leader growth in favorable water years,

Figure 1—*Purshia*, mexicana, cliffrose: achenes:





Figure 2—*Purshia*, bitterbrush: achenes (left) and cleaned seeds (right) of *P. glandulosa*, desert bitterbrush (top) and



and leader length is an indicator of the potential for seed production the following year (McCarty and Price 1942; Young and Young 1986). Fruits may be hand-stripped or beaten into hoppers or other containers when fully ripe; harvesters should take care to protect themselves from the fiberglass-like style hairs in the case of cliffrose. The window of opportunity is quite narrow, as ripe fruits are easily detached by wind and do not persist long on the plant, making close monitoring during ripening advisable. Plants in draws and other areas protected from wind may retain their seeds longer. Maturation dates for antelope bitterbrush have been predicted with reasonable accuracy using elevational and latitudinal predictors (Nord 1965). Well-timed harvests of antelope bitterbrush average 168 to 224 kg/ha (150 to 200 lb/acre) but may range up to 560 kg/ha (500 lb/acre) (Nord 1965). Fill percentages are usually high, although insects or drought stress during filling can damage the crop (Shaw and Mosen 1983). Krannitz (1997a) reported the variation in seed weight from 240 bitterbrush plants representing 10 sites in the southern Okanagan Valley of Canada varied from

5 to 46 mg/seed with the population being skewed toward the small seeds. The representative weights given in table 2 are of cleaned seeds (the smaller fraction is removed in cleaning). Krannitz also found that larger seeds had greater concentrations of nitrogen than smaller seeds and that shrubs that had been browsed most intensively the winter before seed-set had seeds with greater concentrations of magnesium (Krannitz 1997b).

A seed cleaner or barley de-bearder may be used to break the styles from cliffrose achenes and to remove the papery pericarps of bitterbrush species. The achenes (cliffrose) or seeds (bitterbrush species) may be separated from the inert material—which usually comprises from one-third (antelope bitterbrush) to two-thirds (cliffrose) of the total weight—using a fanning mill (Alexander and others 1974; Giunta and others 1978). In cliffrose, the achene is considered the seed unit, as the seed is held tightly within the pericarp and cannot be threshed out without damage. In bitterbrush species, the seeds are easily threshed free of their papery pericarps, and the seed unit is the seed itself. If properly dried (<10% moisture content), seeds of bitterbrush species can be warehouse-stored for 5 to 7 years (Belcher 1985) or even up to 15 years without losing viability (Stevens and others 1981).

Germination and seed testing. Bitterbrush and cliffrose seeds are mostly dormant but the inhibiting mechanism(s) is not understood (Booth 1999; Booth and Sowa 2001; Dreyer and Trousdale 1978; Meyer 1989; Meyer and Monsen 1989; Young and Evans 1976, 1981). Moist chilling is preferred for breaking dormancy (table 3). Although some collections are less dormant than others are—as indicated by germination percentages for untreated or partially treated seeds (table 3) (Booth 1999; Meyer and Monsen 1989)—there is no obvious relationship between collection site and chilling requirement (Meyer and Monsen 1989). Dormancy might be affected by high seed temperature (30 °C) while in the dry state (Meyer 1989) and is certainly affected by imbibition temperature (Booth 1999; Meyer 1989).

Young and Evans (1981) reported the required chilling period was shorter at 5 °C, than at 2 °C for all 3 species, and that adequately chilled seeds could germinate over a wide range of temperatures. A 28- to 30-day chill at 1 to 3 °C is highly recommended (AOSA 1993; Belcher 1985; Booth 1999; Meyer 1989) followed by post-chill incubation at 15 °C (10/20 °C for cliffrose). Desert bitterbrush needs only 14 days of chilling (Belcher 1985). Germination of antelope bitterbrush seeds can be facilitated by 24 hours of soaking in cold (2 °C) water prior to moist chilling, but soaking in

Table 2—*Purshia*, bitterbrush and cliffrose: seed yield data (seeds/weight) for mechanically cleaned seeds*

Species	Mean		Range	
	/kg	/b	/kg	/b
<i>P. glandulosa</i>	50,850	26,540	45,000–90,000	20,300–40,900
<i>P. mexicana</i>	129,000	58,600	108,000–210,000	49,000–95,000
<i>P. tridentata</i>	35,000	15,750	29,000–51,000	13,400–23,200

Sources: Alexander and others (1974), Belcher (1985), Deitschman and others (1974), Meyer (2002), Meyer and others (1988).

Table 3—*Purshia*, bitterbrush and cliffrose: germination data

Species	Mean percentage of initially viable seeds							Samples
	0	2 wk	4 wk	6 wk	8 wk	10 wk	12 wk	
<i>P. glandulosa</i>	—	—	—	93	—	—	100	1*
	10	56	81	100	65	—	32	1†
<i>P. mexicana</i>	6	33	83	94	100	—	—	6
	6	64	91	100	32	—	19	1†
<i>P. tridentata</i>	2	43	88	98	100	—	—	13
	13	60	100	100	36	—	37	1†

Sources: Deitschman and others (1974), Meyer (2002), Meyer and Monsen (1989), Young and Evans (1981).

Note: Values are expressed as percentage of initially viable seeds after moist chilling at to 2 °C for 0 to 12 weeks followed by incubation at 15 °C or 10/20 °C for 4 weeks.

* These seeds were chilled at 3 to 5 °C and germination was scored during chilling.

† Decrease in germination percentage after 6 weeks was due to seed mortality during the test.

warm water (>10 °C), or holding imbibed seeds at warm temperatures, decreases seedling vigor and increases pre-germination seed-weight loss (Booth 1999; Booth and Sowa 2001). Longer, colder chilling periods (28 days, 2 °C vs 14 days, 5 °C) increases seedling vigor (Booth 1999; Booth and Morgan 1993). Recommended germination test periods are 28 days for antelope bitterbrush and cliffrose (AOSA 1993).


Soaking seeds in hydrogen peroxide (Everett and Meeuwig 1975) or a 1 to 3% solution of thiourea (Pearson 1957; Young and Evans 1981) will induce germination but these methods have not proven useful for field plantings. Booth (1999) found thiourea-treated seeds to have the lowest seedling vigor among 8 dormancy-breaking treatments and attributed the lower vigor to residual dormancy and to weight loss resulting from accelerated respiration (Booth 1999; Booth and Sowa 2001).

Tetrazolium (TZ) staining is acceptable for evaluating seed quality of bitterbrush (AOSA 1993; Weber and Weisner 1980). Meyer (2002) found no significant difference between TZ viability estimates and germination percentages after 8 weeks of chilling for either cliffrose or antelope bitterbrush. For TZ viability testing, seeds should be clipped at the cotyledon end (figure 3) and soaked in water for 6 to 24 hours. Then, the embryos can be popped-out of the cut end by gentle finger pressure and immersed in 1% TZ solution for 4 to 12 hours at room temperature before evaluation. Cliffrose must be soaked longer than bitterbrush before the embryos can be popped out.

Field seeding and nursery practice. Bitterbrush species are generally sown in fall or early winter in a mixture with other shrubs and forbs. They are used in upper sagebrush, piñon-juniper woodlands, and mountain brush vegetation types to improve degraded wildlife habitat or revegetate bare roadcuts, gullies, south slopes, and other difficult sites (Alexander and others 1974). Because of the chilling requirement, spring-seeding should be avoided. Seeds may be drilled at a depth of 6 to 12 mm ($1/4$ to $1/2$ in) or deeper. Deeper seeding may provide some protection from rodent depredation, which can be a serious problem (Alexander and others 1974; Evans and others 1983; Vander Wall 1994). Seeding in late fall or early winter, when rodents are less active, may also alleviate this problem.

Broadcast-seeding is generally unsuccessful unless provision is made for covering the seeds. Aerial seeding is not recommended. The seedlings do not compete well with weedy annual grasses such as red brome (*Bromus rubens* L.) and cheatgrass (*B. tectorum* L.), or with heavy stands of perennial grasses. They are sensitive to frost and drought during establishment (Plummer and others 1968). Recommended (drill) seeding rates for cliffrose are 5 to 10% of the shrub mix at 8 to 10 kg/ha (7 to 9 lb/ac) (Alexander

Figure 3—*Purshia*: longitudinal section of *P. mexicana*, cliffrose (**top**) and *P. tridentata*, antelope bitterbrush (**bottom**).



and others 1974; Plummer and others 1968) and 16 to 65 seeds/m (5 to 20 seeds/ft) for bitterbrush. The higher rates are advisable for both species when seeding in crust-forming soils. The most effective method of seeding large areas in conjunction with chaining is with a seed dribbler that drops seeds in front of the bulldozers pulling the chain.

Hand-planting into scalped sites with a tool such as a cased-hole punch planter can be very effective on a small scale (Booth 1995). The purpose of scalping is to control herbaceous competition within a half-meter ($1/2$ -ft) radius of the planting spots. Treating seeds with fungicide, planting seeds in groups, and planting with vermiculite to aid in moisture retention have all improved emergence and establishment of antelope bitterbrush (Booth 1980; Evans and

others 1983; Ferguson and Basile 1967). Good emergence depends on adequate snowcover (Young and others 1993).

Bitterbrush species are readily grown as bareroot or container stock, and outplanting may succeed where direct seeding has failed (Alexander and others 1974). Care must be taken to lift or transplant stock only when the plants are hardened or dormant, as survival of actively growing plants is generally low (Landis and Simonich 1984; Shaw 1984). Plants are easier to handle and have higher survival rates if allowed to reach sufficient size before field transplanting. One-year-old bareroot stock or container seedlings 16 to 20

weeks of age are usually large enough (Alexander and others 1974; Shaw 1984). On more level terrain, a conventional tree-planter may be used (Alexander and others 1974). Transplanting should be carried out at a time and in such a way as to assure that the transplants will have adequate moisture for root development for 4 to 6 weeks after planting. This may be accomplished by planting in very early spring or by watering at the time of planting. Fall-planted seedlings may require supplemental watering. Controlling competition from weedy annual or perennial grasses before planting will enhance survival and first-season growth.

References

- Alexander RR, Jorgensen K, Plummer AP. 1974. *Cowania mexicana* var. *stansburiana* (Torr.) Jeps., cliffrose. In: Schopmeyer CS, tech. coord. Seeds of woody plants in the United States. Agric. Handbk. 450. Washington, DC: USDA Forest Service: 353–355.
- AOSA [Association of Official Seed Analysts]. 1993. Rules for testing seeds. *Journal of Seed Technology* 16(3): 1–113.
- Austin DD, Urness PJ. 1983. Summer use of bitterbrush rangelands by mule deer. In: Tiedemann AR, Johnson KL, comps. Proceedings, Research and Management of Bitterbrush and Cliffrose in Western North America; 1982 May 13–15; Salt Lake City. Gen. Tech. Rep. INT-152. Ogden, UT: USDA Forest Service, Intermountain Forest and Range Experiment Station: 203–212.
- Belcher E. 1985. Handbook on seeds of browse-shrubs and forbs. Tech. Pub. R8-8. Atlanta: USDA Forest Service, Southern Region. 246 p.
- Bishop CJ, Garton EO, Unsworth JW. 2001. Bitterbrush and cheatgrass quality on 3 southwest Idaho winter ranges. *Journal of Range Management* 54:595–602.
- Bond G. 1976. Observations on the root nodules of *Purshia tridentata*. *Proceedings of the Royal Society of London (Series B)* 192: 127–135.
- Booth DT. 1980. Emergence of bitterbrush seedlings on land disturbed by phosphate mining. *Journal of Range Management* 33: 439–441.
- Booth DT. 1995. Cased-hole punch seeder: a tool for increasing plant diversity. In: Abstracts, 48th Annual Meeting. Denver: Society for Range Management: 8.
- Booth DT. 1999. Imbibition temperatures affect bitterbrush seed dormancy and seedling vigor. *Journal of Arid Environments* 48: 35–39.
- Booth DT, Morgan DR. 1993. Post-germination growth related to time-to-germination for four woody plants. *Journal of Seed Technology* 16: 30–38.
- Booth DT, Sowa S. 2001. Respiration in dormant and non-dormant bitterbrush seeds. *Journal of Arid Environments* 43: 91–101.
- Davis KM, Englert JM, Kujawski JL. 2002. Improved conservation plant materials released by NRC and cooperators through September 2002. Beltsville, MD: USDA Agricultural Research Service. 57 p.
- Deitschman GH, Jorgensen KR, Plummer AP. 1974. *Purshia*, bitterbrush. In: Schopmeyer CS, tech. coord. Seeds of woody plants in the United States. Agric. Handbk. 450. Washington DC: USDA Forest Service: 686–688.
- Dreyer DL and Trousdale EK. 1978. Cucurbitacins in *Purshia tridentata*. *Phytochemistry* 17:325–326.
- Evans RA, Young JA, Cluff GJ, McAdoo JK. 1983. Dynamics of antelope bitterbrush seed caches. In: Tiedemann AR, Johnson KL, comps. Proceedings, Research and Management of Bitterbrush and Cliffrose in Western North America. 1982 May 13–15; Salt Lake City. Gen. Tech. Rep. INT-152. Ogden, UT: USDA Forest Service, Intermountain Forest and Range Experiment Station: 195–202.
- Everett RL, Meeuwig RO. 1975. Hydrogen peroxide and thiourea treatment of bitterbrush seed. Note 196. Ogden, UT: USDA Forest Service, Intermountain Forest and Range Experiment Station. 6 p.
- Ferguson RB, Basile JV. 1967. Effect of seedling numbers on bitterbrush survival. *Journal of Range Management* 20: 380–382.
- Geier-Hayes K. 1987. Occurrence of conifer seedlings and their microenvironments on disturbed sites in central Idaho. Res. Pap. INT-383. Ogden, UT: USDA Forest Service, Intermountain Forest and Range Experiment Station. 12 p.
- Giunta B, Stevens R, Jorgensen K, Plummer AP. 1978. Antelope bitterbrush: an important wildland shrub. Pub. 78-12. Salt Lake City: Utah State Division of Wildlife Resources. 48 p.
- Hideyuki I, Tsutomu H, Masateru M, Takashi Y, Mutsuo K, Harukuni T, Takuo O. 1995. Tannins from *Cowania mexicana*, and their inhibitory effects on TPA-induced EBV activation. In: Brouillard R, Jay M, Scalbert A, eds. Polyphenols 94, Paris: Institut National de la Recherche Agronomique (INRA): 419–420.
- Ito H, Miyake M, Nishitani E, Mori K, Hatano T, Okuda T, Konoshima T, Takasaki M, Kozuka M, Mukainaka T, Tokuda H, Nishino H, Yoshida T. 1999. Anti-tumor promoting activity of polyphenols from *Cowania mexicana* and *Coleogyne ramosissima*. *Cancer Letters* 143: 5–13.
- Jabbes M. 2000. Hybridization and its evolutionary consequences in *Purshia* and *Cowania* [dissertation]. Moscow, ID: University of Idaho. 185 p.
- Krannitz PG. 1997a. Variation in magnesium and nitrogen content in seeds of antelope bitterbrush (*Purshia tridentata* (Rosaceae)). *American Midland Naturalist* 138: 306–321.
- Krannitz PG. 1997b. Seed weight variability of antelope bitterbrush (*Purshia tridentata*: Rosaceae). *American Journal of Botany* 84: 1738–1742.
- Kyle NE, Righetti TL. 1996. Observations of shoots and roots from interspecific grafted rosaceous shrubs. *Journal of Range Management* 49: 350–354.
- Landis TD, Simonich EJ. 1984. Producing native plants as container seedlings. In: Murphy PM, ed. The challenge of producing native plants for the Intermountain area. Gen. Tech. Rep. INT-168. Ogden, UT: USDA Forest Service, Intermountain Forest and Range Experiment Station: 16–25.
- Little EL Jr. 1979. Checklist of United States trees (native and naturalized). Agric. Handbk. 541. Washington, DC: USDA Forest Service. 375 p.
- McArthur ED, Stutz HC, Sanderson SC. 1983. Taxonomy, distribution, and cytogenetics of *Purshia*, *Cowania*, and *Fallugia* (Rosaceae, Rosaceae). In: Tiedemann AR, Johnson KL, comps. Proceedings, Research and Management of Bitterbrush and Cliffrose in Western North America; 1982 April 13–15; Salt Lake City. Gen. Tech. Rep. INT-152. Ogden, UT: USDA Forest Service, Intermountain Forest and Range Experiment Station: 4–24.

- McCarty EC, Price R. 1942. Growth and carbohydrate content of important mountain forage plants in central Utah as affected by clipping and grazing. *Tech. Bull.* 818. Washington, DC: USDA. 51 p.
- Meyer SE. 1989. Warm pretreatment effects on antelope bitterbrush (*Purshia tridentata*) germination response to chilling. *Northwest Science* 63: 146–153.
- Meyer SE. 2002. Unpublished data. Provo, UT: USDA Forest Service, Intermountain Research Station.
- Meyer SE, Monsen SB. 1989. Seed germination biology of antelope bitterbrush (*Purshia tridentata*). In: Wallace A, McArthur ED, Haferkamp MR, comps. *Proceedings, Symposium on Shrub Ecophysiology and Biotechnology*; 1987 June 30–July 2; Logan, UT. *Gen. Tech. Rep. INT-256*. Ogden, UT: USDA Forest Service, Intermountain Forest and Range Experiment Station: 147–157.
- Meyer SE, Kitchen SG, Wilson GR, Stevens R. 1988. Proposed rule for *Cowania mexicana*—cliffrose. *Association of Official Seed Analysts Newsletter* 63: 24–25.
- Nakanishi T, Nishi M, Somekawa M, Murata H, Mizuno M, Inuma M, Tanaka T, Murata J. 1994. Structures of new and known cyanoglucosides from a North American plant, *Purshia tridentata* DC. *Chemical Pharmaceutical Bulletin* 42:2251–2255.
- Nelson DL. 1983. The occurrence and nature of *Actinorhizae* on *Cowania stansburiana* and other Rosaceae. In: Tiedemann AR, Johnson KL, comps. *Proceedings, Research and Management of Bitterbrush and Cliffrose in Western North America*; 1982 April 13–15; Salt Lake City. *Gen. Tech. Rep. INT-152*. Ogden, UT: USDA Forest Service, Intermountain Forest and Range Experiment Station: 225–239.
- Ngugi KR, Powell J, Hinds FC, Olsen RA. 1992. Range animal diet composition in southcentral Wyoming. *Journal of Range Management* 45: 542–545.
- Nord EC. 1965. Autecology of bitterbrush in California. *Ecological Monographs* 35: 307–334.
- Pearson BO. 1957. Bitterbrush seed dormancy broken with thiourea. *Journal of Range Management* 10: 41–42.
- Plummer AP, Christensen DR, Monsen SB. 1968. Restoring big game range in Utah. *Pub. 68-3*. Salt Lake City: Utah Division of Fish and Game. 183 p.
- Righetti TL, Chard CH, Munns DN. 1983. Opportunities and approaches for enhancing nitrogen fixation in *Purshia*, *Cowania* and *Fallugia*. In: Tiedemann AR, Johnson KL, comps. *Proceedings, Research and Management of Bitterbrush and Cliffrose in Western North America*; 1982 April 13–15; Salt Lake City. *Gen. Tech. Rep. INT-152*. Ogden, UT: USDA Forest Service, Intermountain Forest and Range Experiment Station: 214–224.
- Sargent CS. 1965. *Manual of the trees of North America (exclusive of Mexico)*. 2 volumes; 2nd corr. ed. New York: Dover. 934 p.
- Scholten GC. 1983. Bitterbrush management on the Boise wildlife management area. In: Tiedemann AR, Johnson KL, comps. *Proceedings, Research and Management of Bitterbrush and Cliffrose in Western North America*; 1982 April 13–15; Salt Lake City. *Gen. Tech. Rep. INT-152*. Ogden, UT: USDA Forest Service, Intermountain Forest and Range Experiment Station: 153–162.
- Shaw N. 1984. Producing bareroot seedlings of native shrubs. In: Murphy PM, ed. *The challenge of producing native plants for the intermountain area*. *Gen. Tech. Rep. INT-168*. Ogden, UT: Forest Service, Intermountain Forest and Range Experiment Station: 6–15.
- Shaw NL, Monsen SB. 1983. Phenology and growth habits of nine antelope bitterbrush, desert bitterbrush, stansbury cliffrose, and apache plume accessions. In: Tiedemann AR, Johnson KL, comps. *Proceedings, Research and Management of Bitterbrush and Cliffrose in Western North America*; 1982 April 13–15; Salt Lake City. *Gen. Tech. Rep. INT-152*. Ogden, UT: USDA Forest Service, Intermountain Forest and Range Experiment Station: 55–69.
- Shaw NL, Monsen SB. 1995. "Lassen" antelope bitterbrush. In: Roundy BA, comp. *Proceedings, Wildland Shrub and Arid Land Restoration Symposium*; 1993 October 19–21; Las Vegas, NV. *Gen. Tech. Rep. INT-GTR-315*. Ogden, UT: USDA Forest Service, Intermountain Research Station: 364–371.
- Stevens R, Jorgensen KR, Davis JN. 1981. Viability of seed from thirty-two shrub and forb species through fifteen years of warehouse storage. *Great Basin Naturalist* 41: 274–277.
- Stutz HC, Thomas KL. 1964. Hybridization and introgression in *Cowania* and *Purshia*. *Evolution* 18: 183–185.
- Tew RK. 1983. Bitterbrush distribution and habitat classification on the Boise National Forest. In: Tiedemann AR, Johnson KL, comps. *Proceedings, Research and Management of Bitterbrush and Cliffrose in Western North America*; 1982 April 13–15; Salt Lake City. *Gen. Tech. Rep. INT-152*. Ogden, UT: USDA Forest Service, Intermountain Forest and Range Experiment Station: 32–36.
- Vander Wall SB. 1994. Seed fate pathways of antelope bitterbrush: dispersal by seed-caching yellow pine chipmunks. *Ecology* 75: 1911–1926.
- Vines RA. 1960. *Trees, shrubs, and woody vines of the Southwest*. Austin: University of Texas Press. 1104 p.
- Weber GP, Weisner LE. 1980. Tetrazolium testing procedures for native shrubs and forbs. *Journal of Seed Technology* 5: 23–34.
- Welsh SL, Atwood ND, Higgins LC, Goodrich S. 1987. *A Utah flora*. Great Basin Naturalist Mem. 9. Provo, Utah: Brigham Young University Press. 894 p.
- Young JA, Evans RA. 1976. Stratification of bitterbrush seeds. *Journal of Range Management* 29: 421–425.
- Young JA, Evans RA. 1981. Germination of seeds of antelope bitterbrush, desert bitterbrush, and cliffrose. Washington, DC: USDA Science and Education Administration. *Agric. Res. Results W-17/January* 1981. 39 p.
- Young JA, Young CG. 1986. Collecting, processing, and germinating seeds of wildland plants. Portland, OR: Timber Press. 236 p.
- Young JA, Wight JR, Mowbray JE. 1993. Field stratification of antelope bitterbrush seeds. *Journal of Range Management* 46: 325–330.